



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 03:41 PM BST

PDB ID : 6EIK
Title : A de novo designed heptameric coiled coil CC-Hept-I24E
Authors : Rhys, G.G.; Burton, A.J.; Dawson, W.M.; Thomas, F.; Woolfson, D.N.
Deposited on : 2017-09-19
Resolution : 1.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4009 (1.54-1.50)
Ramachandran outliers	138981	4148 (1.54-1.50)
Sidechain outliers	138945	4146 (1.54-1.50)
RSRZ outliers	127900	3943 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	32	<div> <div>16%</div> <div> <div></div> <div>66%</div> <div>22%</div> <div>6%</div> <div>6%</div> </div> </div>
1	B	32	<div> <div>3%</div> <div> <div></div> <div>78%</div> <div>16%</div> <div>6%</div> </div> </div>
1	C	32	<div> <div>13%</div> <div> <div></div> <div>75%</div> <div>19%</div> <div>6%</div> </div> </div>
1	D	32	<div> <div>9%</div> <div> <div></div> <div>81%</div> <div>13%</div> <div>6%</div> </div> </div>
1	E	32	<div> <div>16%</div> <div> <div></div> <div>69%</div> <div>19%</div> <div>13%</div> </div> </div>
1	F	32	<div> <div></div> <div> <div></div> <div>78%</div> <div>16%</div> <div>6%</div> </div> </div>
1	G	32	<div> <div>6%</div> <div> <div></div> <div>84%</div> <div>6%</div> <div>6%</div> </div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 1826 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CC-Hept-I24E.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	30	Total	C	N	O	8	1	0
			230	146	43	41			
1	B	30	Total	C	N	O	4	1	0
			234	149	45	40			
1	C	30	Total	C	N	O	6	2	0
			236	150	43	43			
1	D	30	Total	C	N	O	4	0	0
			229	147	42	40			
1	E	28	Total	C	N	O	4	3	0
			232	148	45	39			
1	F	30	Total	C	N	O	0	5	0
			256	163	48	45			
1	G	30	Total	C	N	O	3	2	0
			243	155	47	41			

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	1
			6	2	4		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	16	Total	O	0	0
			16	16		
4	B	25	Total	O	0	0
			25	25		
4	C	15	Total	O	0	0
			15	15		
4	D	18	Total	O	0	0
			18	18		
4	E	17	Total	O	0	0
			17	17		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	F	21	Total	O	0	0
			21	21		
4	G	24	Total	O	0	0
			24	24		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CC-Hept-I24E




- Molecule 1: CC-Hept-I24E




- Molecule 1: CC-Hept-I24E




- Molecule 1: CC-Hept-I24E





- Molecule 1: CC-Hept-I24E



- Molecule 1: CC-Hept-I24E



● Molecule 1: CC-Hept-I24E



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 ₂ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	62.21 Å 62.21 Å 108.68 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.78 – 1.52 40.78 – 1.52	Depositor EDS
% Data completeness (in resolution range)	99.1 (40.78-1.52) 99.1 (40.78-1.52)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.99 (at 1.52 Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.161 , 0.208 0.163 , 0.209	Depositor DCC
R_{free} test set	1635 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	24.0	Xtriage
Anisotropy	0.523	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 68.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.38$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	1826	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 23.68 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.4478e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FMT, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.91	7/229 (3.1%)	1.21	2/305 (0.7%)
1	B	2.37	8/236 (3.4%)	1.50	5/314 (1.6%)
1	C	1.71	3/237 (1.3%)	1.28	3/315 (1.0%)
1	D	1.70	3/228 (1.3%)	1.28	1/304 (0.3%)
1	E	1.97	7/237 (3.0%)	1.80	7/314 (2.2%)
1	F	1.84	7/259 (2.7%)	1.27	2/344 (0.6%)
1	G	1.55	2/244 (0.8%)	1.28	3/323 (0.9%)
All	All	1.88	37/1670 (2.2%)	1.39	23/2219 (1.0%)

All (37) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	2	GLU	CD-OE2	-17.80	1.06	1.25
1	A	23	GLU	CG-CD	-11.53	1.34	1.51
1	E	16	GLU	CD-OE1	-11.39	1.13	1.25
1	B	2	GLU	CD-OE1	10.88	1.37	1.25
1	B	9	GLU	CD-OE2	10.77	1.37	1.25
1	B	9	GLU	CD-OE1	9.07	1.35	1.25
1	E	16	GLU	CG-CD	8.76	1.65	1.51
1	A	2	GLU	CD-OE2	8.59	1.35	1.25
1	E	2	GLU	CD-OE2	8.26	1.34	1.25
1	F	16	GLU	CD-OE1	-8.23	1.16	1.25
1	C	29	ARG	N-CA	-8.09	1.30	1.46
1	G	24	GLU	CG-CD	7.73	1.63	1.51
1	D	2	GLU	CD-OE1	7.64	1.34	1.25
1	F	9	GLU	CD-OE1	-7.52	1.17	1.25
1	E	22[A]	ARG	CZ-NH2	-7.17	1.23	1.33
1	E	22[B]	ARG	CZ-NH2	-7.17	1.23	1.33
1	D	24	GLU	CG-CD	6.99	1.62	1.51
1	A	2	GLU	CG-CD	6.96	1.62	1.51
1	B	9	GLU	CG-CD	6.72	1.62	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	24	GLU	CD-OE2	-6.65	1.18	1.25
1	G	2	GLU	CB-CG	-6.62	1.39	1.52
1	D	26	LYS	CB-CG	-6.51	1.34	1.52
1	E	26	LYS	CB-CG	-6.49	1.35	1.52
1	B	29	ARG	N-CA	-6.36	1.33	1.46
1	F	9	GLU	CD-OE2	6.28	1.32	1.25
1	A	0	ACE	C-N	6.01	1.43	1.33
1	B	22	ARG	CZ-NH2	5.86	1.40	1.33
1	A	12	LYS	CE-NZ	5.76	1.63	1.49
1	E	8	ARG	NE-CZ	5.35	1.40	1.33
1	A	16	GLU	CB-CG	-5.34	1.42	1.52
1	F	9	GLU	CG-CD	5.28	1.59	1.51
1	F	1[A]	GLY	N-CA	5.20	1.53	1.46
1	F	1[B]	GLY	N-CA	5.20	1.53	1.46
1	C	2[A]	GLU	CD-OE2	5.13	1.31	1.25
1	C	2[B]	GLU	CD-OE2	5.13	1.31	1.25
1	B	2	GLU	CG-CD	5.07	1.59	1.51
1	A	23	GLU	CD-OE2	5.05	1.31	1.25

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	22[A]	ARG	NE-CZ-NH2	-11.64	114.48	120.30
1	E	22[B]	ARG	NE-CZ-NH2	-11.64	114.48	120.30
1	B	29	ARG	N-CA-C	10.16	138.43	111.00
1	E	22[A]	ARG	NE-CZ-NH1	9.46	125.03	120.30
1	E	22[B]	ARG	NE-CZ-NH1	9.46	125.03	120.30
1	E	0	ACE	O-C-N	9.46	139.28	123.20
1	G	29	ARG	NE-CZ-NH1	-8.62	115.99	120.30
1	A	8	ARG	NE-CZ-NH1	7.58	124.09	120.30
1	F	8[A]	ARG	NE-CZ-NH1	7.38	123.99	120.30
1	F	8[B]	ARG	NE-CZ-NH1	7.38	123.99	120.30
1	C	15	ARG	NE-CZ-NH1	6.93	123.77	120.30
1	D	8	ARG	NE-CZ-NH1	6.68	123.64	120.30
1	B	8[A]	ARG	NE-CZ-NH1	6.32	123.46	120.30
1	B	8[B]	ARG	NE-CZ-NH1	6.32	123.46	120.30
1	C	28	LEU	C-N-CA	6.01	136.73	121.70
1	B	8[A]	ARG	NE-CZ-NH2	-5.93	117.33	120.30
1	B	8[B]	ARG	NE-CZ-NH2	-5.93	117.33	120.30
1	E	8	ARG	NE-CZ-NH1	5.76	123.18	120.30
1	G	24	GLU	CA-CB-CG	5.71	125.97	113.40
1	C	8	ARG	NE-CZ-NH1	5.67	123.13	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	15	ARG	NE-CZ-NH2	-5.59	117.50	120.30
1	G	24	GLU	OE1-CD-OE2	-5.23	117.02	123.30
1	E	2	GLU	OE1-CD-OE2	-5.01	117.29	123.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	230	0	247	0	0
1	B	234	0	255	0	0
1	C	236	0	251	0	1
1	D	229	0	246	0	0
1	E	232	0	260	0	0
1	F	256	0	279	0	0
1	G	243	0	273	0	0
2	A	6	0	8	0	0
2	B	6	0	8	0	0
2	C	6	0	8	0	1
2	D	6	0	7	0	0
3	C	6	0	4	0	0
4	A	16	0	0	0	0
4	B	25	0	0	0	3
4	C	15	0	0	0	0
4	D	18	0	0	0	0
4	E	17	0	0	0	0
4	F	21	0	0	0	0
4	G	24	0	0	0	1
All	All	1826	0	1846	0	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). Clashscore could not be calculated for this entry.

There are no clashes within the asymmetric unit.

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:225:HOH:O	4:B:225:HOH:O[8_665]	0.77	1.43
2:C:102:GOL:O3	4:B:218:HOH:O[8_665]	1.27	0.93
1:C:19:TRP:CZ2	1:C:19:TRP:CZ2[2_755]	1.84	0.36
4:B:223:HOH:O	4:G:107:HOH:O[5_755]	2.01	0.19

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	29/32 (91%)	28 (97%)	0	1 (3%)	3	0
1	B	29/32 (91%)	28 (97%)	1 (3%)	0	100	100
1	C	30/32 (94%)	30 (100%)	0	0	100	100
1	D	28/32 (88%)	28 (100%)	0	0	100	100
1	E	29/32 (91%)	29 (100%)	0	0	100	100
1	F	32/32 (100%)	32 (100%)	0	0	100	100
1	G	30/32 (94%)	30 (100%)	0	0	100	100
All	All	207/224 (92%)	205 (99%)	1 (0%)	1 (0%)	29	9

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	27	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	19/20 (95%)	16 (84%)	3 (16%)	2	0
1	B	20/20 (100%)	20 (100%)	0	100	100
1	C	20/20 (100%)	20 (100%)	0	100	100
1	D	19/20 (95%)	19 (100%)	0	100	100
1	E	20/20 (100%)	20 (100%)	0	100	100
1	F	22/20 (110%)	22 (100%)	0	100	100
1	G	21/20 (105%)	20 (95%)	1 (5%)	25	4
All	All	141/140 (101%)	137 (97%)	4 (3%)	41	14

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LYS
1	A	12	LYS
1	A	23	GLU
1	G	29	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	FMT	C	101[A]	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	C	101[B]	-	0,2,2	0.00	-	0,1,1	0.00	-
2	GOL	D	101	-	5,5,5	1.23	1 (20%)	5,5,5	1.39	1 (20%)
2	GOL	B	101	-	5,5,5	0.66	0	5,5,5	0.39	0
2	GOL	A	101	-	5,5,5	0.63	0	5,5,5	0.53	0
2	GOL	C	102	-	5,5,5	1.23	1 (20%)	5,5,5	1.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	102	-	-	4/4/4/4	-
2	GOL	B	101	-	-	2/4/4/4	-
2	GOL	A	101	-	-	2/4/4/4	-
2	GOL	D	101	-	-	2/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	101	GOL	O2-C2	-2.55	1.35	1.43
2	C	102	GOL	O3-C3	2.47	1.52	1.42

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	101	GOL	O2-C2-C1	-2.44	98.36	109.12

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	101	GOL	O1-C1-C2-C3
2	B	101	GOL	C1-C2-C3-O3
2	A	101	GOL	O1-C1-C2-C3
2	C	102	GOL	C1-C2-C3-O3
2	C	102	GOL	O2-C2-C3-O3
2	A	101	GOL	O1-C1-C2-O2
2	C	102	GOL	O1-C1-C2-C3
2	B	101	GOL	O2-C2-C3-O3
2	C	102	GOL	O1-C1-C2-O2
2	D	101	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	102	GOL	0	1

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	G	2
1	E	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	G	0:ACE	C	1[A]:GLY	N	3.56
1	E	0:ACE	C	1[B]:GLY	N	1.86
1	G	0:ACE	C	1[B]:GLY	N	1.84

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	29/32 (90%)	0.70	5 (17%) 1 1	22, 33, 85, 97	4 (13%)
1	B	29/32 (90%)	0.04	1 (3%) 45 50	22, 28, 61, 98	2 (6%)
1	C	29/32 (90%)	0.72	4 (13%) 2 2	22, 30, 61, 62	2 (6%)
1	D	29/32 (90%)	0.41	3 (10%) 6 6	22, 30, 59, 70	1 (3%)
1	E	27/32 (84%)	0.72	5 (18%) 1 1	23, 31, 81, 88	1 (3%)
1	F	29/32 (90%)	-0.18	0 100 100	22, 26, 49, 95	0
1	G	29/32 (90%)	0.06	2 (6%) 16 17	21, 28, 48, 58	1 (3%)
All	All	201/224 (89%)	0.35	20 (9%) 7 7	21, 30, 73, 98	11 (5%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	28	LEU	6.9
1	A	25	ALA	6.3
1	A	29	ARG	5.3
1	E	25	ALA	5.1
1	E	19	TRP	4.9
1	A	28	LEU	4.9
1	E	27	ALA	4.9
1	A	19	TRP	4.4
1	C	28	LEU	4.3
1	D	29	ARG	3.6
1	C	27	ALA	3.5
1	C	19	TRP	3.4
1	A	27	ALA	3.1
1	B	28	LEU	3.1
1	G	29	ARG	2.9
1	E	26	LYS	2.6
1	D	27	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	28	LEU	2.2
1	E	1[A]	GLY	2.0
1	C	26	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	A	101	6/6	0.72	0.24	69,74,79,86	0
2	GOL	D	101	6/6	0.82	0.19	48,68,71,73	0
2	GOL	B	101	6/6	0.87	0.15	59,63,76,86	0
3	FMT	C	101[B]	3/3	0.94	0.24	47,47,51,57	3
3	FMT	C	101[A]	3/3	0.94	0.24	33,33,45,63	3
2	GOL	C	102	6/6	0.96	0.20	33,36,70,86	0

6.5 Other polymers [i](#)

There are no such residues in this entry.