



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 28, 2022 – 06:04 PM JST

PDB ID : 7EIW  
Title : Human histidine decarboxylase mutant Y334F reacted with histidine  
Authors : Komori, H.  
Deposited on : 2021-04-01  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.27
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	481	 5% 86% 10% ..
1	B	481	 6% 86% 10% ..

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7650 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Histidine decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	468	Total	C	N	O	S	0	3	0
			3741	2395	638	685	23			
1	B	466	Total	C	N	O	S	0	3	0
			3728	2387	638	680	23			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P19113
A	-2	PRO	-	expression tag	UNP P19113
A	-1	LEU	-	expression tag	UNP P19113
A	0	GLY	-	expression tag	UNP P19113
A	1	SER	-	expression tag	UNP P19113
A	180	SER	CYS	engineered mutation	UNP P19113
A	418	SER	CYS	engineered mutation	UNP P19113
B	-3	GLY	-	expression tag	UNP P19113
B	-2	PRO	-	expression tag	UNP P19113
B	-1	LEU	-	expression tag	UNP P19113
B	0	GLY	-	expression tag	UNP P19113
B	1	SER	-	expression tag	UNP P19113
B	180	SER	CYS	engineered mutation	UNP P19113
B	418	SER	CYS	engineered mutation	UNP P19113

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C<sub>8</sub>H<sub>10</sub>NO<sub>6</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	59	Total	O	0	0
			59	59		
3	B	92	Total	O	0	0
			92	92		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histidine decarboxylase



- Molecule 1: Histidine decarboxylase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.90Å 57.65Å 117.65Å 90.00° 106.30° 90.00°	Depositor
Resolution (Å)	33.22 – 2.10 33.22 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.4 (33.22-2.10) 99.5 (33.22-2.10)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.47 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.212 , 0.252 0.216 , 0.251	Depositor DCC
$R_{free}$ test set	2679 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.1	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 33.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7650	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSX, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.65	0/3837	0.76	0/5201
1	B	0.65	0/3823	0.75	0/5182
All	All	0.65	0/7660	0.76	0/10383

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3741	0	3699	31	0
1	B	3728	0	3695	37	0
2	A	15	0	6	1	0
2	B	15	0	6	2	0
3	A	59	0	0	0	0
3	B	92	0	0	0	0
All	All	7650	0	7406	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:63[B]:ARG:HG3	1:B:63[B]:ARG:HH11	1.31	0.95
1:B:63[B]:ARG:HH11	1:B:63[B]:ARG:CG	1.82	0.92
1:A:330:VAL:HG12	1:B:202:LEU:HD13	1.73	0.71
1:A:91:LEU:HA	1:A:94:MET:CE	2.28	0.64
1:B:238:PRO:HB2	1:B:268:LEU:HD21	1.79	0.63
1:A:91:LEU:HA	1:A:94:MET:HE2	1.80	0.63
1:B:8:ARG:O	1:B:12:ARG:HG3	1.99	0.63
1:B:63[B]:ARG:CG	1:B:63[B]:ARG:NH1	2.50	0.61
1:A:330:VAL:HG12	1:B:202:LEU:CD1	2.31	0.60
1:A:198:GLU:CD	1:B:330:VAL:HG21	2.23	0.58
1:A:91:LEU:HD23	1:A:94:MET:CE	2.32	0.58
1:A:194:HIS:CE1	2:A:1000:PLP:H5A1	2.38	0.58
1:A:91:LEU:HD23	1:A:94:MET:HE1	1.85	0.58
1:B:150:VAL:HG23	1:B:302:ASN:OD1	2.05	0.57
1:B:302:ASN:HB3	1:B:304:SER:OG	2.06	0.55
1:A:82:PRO:O	1:A:305:LYS:HA	2.06	0.55
1:B:232:LYS:NZ	1:B:266:GLU:OE1	2.39	0.54
1:B:302:ASN:HD22	1:B:305:LYS:HE3	1.73	0.54
1:A:119:VAL:O	1:A:122:TRP:HB2	2.08	0.53
1:B:347:MET:SD	1:B:353:LEU:CD1	2.97	0.53
1:A:177:ASP:OD2	1:B:207:LYS:NZ	2.42	0.52
1:B:470:ALA:O	1:B:474:ILE:HD13	2.09	0.52
1:B:347:MET:SD	1:B:353:LEU:HD11	2.51	0.51
1:A:30:VAL:HG21	1:A:76:HIS:HB2	1.92	0.51
1:B:66:MET:N	1:B:67:PRO:CD	2.73	0.51
1:B:82:PRO:O	1:B:305:LYS:HA	2.12	0.49
1:B:63[A]:ARG:NH2	1:B:64:ILE:HD11	2.27	0.49
1:B:38:TYR:OH	1:B:68:GLY:HA3	2.13	0.49
1:A:468:ARG:O	1:A:468:ARG:HD3	2.13	0.48
1:A:6:GLU:O	1:A:10:ARG:HG2	2.14	0.48
1:B:60:ASP:HA	1:B:63[B]:ARG:HB2	1.95	0.48
1:A:321:LYS:HB2	1:A:349:TRP:CH2	2.51	0.46
1:A:354:SER:OG	2:B:1000:PLP:O3P	2.18	0.46
1:A:255:CSX:HG	1:A:258:GLU:HG3	1.81	0.45
1:A:398:PHE:CE2	1:A:413:LEU:HD23	2.51	0.45
1:B:22:LEU:HD12	1:B:87:TRP:CZ3	2.52	0.45
1:A:113:THR:O	1:A:117:MET:HG3	2.17	0.44
1:A:372:LYS:HA	1:A:375:GLN:OE1	2.17	0.44
1:A:392:VAL:HG22	1:A:464:TRP:CZ3	2.52	0.44
1:B:65:ILE:C	1:B:67:PRO:HD2	2.38	0.44
1:A:87:TRP:HB2	1:A:88:PRO:HD3	2.00	0.44

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:160:ALA:HA	1:A:327:THR:OG1	2.17	0.44
1:A:440:ILE:HD11	1:A:445:ILE:HG21	1.98	0.44
1:B:433:LEU:N	1:B:433:LEU:HD23	2.32	0.44
1:A:398:PHE:CE2	1:A:413:LEU:CD2	3.02	0.43
1:B:2:MET:HE1	1:B:7:TYR:HA	2.01	0.43
1:B:63[B]:ARG:NH1	1:B:63[B]:ARG:HG2	2.31	0.43
1:B:127:LEU:HD23	1:B:282:LEU:HD23	2.00	0.43
1:A:353:LEU:HD22	1:B:194:HIS:CE1	2.54	0.42
1:A:199:LYS:HE3	1:B:328:PHE:HB3	2.00	0.42
1:A:311:PHE:CG	1:A:312:ASP:HA	2.55	0.42
1:B:194:HIS:CG	2:B:1000:PLP:C6	3.03	0.42
1:B:279:THR:O	1:B:282:LEU:HG	2.19	0.42
1:B:281:PHE:CE2	1:B:290:LEU:HD13	2.55	0.42
1:A:66:MET:N	1:A:67:PRO:CD	2.82	0.42
1:B:6:GLU:O	1:B:10:ARG:HG3	2.20	0.42
1:B:437:PRO:HB2	1:B:444:LEU:HD11	2.02	0.41
1:B:290:LEU:HD12	1:B:290:LEU:HA	1.92	0.41
1:B:35:GLN:O	1:B:38:TYR:HB3	2.21	0.41
1:B:63[B]:ARG:HG3	1:B:63[B]:ARG:NH1	2.13	0.41
1:A:28:ARG:HD2	1:A:71:HIS:CE1	2.55	0.40
1:A:87:TRP:N	1:A:88:PRO:CD	2.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	466/481 (97%)	445 (96%)	21 (4%)	0	100	100
1	B	464/481 (96%)	453 (98%)	11 (2%)	0	100	100
All	All	930/962 (97%)	898 (97%)	32 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	406/413 (98%)	391 (96%)	15 (4%)	34	35
1	B	405/413 (98%)	394 (97%)	11 (3%)	44	48
All	All	811/826 (98%)	785 (97%)	26 (3%)	39	41

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	93	ASP
1	A	141	GLN
1	A	162	ARG
1	A	172	SER
1	A	179	SER
1	A	226	LYS
1	A	256	LEU
1	A	282	LEU
1	A	291	LYS
1	A	321	LYS
1	A	329	SER
1	A	357	PHE
1	A	440	ILE
1	A	444	LEU
1	A	468	ARG
1	B	2	MET
1	B	10	ARG
1	B	23	SER
1	B	42	GLN
1	B	93	ASP
1	B	162	ARG
1	B	170	LYS
1	B	179	SER
1	B	202	LEU
1	B	357	PHE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	476	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	132	HIS
1	B	216	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CSX	A	255	1	3,6,7	0.68	0	1,6,8	2.31	1 (100%)
1	CSX	B	255	1	3,6,7	0.70	0	1,6,8	3.71	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSX	A	255	1	-	0/1/5/7	-
1	CSX	B	255	1	-	0/1/5/7	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	255	CSX	CA-CB-SG	-3.71	105.25	113.36
1	A	255	CSX	CA-CB-SG	-2.31	108.31	113.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	255	CSX	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PLP	A	1000	1	15,15,16	0.62	0	20,22,23	1.27	1 (5%)
2	PLP	B	1000	1	15,15,16	0.62	0	20,22,23	0.90	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	1000	1	-	5/6/6/8	0/1/1/1
2	PLP	B	1000	1	-	0/6/6/8	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	A	1000	PLP	O4P-C5A-C5	4.67	118.26	109.35
2	B	1000	PLP	O4P-C5A-C5	2.74	114.56	109.35

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1000	PLP	C4-C5-C5A-O4P
2	A	1000	PLP	C6-C5-C5A-O4P
2	A	1000	PLP	C5A-O4P-P-O1P
2	A	1000	PLP	C5A-O4P-P-O2P
2	A	1000	PLP	C5A-O4P-P-O3P

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1000	PLP	1	0
2	B	1000	PLP	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	467/481 (97%)	0.27	22 (4%) 31 37	33, 50, 70, 115	0
1	B	465/481 (96%)	0.24	27 (5%) 23 28	31, 43, 69, 86	0
All	All	932/962 (96%)	0.26	49 (5%) 26 32	31, 48, 70, 115	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	342	VAL	6.1
1	B	54	TRP	5.7
1	A	341	GLY	5.3
1	B	87	TRP	4.7
1	B	43	LEU	4.4
1	A	4	PRO	4.2
1	B	9[A]	GLU	4.0
1	A	2	MET	3.9
1	A	342	VAL	3.9
1	B	58	PHE	3.8
1	B	39	LEU	3.6
1	B	57	ILE	3.3
1	A	343	ALA	3.3
1	A	5	GLU	3.3
1	B	38	TYR	2.9
1	B	322	TYR	2.9
1	A	3	GLU	2.8
1	B	41	ALA	2.8
1	B	42	GLN	2.7
1	B	55	ASP	2.7
1	A	131	GLU	2.7
1	B	15	VAL	2.7
1	B	64	ILE	2.6
1	B	309	VAL	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	303	PRO	2.5
1	A	138	PRO	2.5
1	B	301	PHE	2.5
1	A	277	ALA	2.5
1	B	456	THR	2.4
1	A	54	TRP	2.4
1	B	5	GLU	2.3
1	B	18	ILE	2.3
1	A	96	ALA	2.3
1	A	279	THR	2.2
1	B	304	SER	2.2
1	A	53	SER	2.2
1	B	63[A]	ARG	2.2
1	A	330	VAL	2.2
1	A	378	VAL	2.1
1	A	322	TYR	2.1
1	A	140	SER	2.1
1	B	23	SER	2.1
1	A	303	PRO	2.1
1	B	27	GLU	2.1
1	A	139	SER	2.1
1	A	294	GLU	2.1
1	B	21	TYR	2.1
1	A	477	GLN	2.0
1	B	4	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	CSX	A	255	7/8	0.97	0.09	47,48,51,51	0
1	CSX	B	255	7/8	0.97	0.07	35,36,37,38	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PLP	B	1000	15/16	0.93	0.16	44,54,60,61	0
2	PLP	A	1000	15/16	0.94	0.19	43,51,55,55	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.