



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 04:20 pm BST

PDB ID : 5EJD  
Title : The crystal structure of holo T3CT  
Authors : Zhang, J.R.; Tang, Y.; Zhou, J.H.  
Deposited on : 2015-11-01  
Resolution : 2.49 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	477	<div> <div>0%</div> <div> <div></div> <div>82%</div> <div>8%</div> <div>9%</div> </div> </div>
1	D	477	<div> <div>0%</div> <div> <div></div> <div>84%</div> <div>6%</div> <div>8%</div> </div> </div>
1	F	477	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>6%</div> <div>9%</div> </div> </div>
1	H	477	<div> <div>4%</div> <div> <div></div> <div>84%</div> <div>5%</div> <div>9%</div> </div> </div>
1	J	477	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>7%</div> <div>9%</div> </div> </div>
1	L	477	<div> <div>5%</div> <div> <div></div> <div>83%</div> <div>6%</div> <div>9%</div> </div> </div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	N	477	
1	P	477	
2	A	77	
2	C	77	
2	E	77	
2	G	77	
2	I	77	
2	K	77	
2	M	77	
2	O	77	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	P	501	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 33064 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TqaA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	436	Total	C	N	O	S	0	0	0
			3486	2227	601	645	13			
1	D	437	Total	C	N	O	S	0	0	0
			3493	2232	602	646	13			
1	F	435	Total	C	N	O	S	0	0	0
			3478	2223	600	642	13			
1	H	435	Total	C	N	O	S	0	0	0
			3478	2223	600	642	13			
1	J	436	Total	C	N	O	S	0	0	0
			3486	2227	601	645	13			
1	L	435	Total	C	N	O	S	0	0	0
			3478	2223	600	642	13			
1	N	435	Total	C	N	O	S	0	0	0
			3478	2223	600	642	13			
1	P	436	Total	C	N	O	S	0	0	0
			3486	2227	601	645	13			

- Molecule 2 is a protein called TqaA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	75	Total	C	N	O	S	0	0	0
			554	351	94	107	2			
2	C	74	Total	C	N	O	S	0	0	0
			547	346	93	106	2			
2	E	72	Total	C	N	O	S	0	0	0
			533	337	90	104	2			
2	G	75	Total	C	N	O	S	0	0	0
			554	351	94	107	2			
2	I	72	Total	C	N	O	S	0	0	0
			533	337	90	104	2			
2	K	73	Total	C	N	O	S	0	0	0
			536	340	90	104	2			

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	M	70	Total	C	N	O	S	0	0	0
			517	327	87	101	2			
2	O	74	Total	C	N	O	S	0	0	0
			545	345	92	106	2			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP F1CWE4
C	1	MET	-	initiating methionine	UNP F1CWE4
E	1	MET	-	initiating methionine	UNP F1CWE4
G	1	MET	-	initiating methionine	UNP F1CWE4
I	1	MET	-	initiating methionine	UNP F1CWE4
K	1	MET	-	initiating methionine	UNP F1CWE4
M	1	MET	-	initiating methionine	UNP F1CWE4
O	1	MET	-	initiating methionine	UNP F1CWE4

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	P	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is (R)-3-hydroxy-4-((3-((2-mercaptoethyl)amino)-3-oxopropyl)amino)-2,2-dimethyl-4-oxobutyl hydrogen phosphonate (three-letter code: 5PD) (formula:  $C_{11}H_{23}N_2O_6PS$ ).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	S	0	0
			21	11	2	6	1	1		
4	C	1	Total	C	N	O	P	S	0	0
			21	11	2	6	1	1		
4	E	1	Total	C	N	O	P	S	0	0
			21	11	2	6	1	1		
4	G	1	Total	C	N	O	P	S	0	0
			21	11	2	6	1	1		
4	I	1	Total	C	N	O	P	S	0	0
			21	11	2	6	1	1		
4	K	1	Total	C	N	O	P	S	0	0
			21	11	2	6	1	1		
4	M	1	Total	C	N	O	P	S	0	0
			21	11	2	6	1	1		
4	O	1	Total	C	N	O	P	S	0	0
			21	11	2	6	1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	112	Total	O	0	0
			112	112		
5	A	15	Total	O	0	0
			15	15		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	118	Total 118	O 118	0	0
5	C	8	Total 8	O 8	0	0
5	F	73	Total 73	O 73	0	0
5	E	6	Total 6	O 6	0	0
5	H	69	Total 69	O 69	0	0
5	G	6	Total 6	O 6	0	0
5	J	70	Total 70	O 70	0	0
5	I	12	Total 12	O 12	0	0
5	L	63	Total 63	O 63	0	0
5	K	2	Total 2	O 2	0	0
5	N	55	Total 55	O 55	0	0
5	M	5	Total 5	O 5	0	0
5	P	76	Total 76	O 76	0	0
5	O	6	Total 6	O 6	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: TqaA



#### • Molecule 1: TqaA




#### • Molecule 1: TqaA







- Molecule 1: TqaA




- Molecule 1: TqaA



- Molecule 1: TqaA





- Molecule 1: TqaA







• Molecule 1: TqaA




• Molecule 2: TqaA




• Molecule 2: TqaA




• Molecule 2: TqaA



• Molecule 2: TqaA




• Molecule 2: TqaA








● Molecule 2: TqaA



● Molecule 2: TqaA



● Molecule 2: TqaA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.19Å 204.76Å 126.05Å 90.00° 91.10° 90.00°	Depositor
Resolution (Å)	50.00 – 2.49 34.72 – 2.49	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.00-2.49) 99.3 (34.72-2.49)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.10 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.194 , 0.222 0.198 , 0.226	Depositor DCC
$R_{free}$ test set	9162 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	42.4	Xtriage
Anisotropy	0.171	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 32.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.055 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	33064	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 5PD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	B	0.73	0/3582	0.82	9/4892 (0.2%)
1	D	0.73	0/3590	0.84	11/4904 (0.2%)
1	F	0.68	0/3574	0.80	7/4881 (0.1%)
1	H	0.67	0/3574	0.80	7/4881 (0.1%)
1	J	0.68	0/3582	0.83	9/4892 (0.2%)
1	L	0.68	0/3574	0.82	11/4881 (0.2%)
1	N	0.67	1/3574 (0.0%)	0.83	11/4881 (0.2%)
1	P	0.69	0/3582	0.79	5/4892 (0.1%)
2	A	0.76	0/564	0.81	2/764 (0.3%)
2	C	0.73	0/557	0.81	1/754 (0.1%)
2	E	0.68	0/543	0.75	1/736 (0.1%)
2	G	0.75	0/564	1.04	7/764 (0.9%)
2	I	0.66	0/543	0.87	4/736 (0.5%)
2	K	0.67	0/546	0.96	5/741 (0.7%)
2	M	0.69	0/527	0.93	5/714 (0.7%)
2	O	0.65	0/555	0.80	2/753 (0.3%)
All	All	0.69	1/33031 (0.0%)	0.83	97/45066 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	N	0	1
2	C	0	1
2	K	0	1
All	All	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	N	345	TRP	CB-CG	5.96	1.60	1.50

The worst 5 of 97 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	N	442	ASP	CB-CG-OD1	9.44	126.80	118.30
1	D	314	ARG	NE-CZ-NH2	-9.35	115.62	120.30
1	H	314	ARG	NE-CZ-NH1	9.22	124.91	120.30
1	B	314	ARG	NE-CZ-NH1	9.11	124.86	120.30
1	D	314	ARG	NE-CZ-NH1	9.07	124.84	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	73	SER	Peptide
2	K	74	VAL	Peptide
1	N	481	ARG	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3486	0	3415	23	0
1	D	3493	0	3422	15	0
1	F	3478	0	3411	18	0
1	H	3478	0	3411	16	0
1	J	3486	0	3415	21	0
1	L	3478	0	3411	20	0
1	N	3478	0	3411	16	0
1	P	3486	0	3415	17	0
2	A	554	0	542	6	0
2	C	547	0	533	8	0
2	E	533	0	515	2	0
2	G	554	0	542	8	0
2	I	533	0	515	2	0
2	K	536	0	521	11	0
2	M	517	0	498	10	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	O	545	0	529	2	0
3	B	6	0	8	0	0
3	D	6	0	8	0	0
3	P	6	0	8	0	0
4	A	21	0	0	0	0
4	C	21	0	0	0	0
4	E	21	0	0	1	0
4	G	21	0	0	0	0
4	I	21	0	0	0	0
4	K	21	0	0	1	0
4	M	21	0	0	0	0
4	O	21	0	0	1	0
5	A	15	0	0	0	0
5	B	112	0	0	6	0
5	C	8	0	0	1	0
5	D	118	0	0	4	0
5	E	6	0	0	0	0
5	F	73	0	0	1	0
5	G	6	0	0	0	0
5	H	69	0	0	3	0
5	I	12	0	0	1	0
5	J	70	0	0	1	0
5	K	2	0	0	0	0
5	L	63	0	0	1	0
5	M	5	0	0	0	0
5	N	55	0	0	0	0
5	O	6	0	0	0	0
5	P	76	0	0	1	0
All	All	33064	0	31530	182	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 182 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:265:GLN:N	1:J:265:GLN:OE1	1.89	1.05
1:B:201:THR:HG22	5:B:617:HOH:O	1.58	1.04
1:P:473:LYS:NZ	1:P:484:ASP:OD1	1.97	0.97
1:B:201:THR:CG2	5:B:617:HOH:O	2.09	0.94
1:B:20:GLU:N	1:B:20:GLU:OE1	2.02	0.92

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	430/477 (90%)	421 (98%)	8 (2%)	1 (0%)	47	68
1	D	431/477 (90%)	421 (98%)	9 (2%)	1 (0%)	47	68
1	F	429/477 (90%)	416 (97%)	12 (3%)	1 (0%)	47	68
1	H	429/477 (90%)	419 (98%)	9 (2%)	1 (0%)	47	68
1	J	430/477 (90%)	419 (97%)	10 (2%)	1 (0%)	47	68
1	L	429/477 (90%)	417 (97%)	9 (2%)	3 (1%)	22	39
1	N	429/477 (90%)	418 (97%)	9 (2%)	2 (0%)	29	48
1	P	430/477 (90%)	420 (98%)	9 (2%)	1 (0%)	47	68
2	A	73/77 (95%)	73 (100%)	0	0	100	100
2	C	72/77 (94%)	69 (96%)	2 (3%)	1 (1%)	11	20
2	E	70/77 (91%)	68 (97%)	1 (1%)	1 (1%)	11	20
2	G	73/77 (95%)	69 (94%)	3 (4%)	1 (1%)	11	20
2	I	70/77 (91%)	69 (99%)	1 (1%)	0	100	100
2	K	71/77 (92%)	67 (94%)	1 (1%)	3 (4%)	3	3
2	M	68/77 (88%)	66 (97%)	0	2 (3%)	4	6
2	O	72/77 (94%)	71 (99%)	1 (1%)	0	100	100
All	All	4006/4432 (90%)	3903 (97%)	84 (2%)	19 (0%)	29	48

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	74	VAL
1	F	343	THR
2	E	73	SER
1	N	343	THR

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	L	482	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	394/425 (93%)	379 (96%)	15 (4%)	33	58
1	D	395/425 (93%)	383 (97%)	12 (3%)	41	68
1	F	393/425 (92%)	383 (98%)	10 (2%)	47	73
1	H	393/425 (92%)	382 (97%)	11 (3%)	43	70
1	J	394/425 (93%)	380 (96%)	14 (4%)	35	61
1	L	393/425 (92%)	381 (97%)	12 (3%)	40	67
1	N	393/425 (92%)	379 (96%)	14 (4%)	35	61
1	P	394/425 (93%)	380 (96%)	14 (4%)	35	61
2	A	57/58 (98%)	54 (95%)	3 (5%)	22	43
2	C	56/58 (97%)	54 (96%)	2 (4%)	35	61
2	E	55/58 (95%)	53 (96%)	2 (4%)	35	61
2	G	57/58 (98%)	53 (93%)	4 (7%)	15	29
2	I	55/58 (95%)	53 (96%)	2 (4%)	35	61
2	K	55/58 (95%)	51 (93%)	4 (7%)	14	27
2	M	53/58 (91%)	50 (94%)	3 (6%)	20	39
2	O	56/58 (97%)	51 (91%)	5 (9%)	9	19
All	All	3593/3864 (93%)	3466 (96%)	127 (4%)	36	62

5 of 127 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	G	44	MET
1	J	481	ARG
1	P	363	SER
1	J	64	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	J	174	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	J	60	HIS
1	J	69	HIS
1	L	200	GLN
1	J	34	HIS
1	L	390	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	5PD	O	101	2	13,20,20	0.99	0	18,26,26	2.32	7 (38%)
4	5PD	I	101	2	13,20,20	0.79	0	18,26,26	1.67	3 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	5PD	K	101	2	13,20,20	0.80	0	18,26,26	4.71	9 (50%)
3	GOL	P	501	-	5,5,5	1.07	0	5,5,5	0.83	0
3	GOL	B	501	-	5,5,5	0.68	0	5,5,5	0.64	0
3	GOL	D	501	-	5,5,5	0.77	0	5,5,5	0.49	0
4	5PD	E	101	2	13,20,20	0.93	1 (7%)	18,26,26	5.04	10 (55%)
4	5PD	G	101	2	13,20,20	1.09	1 (7%)	18,26,26	3.28	7 (38%)
4	5PD	A	101	2	13,20,20	1.53	2 (15%)	18,26,26	5.73	9 (50%)
4	5PD	C	101	2	13,20,20	0.77	0	18,26,26	2.24	6 (33%)
4	5PD	M	101	2	13,20,20	1.25	2 (15%)	18,26,26	3.88	10 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	5PD	O	101	2	-	3/24/26/26	-
4	5PD	I	101	2	-	3/24/26/26	-
4	5PD	K	101	2	-	11/24/26/26	-
3	GOL	P	501	-	-	2/4/4/4	-
3	GOL	B	501	-	-	0/4/4/4	-
3	GOL	D	501	-	-	3/4/4/4	-
4	5PD	E	101	2	-	12/24/26/26	-
4	5PD	G	101	2	-	4/24/26/26	-
4	5PD	A	101	2	-	6/24/26/26	-
4	5PD	C	101	2	-	2/24/26/26	-
4	5PD	M	101	2	-	5/24/26/26	-

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	101	5PD	O13-C12	3.47	1.48	1.42
4	M	101	5PD	C15-C14	-3.05	1.47	1.53
4	A	101	5PD	C08-N09	-2.83	1.39	1.46
4	G	101	5PD	C07-C02	-2.35	1.46	1.51
4	M	101	5PD	C16-C14	-2.24	1.48	1.53

The worst 5 of 61 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	101	5PD	C15-C14-C17	-14.92	83.89	108.23
4	K	101	5PD	C15-C14-C17	-12.54	87.77	108.23
4	A	101	5PD	C16-C14-C17	-11.68	89.18	108.23
4	A	101	5PD	C15-C14-C17	-10.23	91.54	108.23
4	E	101	5PD	C15-C14-C12	10.14	126.40	108.82

There are no chirality outliers.

5 of 51 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	O	101	5PD	C02-C07-C08-N09
3	P	501	GOL	O1-C1-C2-C3
4	K	101	5PD	O11-C10-C12-C14
4	K	101	5PD	O11-C10-C12-O13
4	K	101	5PD	N09-C10-C12-C14


There are no ring outliers.

3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	O	101	5PD	1	0
4	K	101	5PD	1	0
4	E	101	5PD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## Ligand 5PD G 101



## Ligand 5PD A 101





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	436/477 (91%)	-0.04	5 (1%) 80 82	20, 35, 62, 80	0
1	D	437/477 (91%)	-0.09	3 (0%) 87 89	20, 34, 63, 81	0
1	F	435/477 (91%)	-0.04	11 (2%) 57 61	22, 41, 72, 115	0
1	H	435/477 (91%)	0.05	18 (4%) 37 40	25, 43, 75, 114	0
1	J	436/477 (91%)	0.11	13 (2%) 50 53	25, 43, 76, 116	0
1	L	435/477 (91%)	0.24	23 (5%) 26 28	24, 47, 86, 116	0
1	N	435/477 (91%)	0.11	17 (3%) 39 42	25, 45, 76, 129	0
1	P	436/477 (91%)	0.01	10 (2%) 60 63	23, 41, 70, 103	0
2	A	75/77 (97%)	-0.13	0 100 100	23, 37, 61, 81	0
2	C	74/77 (96%)	-0.07	1 (1%) 75 77	23, 38, 61, 81	0
2	E	72/77 (93%)	-0.20	0 100 100	26, 42, 64, 80	0
2	G	75/77 (97%)	-0.14	0 100 100	28, 41, 67, 76	0
2	I	72/77 (93%)	0.03	1 (1%) 75 77	25, 39, 57, 86	0
2	K	73/77 (94%)	0.52	7 (9%) 8 7	34, 58, 89, 105	0
2	M	70/77 (90%)	0.34	3 (4%) 35 38	31, 56, 87, 105	0
2	O	74/77 (96%)	0.17	4 (5%) 25 27	28, 47, 75, 85	0
All	All	4070/4432 (91%)	0.05	116 (2%) 51 55	20, 41, 74, 129	0

The worst 5 of 116 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	342	SER	5.2
1	P	342	SER	5.0
1	L	340	GLU	4.6
1	H	341	ASP	4.0
1	N	342	SER	3.9

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	P	501	6/6	0.77	0.44	63,66,75,77	0
3	GOL	D	501	6/6	0.84	0.53	58,63,67,70	0
3	GOL	B	501	6/6	0.86	0.28	51,53,56,61	0
4	5PD	K	101	21/21	0.94	0.26	37,50,65,89	0
4	5PD	E	101	21/21	0.94	0.34	28,38,61,74	0
4	5PD	G	101	21/21	0.97	0.22	27,34,59,63	0
4	5PD	M	101	21/21	0.97	0.24	34,44,62,71	0
4	5PD	I	101	21/21	0.98	0.29	28,32,60,64	0
4	5PD	A	101	21/21	0.98	0.26	19,24,49,56	0
4	5PD	C	101	21/21	0.98	0.24	21,24,53,63	0
4	5PD	O	101	21/21	0.98	0.22	25,32,53,64	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 5PD K 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around 5PD E 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 5PD G 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around 5PD M 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around 5PD I 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around 5PD A 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 5PD C 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 5PD O 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.