



# Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 01:25 am BST

PDB ID : 6EKM  
Title : Crystal structure of mammalian Rev7 in complex with human Rev3 second binding site  
Authors : Huber, F.; Tropia, L.; Emamzadah, S.; Halazonetis, T.  
Deposited on : 2017-09-26  
Resolution : 2.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	211	<div> <div>3%</div> <div> <div></div> <div>57%</div> <div>32%</div> <div>5%</div> <div>6%</div> </div> </div>
2	B	28	<div> <div>4%</div> <div> <div></div> <div>50%</div> <div>29%</div> <div>11%</div> <div>11%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1839 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitotic spindle assembly checkpoint protein MAD2B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	199	Total	C	N	O	S	0	0	0
			1619	1045	274	292	8			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	11	SER	PHE	engineered mutation	UNP Q9D752
A	12	ALA	GLY	engineered mutation	UNP Q9D752
A	132	LYS	VAL	engineered mutation	UNP Q9D752
A	133	VAL	CYS	engineered mutation	UNP Q9D752
A	135	LYS	ALA	engineered mutation	UNP Q9D752

- Molecule 2 is a protein called DNA polymerase zeta catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	25	Total	C	N	O	S	0	0	0
			199	129	36	31	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1987	MET	-	initiating methionine	UNP O60673
B	1988	GLY	-	expression tag	UNP O60673


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	17	Total	O	0	0
			17	17		
3	B	4	Total	O	0	0
			4	4		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitotic spindle assembly checkpoint protein MAD2B



- Molecule 2: DNA polymerase zeta catalytic subunit



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	39.20 Å 48.64 Å 116.81 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.40 – 2.76 44.90 – 2.76	Depositor EDS
% Data completeness (in resolution range)	99.5 (58.40-2.76) 99.5 (44.90-2.76)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.18	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.49 (at 2.77 Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.230 , 0.258 0.224 , 0.243	Depositor DCC
$R_{free}$ test set	298 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	50.4	Xtriage
Anisotropy	0.146	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 54.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	1839	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/1653	0.65	0/2246
2	B	0.43	0/202	0.66	0/271
All	All	0.38	0/1855	0.66	0/2517

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1619	0	1671	63	0
2	B	199	0	221	11	0
3	A	17	0	0	2	0
3	B	4	0	0	0	0
All	All	1839	0	1892	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:183:ASP:HB3	1:A:184:PRO:HA	1.35	1.05

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:47:LYS:HE2	1:A:47:LYS:HA	1.64	0.79
1:A:28:HIS:HD2	1:A:55:SER:H	1.30	0.79
1:A:183:ASP:HB3	1:A:184:PRO:CA	2.11	0.78
1:A:78:ASN:OD1	1:A:104:GLN:HB2	1.85	0.76
1:A:48:TYR:O	1:A:124:ARG:HD3	1.93	0.69
1:A:184:PRO:HB2	1:A:205:GLU:O	1.92	0.69
1:A:15:VAL:HG11	1:A:104:GLN:HE21	1.61	0.66
1:A:48:TYR:HD1	1:A:120:GLU:HG3	1.62	0.64
1:A:119:VAL:HG21	1:A:196:ILE:HD11	1.78	0.64
1:A:162:LYS:HZ2	1:A:162:LYS:H	1.47	0.61
1:A:116:LEU:HA	1:A:119:VAL:HG23	1.81	0.61
1:A:185:ARG:CZ	1:A:187:ILE:HD11	2.31	0.61
1:A:176:GLU:O	1:A:180:HIS:HB2	2.01	0.60
1:A:173:LEU:HD21	2:B:1992:ILE:HD11	1.84	0.60
1:A:23:LEU:O	1:A:27:VAL:HG23	2.03	0.59
1:A:173:LEU:HD21	2:B:1992:ILE:CG1	2.32	0.58
2:B:2011:GLN:NE2	2:B:2011:GLN:HA	2.18	0.58
1:A:93:ARG:HH22	1:A:180:HIS:HB3	1.69	0.57
1:A:79:ASP:O	1:A:152:THR:HG22	2.05	0.56
1:A:74:LEU:HD13	1:A:152:THR:HG21	1.89	0.55
1:A:172:ILE:HG23	2:B:1997:CYS:SG	2.46	0.55
1:A:162:LYS:NZ	1:A:162:LYS:H	2.04	0.54
1:A:27:VAL:O	1:A:31:LEU:HG	2.07	0.54
1:A:28:HIS:CD2	1:A:55:SER:H	2.17	0.54
1:A:105:PRO:HG2	1:A:195:ASP:O	2.07	0.54
1:A:162:LYS:O	1:A:166:ILE:HG12	2.08	0.53
2:B:2011:GLN:C	2:B:2013:LYS:H	2.13	0.52
1:A:15:VAL:HG11	1:A:104:GLN:NE2	2.25	0.51
1:A:173:LEU:HD21	2:B:1992:ILE:CD1	2.42	0.50
2:B:1991:LYS:HB3	2:B:1991:LYS:NZ	2.26	0.50
1:A:186:LEU:HD23	1:A:204:GLU:HG2	1.94	0.50
1:A:72:LYS:O	1:A:76:GLU:HG3	2.12	0.49
1:A:125:ALA:HB1	1:A:190:LYS:HE3	1.93	0.49
2:B:2011:GLN:HE21	2:B:2011:GLN:HA	1.76	0.49
1:A:47:LYS:HA	1:A:47:LYS:CE	2.39	0.48
1:A:69:HIS:O	1:A:163:ILE:HD13	2.13	0.48
1:A:160:MET:HB3	1:A:164:GLN:HE21	1.79	0.48
1:A:123:LEU:HD23	1:A:192:MET:CE	2.44	0.48
1:A:183:ASP:CB	1:A:184:PRO:CA	2.90	0.46
1:A:173:LEU:HD21	2:B:1992:ILE:HG13	1.96	0.46
1:A:137:LEU:HD21	1:A:187:ILE:HD12	1.97	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:69:HIS:ND1	1:A:167:LYS:HE3	2.32	0.45
1:A:75:LEU:HD23	1:A:80:VAL:HG21	1.97	0.45
1:A:160:MET:O	1:A:164:GLN:HG3	2.16	0.45
2:B:1990:LYS:HG3	2:B:1991:LYS:H	1.81	0.45
1:A:180:HIS:HE1	3:A:316:HOH:O	1.97	0.45
1:A:126:PHE:O	1:A:130:ILE:HG13	2.16	0.45
1:A:119:VAL:O	1:A:123:LEU:HG	2.17	0.45
1:A:19:LEU:HD12	1:A:19:LEU:HA	1.85	0.44
1:A:154:GLU:O	1:A:155:ALA:HB2	2.18	0.44
1:A:175:ASP:N	1:A:175:ASP:OD1	2.47	0.43
1:A:86:VAL:HG22	1:A:97:LYS:HG2	2.01	0.43
1:A:50:VAL:HG13	1:A:51:PRO:HD2	2.00	0.43
1:A:115:LEU:O	1:A:118:HIS:HB3	2.17	0.43
1:A:94:PRO:HB2	1:A:181:MET:HE3	2.01	0.43
1:A:193:THR:O	1:A:194:SER:HB3	2.19	0.42
1:A:40:GLY:HA3	2:B:2010:LEU:HD11	2.01	0.42
1:A:90:LYS:C	1:A:92:HIS:H	2.22	0.42
1:A:93:ARG:HG2	1:A:93:ARG:HH11	1.85	0.42
1:A:97:LYS:HB2	1:A:204:GLU:HB2	2.02	0.42
1:A:116:LEU:C	1:A:118:HIS:N	2.73	0.41
1:A:182:HIS:O	1:A:183:ASP:CB	2.69	0.41
1:A:82:LYS:HG2	1:A:101:GLU:HG2	2.02	0.41
1:A:44:LYS:HB2	3:A:310:HOH:O	2.21	0.41
1:A:93:ARG:HA	1:A:94:PRO:HD3	1.97	0.41
1:A:75:LEU:HD23	1:A:80:VAL:CG2	2.50	0.40
1:A:32:TYR:CE2	1:A:53:GLN:HG3	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/211 (93%)	179 (91%)	13 (7%)	5 (2%)	5	9
2	B	23/28 (82%)	22 (96%)	1 (4%)	0	100	100
All	All	220/239 (92%)	201 (91%)	14 (6%)	5 (2%)	6	10

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	155	ALA
1	A	183	ASP
1	A	154	GLU
1	A	112	SER
1	A	184	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/201 (94%)	176 (93%)	13 (7%)	15	27
2	B	23/25 (92%)	16 (70%)	7 (30%)	0	0
All	All	212/226 (94%)	192 (91%)	20 (9%)	8	15

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	LYS
1	A	47	LYS
1	A	81	GLU
1	A	107	LEU
1	A	109	SER
1	A	133	VAL
1	A	162	LYS
1	A	175	ASP
1	A	176	GLU
1	A	181	MET
1	A	184	PRO

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	193	THR
1	A	195	ASP
2	B	1990	LYS
2	B	1999	CYS
2	B	2003	ARG
2	B	2004	GLN
2	B	2007	GLN
2	B	2010	LEU
2	B	2013	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	ASN
1	A	53	GLN
1	A	104	GLN
1	A	139	HIS
1	A	177	GLN
2	B	2004	GLN
2	B	2007	GLN
2	B	2011	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	199/211 (94%)	0.07	6 (3%) 50 59	26, 51, 95, 107	0
2	B	25/28 (89%)	0.12	1 (4%) 38 45	39, 52, 90, 94	0
All	All	224/239 (93%)	0.08	7 (3%) 49 58	26, 52, 94, 107	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	113	ASP	5.5
1	A	182	HIS	4.9
2	B	1989	ASP	4.4
1	A	114	SER	4.1
1	A	93	ARG	2.6
1	A	195	ASP	2.5
1	A	117	SER	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.