



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 11:05 am BST

PDB ID : 3EMZ  
Title : Crystal structure of xylanase XynB from *Paenibacillus barcinonensis* complexed with a conduramine derivative  
Authors : Sanz-Aparicio, J.; Isorna, P.  
Deposited on : 2008-09-25  
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6189 (2.10-2.06)
Clashscore	141614	6738 (2.10-2.06)
Ramachandran outliers	138981	6663 (2.10-2.06)
Sidechain outliers	138945	6664 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	331	

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3007 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endo-1,4-beta-xylanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	331	Total	C	N	O	S	0	0	0
			2715	1717	471	517	10			

- Molecule 2 is (1S,2S,3R,6R)-6-[(4-phenoxybenzyl)amino]cyclohex-4-ene-1,2,3-triol (three-letter code: HXH) (formula: C<sub>19</sub>H<sub>21</sub>NO<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	7	0
			24	19	1	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	268	Total	O	0	0
			268	268		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Endo-1,4-beta-xylanase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	44.02Å 79.72Å 91.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	18.85 – 2.08 18.85 – 1.98	Depositor EDS
% Data completeness (in resolution range)	99.6 (18.85-2.08) 91.2 (18.85-1.98)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.10 (at 1.97Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.216 , 0.264 0.324 , 0.338	Depositor DCC
$R_{free}$ test set	1439 reflections (6.84%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.0	Xtriage
Anisotropy	0.469	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.44 , 45.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.84	EDS
Total number of atoms	3007	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HXH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/2785	0.62	1/3778 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	239	VAL	N-CA-C	-6.20	94.26	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	176	TYR	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2715	0	2574	78	0
2	A	24	0	21	1	0
3	A	268	0	0	9	0
All	All	3007	0	2595	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:122:TYR:HB3	1:A:126:ILE:HD13	1.60	0.81
1:A:5:ILE:HG21	1:A:321:PRO:HG3	1.62	0.80
1:A:271:TYR:HB3	1:A:328:ILE:HD11	1.63	0.79
1:A:3:THR:HG23	1:A:4:GLU:H	1.45	0.79
1:A:81:HIS:O	3:A:1001:HOH:O	2.04	0.75
1:A:314:VAL:HG11	1:A:328:ILE:HD12	1.69	0.73
1:A:18:ILE:H	1:A:39:ASN:HD21	1.39	0.69
1:A:331:GLN:O	1:A:332:ASP:HB2	1.93	0.68
1:A:325:PHE:CZ	1:A:329:ILE:HD11	2.29	0.68
1:A:28:GLN:HB2	3:A:1256:HOH:O	1.95	0.67
1:A:82:THR:H	1:A:130:ASP:HB2	1.61	0.64
1:A:270:ARG:O	1:A:274:ILE:HG12	1.99	0.63
1:A:23:HIS:O	1:A:26:MET:HG3	2.00	0.61
1:A:271:TYR:CB	1:A:328:ILE:HD11	2.30	0.61
1:A:83:LEU:HD13	1:A:115:ILE:HD12	1.83	0.61
1:A:111:LEU:O	1:A:115:ILE:HG12	2.01	0.61
1:A:2:SER:N	1:A:36:LYS:HZ2	2.01	0.58
1:A:122:TYR:CB	1:A:126:ILE:HD13	2.32	0.57
1:A:48:LYS:HG2	1:A:82:THR:CG2	2.34	0.57
1:A:81:HIS:HB3	3:A:1001:HOH:O	2.04	0.57
1:A:5:ILE:HD11	3:A:1140:HOH:O	2.05	0.56
1:A:17:LYS:HE2	3:A:1168:HOH:O	2.05	0.56
1:A:271:TYR:CG	1:A:328:ILE:HD11	2.42	0.55
1:A:218:PRO:O	1:A:270:ARG:NH2	2.30	0.55
1:A:271:TYR:CD1	1:A:328:ILE:HD11	2.42	0.55
1:A:137:GLU:HG2	1:A:145:ARG:HA	1.88	0.54
1:A:55:ARG:HH21	1:A:58:GLU:CD	2.12	0.53
1:A:29:THR:HG23	1:A:30:GLU:HG3	1.90	0.52
1:A:17:LYS:C	1:A:18:ILE:HD12	2.30	0.51
1:A:32:GLU:HG3	3:A:1219:HOH:O	2.10	0.51
1:A:50:GLU:HG2	1:A:88:GLN:O	2.11	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:292:GLY:HA3	1:A:313:PHE:CZ	2.46	0.50
1:A:299:TRP:HH2	2:A:1000:HXX:H16	1.76	0.50
1:A:241:GLU:HG2	1:A:291:TRP:CE3	2.47	0.50
1:A:217:GLY:C	1:A:270:ARG:NH2	2.65	0.50
1:A:320:GLN:HG3	1:A:321:PRO:HD2	1.94	0.49
1:A:18:ILE:H	1:A:39:ASN:ND2	2.07	0.49
1:A:211:GLY:HA2	1:A:213:TRP:CE2	2.48	0.48
1:A:36:LYS:NZ	3:A:1205:HOH:O	2.47	0.48
1:A:136:ILE:CD1	1:A:190:ILE:HA	2.43	0.48
1:A:136:ILE:HD13	1:A:190:ILE:HA	1.96	0.48
1:A:64:ALA:O	1:A:68:VAL:HG23	2.14	0.48
1:A:137:GLU:HG3	1:A:142:LEU:HD22	1.96	0.47
1:A:49:PHE:CD1	1:A:90:PRO:HD2	2.50	0.47
1:A:238:HIS:CD2	1:A:287:SER:HB3	2.50	0.47
1:A:314:VAL:CG1	1:A:328:ILE:HD12	2.43	0.46
1:A:105:GLU:H	1:A:105:GLU:CD	2.18	0.46
1:A:166:HIS:HD2	3:A:1048:HOH:O	1.97	0.46
1:A:266:LEU:O	1:A:270:ARG:HB2	2.14	0.46
1:A:53:HIS:HD2	1:A:114:HIS:ND1	2.13	0.46
1:A:204:VAL:HG13	1:A:204:VAL:O	2.17	0.45
1:A:217:GLY:C	1:A:270:ARG:HH21	2.20	0.45
1:A:5:ILE:HG22	1:A:326:TRP:CH2	2.52	0.45
1:A:45:ASN:O	1:A:51:GLU:HG3	2.17	0.45
1:A:139:LYS:NZ	1:A:141:ASP:OD1	2.50	0.44
1:A:82:THR:OG1	1:A:85:TRP:HB2	2.17	0.44
1:A:53:HIS:HE1	1:A:56:GLU:O	2.00	0.44
1:A:17:LYS:O	1:A:18:ILE:HD12	2.18	0.44
1:A:21:ALA:HB2	1:A:291:TRP:CE3	2.52	0.43
1:A:3:THR:HG23	1:A:4:GLU:N	2.23	0.43
1:A:55:ARG:HB2	1:A:58:GLU:HB3	2.01	0.43
1:A:5:ILE:HA	1:A:6:PRO:HD3	1.85	0.43
1:A:259:PRO:HD3	1:A:311:TRP:CD2	2.54	0.43
1:A:249:HIS:O	1:A:250:GLU:HG2	2.18	0.43
1:A:5:ILE:HD11	1:A:37:HIS:HE2	1.84	0.43
1:A:12:TYR:HB2	1:A:18:ILE:CD1	2.49	0.42
1:A:123:LYS:O	1:A:124:ASP:HB2	2.20	0.42
1:A:174:LEU:HD12	1:A:203:PRO:O	2.20	0.42
1:A:11:SER:O	1:A:332:ASP:N	2.49	0.42
1:A:212:HIS:HA	1:A:243:ASP:OD1	2.17	0.42
1:A:238:HIS:CG	1:A:287:SER:HB3	2.55	0.41
1:A:148:LYS:HG3	3:A:1101:HOH:O	2.20	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223:ILE:O	1:A:227:ILE:HG12	2.21	0.41
1:A:5:ILE:HG13	1:A:5:ILE:O	2.20	0.41
1:A:39:ASN:HD22	1:A:39:ASN:C	2.25	0.40
1:A:12:TYR:HB2	1:A:18:ILE:HD13	2.02	0.40
1:A:180:ASN:HA	1:A:180:ASN:HD22	1.66	0.40
1:A:39:ASN:ND2	1:A:39:ASN:H	2.19	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	329/331 (99%)	312 (95%)	15 (5%)	2 (1%)	25	20

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	82	THR
1	A	3	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/288 (100%)	278 (96%)	10 (4%)	36	36

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	THR
1	A	26	MET
1	A	28	GLN
1	A	39	ASN
1	A	53	HIS
1	A	58	GLU
1	A	142	LEU
1	A	268	GLN
1	A	312	PRO
1	A	331	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	ASN
1	A	53	HIS
1	A	87	ASN
1	A	160	GLN
1	A	166	HIS
1	A	180	ASN
1	A	192	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	HXH	A	1000	-	26,26,26	2.42	7 (26%)	30,35,35	1.25	3 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HXH	A	1000	-	-	1/9/26/26	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1000	HXH	C4-C5	-8.53	1.34	1.50
2	A	1000	HXH	O20-C14	3.73	1.47	1.39
2	A	1000	HXH	C10-N1	3.53	1.57	1.46
2	A	1000	HXH	O20-C21	3.42	1.47	1.39
2	A	1000	HXH	C25-C26	3.10	1.45	1.38
2	A	1000	HXH	C13-C12	2.80	1.43	1.38
2	A	1000	HXH	C16-C15	2.23	1.42	1.38

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	HXH	C21-O20-C14	3.89	127.91	118.80
2	A	1000	HXH	C6-C1-N1	3.37	116.13	110.36
2	A	1000	HXH	C1-C6-C5	-2.63	110.39	120.00

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1000	HXH	C11-C10-N1-C1

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1000	HXH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


### 6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.