



wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2020 – 06:52 am BST

PDB ID : 2EQ9
Title : Crystal structure of lipoamide dehydrogenase from thermus thermophilus HB8 with psbdb
Authors : Nakai, T.; Kamiya, N.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2007-03-30
Resolution : 2.09 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	464	<div> <div>4%</div> <div> <div></div> <div>81%</div> <div>18%</div> <div>..</div> </div> </div>
1	B	464	<div> <div>%</div> <div> <div></div> <div>83%</div> <div>15%</div> <div>..</div> </div> </div>
1	D	464	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>13%</div> <div>..</div> </div> </div>
1	E	464	<div> <div>4%</div> <div> <div></div> <div>84%</div> <div>14%</div> <div>..</div> </div> </div>
1	G	464	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>16%</div> <div>..</div> </div> </div>
1	H	464	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>16%</div> <div>..</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	J	464	<div><div>8%</div><div>79%</div><div>19%</div><div>..</div></div>
1	K	464	<div><div>13%</div><div>75%</div><div>23%</div><div>..</div></div>
2	C	41	<div><div>20%</div><div>76%</div><div>20%</div><div>..</div></div>
2	F	41	<div><div>63%</div><div>63%</div><div>34%</div><div>.</div></div>
2	I	41	<div><div>51%</div><div>66%</div><div>24%</div><div>7%</div><div>.</div></div>
2	L	41	<div><div>59%</div><div>61%</div><div>32%</div><div>5%</div><div>.</div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 29827 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyruvate dehydrogenase complex, dihydrolipoamide dehydrogenase E3 component.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	460	Total	C	N	O	S	0	0	0
			3409	2170	596	634	9			
1	B	460	Total	C	N	O	S	0	0	0
			3397	2162	594	632	9			
1	D	460	Total	C	N	O	S	0	0	0
			3413	2173	597	634	9			
1	E	460	Total	C	N	O	S	0	0	0
			3405	2167	595	634	9			
1	G	460	Total	C	N	O	S	0	0	0
			3409	2170	596	634	9			
1	H	460	Total	C	N	O	S	0	0	0
			3397	2162	594	632	9			
1	J	460	Total	C	N	O	S	0	0	0
			3409	2170	596	634	9			
1	K	460	Total	C	N	O	S	0	0	0
			3397	2162	594	632	9			


- Molecule 2 is a protein called Pyruvate dehydrogenase complex, dihydrolipoamide acetyltransferase E2 component.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	40	Total	C	N	O	S	0	0	0
			290	184	53	52	1			
2	F	40	Total	C	N	O	S	0	0	0
			290	184	53	52	1			
2	I	40	Total	C	N	O	S	0	0	0
			290	184	53	52	1			
2	L	40	Total	C	N	O	S	0	0	0
			290	184	53	52	1			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	129	MET	-	INITIATING METHIONINE	UNP Q5SLR1
F	129	MET	-	INITIATING METHIONINE	UNP Q5SLR1
I	129	MET	-	INITIATING METHIONINE	UNP Q5SLR1
L	129	MET	-	INITIATING METHIONINE	UNP Q5SLR1

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	E	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	G	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	H	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	J	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	K	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	114	Total 114	O 114	0	0
4	B	138	Total 138	O 138	0	0
4	C	4	Total 4	O 4	0	0
4	D	158	Total 158	O 158	0	0
4	E	113	Total 113	O 113	0	0
4	F	6	Total 6	O 6	0	0
4	G	153	Total 153	O 153	0	0
4	H	145	Total 145	O 145	0	0
4	I	2	Total 2	O 2	0	0
4	J	117	Total 117	O 117	0	0
4	K	55	Total 55	O 55	0	0
4	L	2	Total 2	O 2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Pyruvate dehydrogenase complex, dihydrolipoamide dehydrogenase E3 component




- Molecule 1: Pyruvate dehydrogenase complex, dihydrolipoamide dehydrogenase E3 component





- Molecule 1: Pyruvate dehydrogenase complex, dihydrolipoamide dehydrogenase E3 component









- Molecule 1: Pyruvate dehydrogenase complex, dihydrolipoamide dehydrogenase E3 component




- Molecule 1: Pyruvate dehydrogenase complex, dihydrolipoamide dehydrogenase E3 component





- Molecule 1: Pyruvate dehydrogenase complex, dihydrolipoamide dehydrogenase E3 component









- Molecule 1: Pyruvate dehydrogenase complex, dihydrolipoamide dehydrogenase E3 component



- Molecule 1: Pyruvate dehydrogenase complex, dihydrolipoamide dehydrogenase E3 component



- Molecule 2: Pyruvate dehydrogenase complex, dihydrolipoamide acetyltransferase E2 component




- Molecule 2: Pyruvate dehydrogenase complex, dihydrolipoamide acetyltransferase E2 component






- Molecule 2: Pyruvate dehydrogenase complex, dihydrolipoamide acetyltransferase E2 component



- Molecule 2: Pyruvate dehydrogenase complex, dihydrolipoamide acetyltransferase E2 component



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	88.78Å 88.88Å 144.15Å 89.26° 87.01° 70.84°	Depositor
Resolution (Å)	43.62 – 2.09 43.62 – 2.09	Depositor EDS
% Data completeness (in resolution range)	96.6 (43.62-2.09) 96.7 (43.62-2.09)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 2.08Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.217 , 0.257 0.207 , 0.246	Depositor DCC
R_{free} test set	11948 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	35.1	Xtriage
Anisotropy	0.154	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.059 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	29827	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.31	0/3464	0.61	0/4683
1	B	0.32	0/3452	0.61	0/4670
1	D	0.34	0/3468	0.62	0/4687
1	E	0.33	0/3460	0.61	0/4679
1	G	0.34	0/3464	0.62	0/4683
1	H	0.33	0/3452	0.61	1/4670 (0.0%)
1	J	0.31	0/3464	0.60	0/4683
1	K	0.29	0/3452	0.58	0/4670
2	C	0.26	0/294	0.60	1/399 (0.3%)
2	F	0.25	0/294	0.59	1/399 (0.3%)
2	I	0.28	0/294	0.56	0/399
2	L	0.26	0/294	0.57	0/399
All	All	0.32	0/28852	0.61	3/39021 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	156	GLY	N-CA-C	5.69	127.33	113.10
2	F	156	GLY	N-CA-C	5.51	126.86	113.10
1	H	52	CYS	CA-CB-SG	5.46	123.83	114.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3409	0	3519	66	0
1	B	3397	0	3493	61	0
1	D	3413	0	3530	45	0
1	E	3405	0	3508	52	0
1	G	3409	0	3519	52	0
1	H	3397	0	3493	61	0
1	J	3409	0	3519	78	0
1	K	3397	0	3493	91	0
2	C	290	0	300	8	0
2	F	290	0	300	9	0
2	I	290	0	300	12	0
2	L	290	0	300	12	0
3	A	53	0	31	2	0
3	B	53	0	31	0	0
3	D	53	0	31	1	0
3	E	53	0	31	0	0
3	G	53	0	31	1	0
3	H	53	0	31	0	0
3	J	53	0	31	0	0
3	K	53	0	31	3	0
4	A	114	0	0	1	0
4	B	138	0	0	1	0
4	C	4	0	0	0	0
4	D	158	0	0	0	0
4	E	113	0	0	1	0
4	F	6	0	0	0	0
4	G	153	0	0	2	0
4	H	145	0	0	2	0
4	I	2	0	0	0	0
4	J	117	0	0	1	0
4	K	55	0	0	2	0
4	L	2	0	0	0	0
All	All	29827	0	29522	530	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 530 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:388:ALA:HB1	1:B:394:ALA:HB2	1.39	1.04
1:D:233:ARG:HH11	1:D:233:ARG:HG2	1.24	1.00
1:J:11:ASP:HA	1:J:141:LYS:HD3	1.42	0.99
1:A:141:LYS:H	1:A:141:LYS:HD2	1.36	0.88
1:H:439:LEU:HD21	1:H:453:LEU:HD22	1.54	0.88

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	458/464 (99%)	443 (97%)	14 (3%)	1 (0%)	47	49
1	B	458/464 (99%)	444 (97%)	14 (3%)	0	100	100
1	D	458/464 (99%)	440 (96%)	17 (4%)	1 (0%)	47	49
1	E	458/464 (99%)	445 (97%)	13 (3%)	0	100	100
1	G	458/464 (99%)	449 (98%)	9 (2%)	0	100	100
1	H	458/464 (99%)	449 (98%)	9 (2%)	0	100	100
1	J	458/464 (99%)	434 (95%)	24 (5%)	0	100	100
1	K	458/464 (99%)	433 (94%)	23 (5%)	2 (0%)	34	32
2	C	38/41 (93%)	35 (92%)	3 (8%)	0	100	100
2	F	38/41 (93%)	34 (90%)	3 (8%)	1 (3%)	5	2
2	I	38/41 (93%)	35 (92%)	3 (8%)	0	100	100
2	L	38/41 (93%)	30 (79%)	8 (21%)	0	100	100
All	All	3816/3876 (98%)	3671 (96%)	140 (4%)	5 (0%)	51	54

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	284	GLU
1	D	284	GLU
1	K	279	GLU
1	K	148	GLY
2	F	153	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/349 (98%)	338 (99%)	4 (1%)	71	77
1	B	339/349 (97%)	334 (98%)	5 (2%)	65	71
1	D	343/349 (98%)	335 (98%)	8 (2%)	50	55
1	E	341/349 (98%)	337 (99%)	4 (1%)	71	77
1	G	342/349 (98%)	338 (99%)	4 (1%)	71	77
1	H	339/349 (97%)	333 (98%)	6 (2%)	59	65
1	J	342/349 (98%)	336 (98%)	6 (2%)	59	65
1	K	339/349 (97%)	336 (99%)	3 (1%)	78	84
2	C	28/31 (90%)	26 (93%)	2 (7%)	14	11
2	F	28/31 (90%)	26 (93%)	2 (7%)	14	11
2	I	28/31 (90%)	25 (89%)	3 (11%)	6	3
2	L	28/31 (90%)	25 (89%)	3 (11%)	6	3
All	All	2839/2916 (97%)	2789 (98%)	50 (2%)	59	65

5 of 50 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	453	LEU
1	G	233	ARG
1	K	233	ARG
2	F	129	MET
1	G	56	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	418	GLN
1	H	214	GLN
1	J	30	GLN
1	E	418	GLN
1	J	23	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FAD	H	5482	-	51,58,58	2.36	19 (37%)	60,89,89	1.69	8 (13%)
3	FAD	D	2482	-	51,58,58	2.32	17 (33%)	60,89,89	1.70	10 (16%)
3	FAD	B	1482	-	51,58,58	2.32	17 (33%)	60,89,89	1.72	10 (16%)
3	FAD	A	8482	-	51,58,58	2.33	15 (29%)	60,89,89	1.71	11 (18%)
3	FAD	G	4482	-	51,58,58	2.33	18 (35%)	60,89,89	1.72	10 (16%)
3	FAD	E	3482	-	51,58,58	2.36	16 (31%)	60,89,89	1.73	10 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FAD	K	7482	-	51,58,58	2.48	17 (33%)	60,89,89	1.73	10 (16%)
3	FAD	J	6482	-	51,58,58	2.35	17 (33%)	60,89,89	1.73	11 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	H	5482	-	-	2/30/50/50	0/6/6/6
3	FAD	D	2482	-	-	2/30/50/50	0/6/6/6
3	FAD	B	1482	-	-	2/30/50/50	0/6/6/6
3	FAD	A	8482	-	-	2/30/50/50	0/6/6/6
3	FAD	G	4482	-	-	3/30/50/50	0/6/6/6
3	FAD	E	3482	-	-	2/30/50/50	0/6/6/6
3	FAD	K	7482	-	-	3/30/50/50	0/6/6/6
3	FAD	J	6482	-	-	2/30/50/50	0/6/6/6

The worst 5 of 136 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	K	7482	FAD	C4X-C10	9.22	1.48	1.38
3	H	5482	FAD	C4X-C10	8.95	1.47	1.38
3	E	3482	FAD	C4X-C10	8.91	1.47	1.38
3	J	6482	FAD	C4X-C10	8.67	1.47	1.38
3	D	2482	FAD	C4X-C10	8.63	1.47	1.38

The worst 5 of 80 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	K	7482	FAD	C4-N3-C2	7.89	121.81	115.14
3	E	3482	FAD	C4-N3-C2	7.79	121.72	115.14
3	B	1482	FAD	C4-N3-C2	7.76	121.69	115.14
3	H	5482	FAD	C4-N3-C2	7.70	121.64	115.14
3	A	8482	FAD	C4-N3-C2	7.63	121.59	115.14

There are no chirality outliers.

5 of 18 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	2482	FAD	PA-O3P-P-O5'
3	B	1482	FAD	PA-O3P-P-O5'
3	A	8482	FAD	PA-O3P-P-O5'
3	E	3482	FAD	PA-O3P-P-O5'
3	H	5482	FAD	PA-O3P-P-O5'


There are no ring outliers.


4 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	2482	FAD	1	0
3	A	8482	FAD	2	0
3	G	4482	FAD	1	0
3	K	7482	FAD	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	460/464 (99%)	0.11	20 (4%) 35 41	25, 38, 63, 79	0
1	B	460/464 (99%)	-0.09	4 (0%) 84 86	24, 38, 55, 69	0
1	D	460/464 (99%)	-0.06	10 (2%) 62 66	23, 34, 62, 80	0
1	E	460/464 (99%)	0.10	19 (4%) 37 43	22, 38, 70, 85	0
1	G	460/464 (99%)	-0.06	9 (1%) 65 69	24, 36, 61, 76	0
1	H	460/464 (99%)	-0.09	7 (1%) 73 77	24, 35, 55, 73	0
1	J	460/464 (99%)	0.25	36 (7%) 13 17	24, 40, 77, 99	0
1	K	460/464 (99%)	0.65	61 (13%) 3 4	26, 49, 87, 101	0
2	C	40/41 (97%)	0.94	8 (20%) 1 1	33, 55, 74, 76	0
2	F	40/41 (97%)	2.86	26 (65%) 0 0	54, 74, 86, 90	0
2	I	40/41 (97%)	2.69	21 (52%) 0 0	45, 76, 91, 91	0
2	L	40/41 (97%)	2.45	24 (60%) 0 0	47, 77, 87, 88	0
All	All	3840/3876 (99%)	0.19	245 (6%) 19 24	22, 38, 73, 101	0

The worst 5 of 245 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	I	155	LEU	15.1
2	I	129	MET	8.9
2	F	155	LEU	7.9
1	K	120	ARG	6.8
2	I	130	LEU	6.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	FAD	K	7482	53/53	0.93	0.13	40,45,56,56	0
3	FAD	J	6482	53/53	0.95	0.11	30,34,40,41	0
3	FAD	B	1482	53/53	0.96	0.11	33,37,43,44	0
3	FAD	G	4482	53/53	0.96	0.10	29,34,42,44	0
3	FAD	D	2482	53/53	0.97	0.11	26,30,40,41	0
3	FAD	E	3482	53/53	0.97	0.13	31,36,43,43	0
3	FAD	H	5482	53/53	0.97	0.13	25,29,32,33	0
3	FAD	A	8482	53/53	0.97	0.12	28,34,40,41	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD K 7482:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD J 6482:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD B 1482:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD G 4482:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD D 2482:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD E 3482:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD H 5482:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD A 8482:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.