



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 7, 2020 – 08:52 PM BST

PDB ID : 3EQN
Title : Crystal structure of beta-1,3-glucanase from Phanerochaete chrysosporium (Lam55A)
Authors : Ishida, T.; Fushinobu, S.; Kawai, R.; Kitaoka, M.; Igarashi, K.; Samejima, M.
Deposited on : 2008-10-01
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	758	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 91%, green 8%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 91% 8% </div> </div>
1	B	758	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 1%, yellow 93%, green 6%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 2% 93% 6% </div> </div>
2	C	3	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 67%, yellow 33%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 67% 33% </div> </div>
2	D	3	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 67%, yellow 33%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 67% 33% </div> </div>

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 13016 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucan 1,3-beta-glucosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	752	Total	C	N	O	S	0	0	0
			5611	3526	973	1096	16			
1	B	752	Total	C	N	O	S	0	0	0
			5611	3526	973	1096	16			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLU	-	expression tag	UNP Q2Z1W1
A	-4	ALA	-	expression tag	UNP Q2Z1W1
A	-3	GLU	-	expression tag	UNP Q2Z1W1
A	-2	ALA	-	expression tag	UNP Q2Z1W1
A	-1	GLU	-	expression tag	UNP Q2Z1W1
A	0	PHE	-	expression tag	UNP Q2Z1W1
B	-5	GLU	-	expression tag	UNP Q2Z1W1
B	-4	ALA	-	expression tag	UNP Q2Z1W1
B	-3	GLU	-	expression tag	UNP Q2Z1W1
B	-2	ALA	-	expression tag	UNP Q2Z1W1
B	-1	GLU	-	expression tag	UNP Q2Z1W1
B	0	PHE	-	expression tag	UNP Q2Z1W1

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	3	Total	C	N	O	0	0	0
			39	22	2	15			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	3	Total	C	N	O	0	0	0
			39	22	2	15			


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	3	Total	Zn	0	0
			3	3		
3	A	3	Total	Zn	0	0
			3	3		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Na	0	0
			2	2		
4	A	2	Total	Na	0	0
			2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	811	Total	O	0	0
			811	811		
7	B	817	Total	O	0	0
			817	817		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glucan 1,3-beta-glucosidase



- Molecule 1: Glucan 1,3-beta-glucosidase



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	66.28Å 67.15Å 105.18Å 81.04° 76.34° 61.45°	Depositor
Resolution (Å)	27.21 – 1.70 27.21 – 1.70	Depositor EDS
% Data completeness (in resolution range)	88.6 (27.21-1.70) 88.6 (27.21-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.14 (at 1.71Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.149 , 0.184 0.148 , 0.184	Depositor DCC
R_{free} test set	7544 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	14.5	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 55.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13016	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, BMA, NAG, NA, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/5755	0.62	0/7869
1	B	0.55	0/5755	0.62	0/7869
All	All	0.55	0/11510	0.62	0/15738

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	301	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5611	0	5302	36	0
1	B	5611	0	5302	29	0
2	C	39	0	34	0	0
2	D	39	0	34	0	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	A	42	0	56	1	0
5	B	24	0	32	2	0
6	A	4	0	3	0	0
6	B	8	0	6	1	0
7	A	811	0	0	7	0
7	B	817	0	0	6	0
All	All	13016	0	10769	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:763:GOL:H32	7:B:1448:HOH:O	1.76	0.84
1:B:523:HIS:HD2	1:B:570:TRP:H	1.31	0.78
1:A:373:ASP:HA	1:A:514:GLN:HE22	1.56	0.70
1:B:672:HIS:NE2	7:B:1309:HOH:O	2.23	0.70
1:B:176:GLN:H	1:B:176:GLN:HE21	1.42	0.66
1:A:176:GLN:H	1:A:176:GLN:HE21	1.44	0.63
1:A:257:ASN:ND2	7:A:1465:HOH:O	2.33	0.62
1:A:186:GLN:HE21	1:A:186:GLN:HA	1.66	0.60
1:B:505:TRP:HB3	1:B:558:LEU:HD23	1.85	0.58
1:B:301:SER:O	1:B:302:SER:HB2	2.04	0.58
1:A:336:THR:O	1:B:303:GLY:HA2	2.03	0.57
1:B:229:VAL:CG1	1:B:232:LEU:HD22	2.36	0.55
1:B:660:GLN:HG2	7:B:791:HOH:O	2.05	0.54
5:B:763:GOL:H31	7:B:828:HOH:O	2.06	0.54
1:B:523:HIS:HE1	7:B:846:HOH:O	1.92	0.53
1:A:505:TRP:HB3	1:A:558:LEU:HD23	1.89	0.53
1:B:660:GLN:HG3	1:B:660:GLN:O	2.09	0.51
1:A:616:GLN:HG3	1:A:632:THR:HB	1.92	0.51
7:A:1261:HOH:O	1:B:302:SER:HB3	2.09	0.51
1:A:623:ALA:HA	1:A:672:HIS:O	2.11	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:553:PHE:O	1:A:591:ASN:HB3	2.13	0.49
1:A:471:PRO:HA	1:A:501:ILE:O	2.13	0.49
1:A:145:ASN:HD22	1:A:203:GLY:HA2	1.79	0.47
1:B:547:ASN:HD22	1:B:550:PHE:HE2	1.62	0.47
1:B:196:GLN:HA	1:B:218:ILE:O	2.14	0.47
1:B:145:ASN:HD22	1:B:203:GLY:HA2	1.79	0.47
1:A:235:ASN:HD22	1:A:257:ASN:HB2	1.80	0.46
1:A:289:THR:HG23	7:A:1465:HOH:O	2.13	0.46
1:B:616:GLN:HG3	1:B:632:THR:HB	1.97	0.46
1:B:553:PHE:O	1:B:591:ASN:HB3	2.16	0.46
1:B:623:ALA:HA	1:B:672:HIS:O	2.16	0.46
1:A:166:SER:HA	1:A:194:GLN:HB2	1.99	0.45
1:A:344:GLN:HA	1:A:358:THR:O	2.17	0.45
1:B:455:VAL:HG22	1:B:490:ILE:HB	1.97	0.45
1:A:455:VAL:HG22	1:A:490:ILE:HB	1.97	0.45
1:B:229:VAL:HG11	1:B:232:LEU:HD22	1.98	0.45
1:A:196:GLN:HA	1:A:218:ILE:O	2.17	0.44
1:A:547:ASN:HD22	1:A:550:PHE:HE2	1.65	0.44
1:A:168:THR:HA	1:A:196:GLN:O	2.16	0.44
1:A:284:ILE:HB	1:A:677:PHE:CE2	2.52	0.44
1:B:471:PRO:HA	1:B:501:ILE:O	2.17	0.44
1:B:174:VAL:HB	1:B:178:THR:HG21	2.00	0.44
1:B:168:THR:HA	1:B:196:GLN:O	2.18	0.44
1:A:253:ARG:HA	1:A:285:ASP:O	2.18	0.43
1:B:628:GLY:HA2	1:B:629:LEU:HA	1.91	0.43
1:A:290:ASN:O	5:A:767:GOL:H31	2.18	0.43
1:A:30:ILE:HD11	1:A:405:ASP:HA	2.01	0.42
1:A:229:VAL:CG1	1:A:232:LEU:HD22	2.49	0.42
1:A:229:VAL:HG11	1:A:232:LEU:HD22	2.01	0.42
1:A:96:VAL:HG21	1:A:129:LEU:HD21	2.00	0.42
1:A:459:THR:HB	1:A:494:ARG:HB3	2.02	0.42
1:B:523:HIS:HD2	1:B:570:TRP:N	2.08	0.42
1:A:237:ALA:HB2	7:A:823:HOH:O	2.19	0.42
1:A:217:ASN:O	1:A:238:ASN:HB3	2.20	0.42
1:A:290:ASN:ND2	7:A:1014:HOH:O	2.36	0.42
1:B:75:GLN:HG3	1:B:142:TYR:CE1	2.55	0.41
6:B:766:ACT:O	7:B:787:HOH:O	2.22	0.41
1:B:365:ILE:HG12	1:B:714:TYR:CZ	2.56	0.41
1:B:75:GLN:HG3	1:B:142:TYR:CZ	2.56	0.41
1:A:174:VAL:HB	1:A:178:THR:HG21	2.03	0.41
1:A:613:ILE:O	1:A:663:ALA:HB2	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:344:GLN:HA	1:B:358:THR:O	2.21	0.41
1:A:672:HIS:NE2	7:A:1521:HOH:O	2.23	0.41
1:A:365:ILE:HG12	1:A:714:TYR:CZ	2.57	0.40
1:B:152:SER:HA	1:B:179:SER:O	2.21	0.40
1:A:254:ILE:HG12	7:A:1016:HOH:O	2.20	0.40
1:A:231:ASN:HA	1:A:253:ARG:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	750/758 (99%)	717 (96%)	32 (4%)	1 (0%)	51	33
1	B	750/758 (99%)	716 (96%)	32 (4%)	2 (0%)	41	24
All	All	1500/1516 (99%)	1433 (96%)	64 (4%)	3 (0%)	47	30

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	452	VAL
1	B	702	ILE
1	A	702	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	590/594 (99%)	584 (99%)	6 (1%)	76	67
1	B	590/594 (99%)	583 (99%)	7 (1%)	71	59
All	All	1180/1188 (99%)	1167 (99%)	13 (1%)	73	63

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	176	GLN
1	A	186	GLN
1	A	227	PHE
1	A	292	GLN
1	A	448	ILE
1	A	466	TYR
1	B	26	GLN
1	B	176	GLN
1	B	227	PHE
1	B	292	GLN
1	B	352	ASN
1	B	466	TYR
1	B	642	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (36) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	107	GLN
1	A	114	ASN
1	A	145	ASN
1	A	176	GLN
1	A	186	GLN
1	A	196	GLN
1	A	214	ASN
1	A	235	ASN
1	A	242	ASN
1	A	257	ASN
1	A	260	GLN
1	A	292	GLN
1	A	346	ASN
1	A	468	ASN
1	A	514	GLN
1	A	547	ASN
1	A	631	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	642	ASN
1	A	724	GLN
1	A	738	ASN
1	B	48	ASN
1	B	114	ASN
1	B	145	ASN
1	B	176	GLN
1	B	242	ASN
1	B	292	GLN
1	B	346	ASN
1	B	364	ASN
1	B	467	ASN
1	B	523	HIS
1	B	547	ASN
1	B	599	GLN
1	B	631	GLN
1	B	724	GLN
1	B	728	ASN
1	B	738	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	1,2	14,14,15	0.85	0	17,19,21	1.29	2 (11%)
2	NAG	C	2	2	14,14,15	0.42	0	17,19,21	1.01	0
2	BMA	C	3	2	11,11,12	0.60	0	15,15,17	0.57	0
2	NAG	D	1	1,2	14,14,15	0.70	0	17,19,21	1.11	0
2	NAG	D	2	2	14,14,15	0.44	0	17,19,21	0.79	0
2	BMA	D	3	2	11,11,12	0.71	0	15,15,17	0.84	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	NAG	D	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1
2	BMA	D	3	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	3	BMA	C1-O5-C5	2.65	115.78	112.19
2	C	1	NAG	C2-N2-C7	2.43	126.36	122.90
2	C	1	NAG	C1-C2-N2	2.07	114.03	110.49



There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

Of 24 ligands modelled in this entry, 10 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	A	766	-	5,5,5	0.48	0	5,5,5	0.46	0
5	GOL	B	764	-	5,5,5	0.26	0	5,5,5	0.61	0
5	GOL	B	763	-	5,5,5	0.57	0	5,5,5	1.05	0
5	GOL	A	763	-	5,5,5	0.36	0	5,5,5	0.18	0
5	GOL	A	767	-	5,5,5	0.34	0	5,5,5	0.37	0
5	GOL	B	762	-	5,5,5	0.34	0	5,5,5	0.15	0
5	GOL	A	764	-	5,5,5	0.38	0	5,5,5	0.28	0
5	GOL	B	761	-	5,5,5	0.33	0	5,5,5	0.35	0
6	ACT	B	765	3	1,3,3	1.61	0	0,3,3	0.00	-
6	ACT	A	768	3	1,3,3	0.66	0	0,3,3	0.00	-
5	GOL	A	761	-	5,5,5	0.40	0	5,5,5	0.29	0
6	ACT	B	766	-	1,3,3	1.65	0	0,3,3	0.00	-
5	GOL	A	762	-	5,5,5	0.29	0	5,5,5	0.46	0
5	GOL	A	765	-	5,5,5	0.34	0	5,5,5	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	766	-	-	1/4/4/4	-
5	GOL	B	764	-	-	0/4/4/4	-
5	GOL	B	763	-	-	3/4/4/4	-
5	GOL	A	763	-	-	2/4/4/4	-
5	GOL	A	767	-	-	2/4/4/4	-
5	GOL	B	762	-	-	0/4/4/4	-
5	GOL	A	764	-	-	0/4/4/4	-
5	GOL	B	761	-	-	0/4/4/4	-
5	GOL	A	761	-	-	0/4/4/4	-
5	GOL	A	762	-	-	0/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	765	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	763	GOL	O1-C1-C2-C3
5	A	767	GOL	O1-C1-C2-O2
5	A	767	GOL	O1-C1-C2-C3
5	A	765	GOL	O1-C1-C2-C3
5	A	766	GOL	C1-C2-C3-O3
5	A	763	GOL	O1-C1-C2-C3
5	B	763	GOL	O1-C1-C2-O2
5	A	765	GOL	O2-C2-C3-O3
5	B	763	GOL	C1-C2-C3-O3
5	A	763	GOL	O1-C1-C2-O2
5	A	765	GOL	C1-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	763	GOL	2	0
5	A	767	GOL	1	0
6	B	766	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	752/758 (99%)	-0.33	10 (1%) 77 81	8, 14, 23, 34	0
1	B	752/758 (99%)	-0.28	13 (1%) 70 74	8, 13, 24, 31	0
All	All	1504/1516 (99%)	-0.30	23 (1%) 73 77	8, 14, 23, 34	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	138	GLY	5.0
1	B	138	GLY	4.8
1	B	139	GLY	4.2
1	A	137	ALA	4.1
1	A	139	GLY	3.8
1	B	327	GLY	3.8
1	B	137	ALA	3.8
1	A	5	CYS	3.4
1	A	7	SER	3.2
1	A	302	SER	3.1
1	B	273	THR	3.0
1	B	303	GLY	3.0
1	B	302	SER	2.9
1	B	164	SER	2.9
1	A	273	THR	2.5
1	B	136	LEU	2.5
1	B	16	PRO	2.4
1	A	6	SER	2.3
1	A	407	HIS	2.2
1	A	8	PRO	2.2
1	B	6	SER	2.1
1	B	11	HIS	2.1
1	B	691	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BMA	C	3	11/12	0.46	0.35	41,43,45,46	0
2	NAG	C	2	14/15	0.69	0.30	33,37,40,40	0
2	BMA	D	3	11/12	0.77	0.27	33,37,39,40	0
2	NAG	D	2	14/15	0.89	0.16	23,26,29,29	0
2	NAG	C	1	14/15	0.91	0.12	14,18,22,28	0
2	NAG	D	1	14/15	0.95	0.09	11,14,16,20	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	A	767	6/6	0.74	0.17	37,38,38,41	0
6	ACT	B	766	4/4	0.82	0.15	37,37,38,38	0
5	GOL	B	763	6/6	0.90	0.17	21,25,27,29	0
5	GOL	B	764	6/6	0.91	0.14	19,25,26,27	0
5	GOL	A	765	6/6	0.93	0.13	19,25,25,26	0
5	GOL	A	766	6/6	0.93	0.10	22,25,25,25	0
5	GOL	A	763	6/6	0.93	0.13	25,27,28,29	0
6	ACT	A	768	4/4	0.93	0.12	20,21,21,21	0
5	GOL	A	761	6/6	0.95	0.09	16,17,20,22	0
5	GOL	A	762	6/6	0.97	0.07	13,16,17,17	0
5	GOL	B	761	6/6	0.97	0.07	14,15,19,23	0
4	NA	B	760	1/1	0.97	0.06	29,29,29,29	0
6	ACT	B	765	4/4	0.97	0.08	17,17,17,17	0
4	NA	A	759	1/1	0.97	0.11	19,19,19,19	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	GOL	A	764	6/6	0.98	0.07	10,12,13,14	0
4	NA	A	760	1/1	0.98	0.14	29,29,29,29	0
5	GOL	B	762	6/6	0.98	0.07	11,14,16,16	0
4	NA	B	759	1/1	0.99	0.07	21,21,21,21	0
3	ZN	A	756	1/1	0.99	0.13	37,37,37,37	0
3	ZN	B	758	1/1	1.00	0.03	14,14,14,14	0
3	ZN	B	756	1/1	1.00	0.02	17,17,17,17	0
3	ZN	A	758	1/1	1.00	0.02	13,13,13,13	0
3	ZN	A	757	1/1	1.00	0.03	20,20,20,20	0
3	ZN	B	757	1/1	1.00	0.05	21,21,21,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.