



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 03:32 am BST

PDB ID : 4EQX
Title : Crystal Structure of the C43S Mutant of Staphylococcus aureus CoADR
Authors : Edwards, J.S.; Wallace, B.D.; Wallen, J.R.; Claiborne, A.; Redinbo, M.R.
Deposited on : 2012-04-19
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	437	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>12%</div> <div>.</div> </div> </div>
1	B	437	<div> <div>%</div> <div> <div></div> <div>86%</div> <div>13%</div> <div>.</div> </div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8091 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Coenzyme A disulfide reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	437	Total	C	N	O	S	0	10	0
			3521	2241	594	674	12			
1	B	437	Total	C	N	O	S	0	11	0
			3519	2239	592	676	12			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	43	SER	CYS	engineered mutation	UNP Q2FIA5
B	43	SER	CYS	engineered mutation	UNP Q2FIA5

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Cl	0	0
			2	2		
4	A	2	Total	Cl	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	479	Total	O	0	0
			479	479		
5	B	460	Total	O	0	0
			460	460		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Coenzyme A disulfide reductase



- Molecule 1: Coenzyme A disulfide reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.96Å 65.38Å 94.47Å 90.00° 104.83° 90.00°	Depositor
Resolution (Å)	48.85 – 1.70 48.83 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.1 (48.85-1.70) 99.1 (48.83-1.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.89 (at 1.70Å)	Xtriage
Refinement program	PHENIX, REFMAC 5.5.0109	Depositor
R, R_{free}	0.193 , 0.233 0.187 , 0.193	Depositor DCC
R_{free} test set	4872 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	21.9	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.38$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8091	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, FAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.42	8/3615 (0.2%)	1.15	13/4893 (0.3%)
1	B	1.40	15/3621 (0.4%)	1.16	12/4899 (0.2%)
All	All	1.41	23/7236 (0.3%)	1.16	25/9792 (0.3%)

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	26	GLU	CD-OE2	9.73	1.36	1.25
1	B	26	GLU	CG-CD	8.97	1.65	1.51
1	B	26	GLU	CB-CG	8.06	1.67	1.52
1	A	6	VAL	CB-CG2	6.24	1.66	1.52
1	B	415	PHE	CD1-CE1	6.19	1.51	1.39
1	A	160	SER	CB-OG	-6.07	1.34	1.42
1	A	158	TYR	CG-CD1	5.97	1.47	1.39
1	B	255	ARG	CG-CD	5.87	1.66	1.51
1	A	83	ALA	CA-CB	5.86	1.64	1.52
1	B	416	GLU	CD-OE2	5.81	1.32	1.25
1	B	333	PHE	CE2-CZ	5.78	1.48	1.37
1	B	40	PHE	CD2-CE2	5.70	1.50	1.39
1	A	54	GLU	CG-CD	5.70	1.60	1.51
1	B	165	GLU	C-O	5.48	1.33	1.23
1	B	40	PHE	CD1-CE1	5.35	1.50	1.39
1	B	419	TYR	CE2-CZ	5.27	1.45	1.38
1	A	331	TYR	CG-CD1	5.22	1.46	1.39
1	B	420	ALA	CA-CB	5.14	1.63	1.52
1	B	40	PHE	CE1-CZ	5.12	1.47	1.37
1	B	385	ALA	CA-CB	5.11	1.63	1.52
1	B	154	VAL	CB-CG2	5.07	1.63	1.52
1	A	362	TYR	CE2-CZ	5.05	1.45	1.38
1	A	181	ASP	CB-CG	-5.03	1.41	1.51

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	96	ARG	NE-CZ-NH1	10.22	125.41	120.30
1	B	207	ARG	NE-CZ-NH2	-8.39	116.11	120.30
1	B	96	ARG	NE-CZ-NH2	-7.91	116.35	120.30
1	A	24	ASP	CB-CG-OD1	7.56	125.10	118.30
1	A	170	ARG	NE-CZ-NH1	6.85	123.72	120.30
1	B	428	ASP	CB-CG-OD2	-6.84	112.14	118.30
1	A	86	ASP	CB-CG-OD1	6.63	124.27	118.30
1	B	21	ARG	NE-CZ-NH2	-6.50	117.05	120.30
1	B	288	LEU	CB-CG-CD1	-6.47	99.99	111.00
1	A	330	ASP	CB-CG-OD2	-6.20	112.72	118.30
1	B	207	ARG	NE-CZ-NH1	5.99	123.30	120.30
1	A	181	ASP	CB-CG-OD1	-5.95	112.95	118.30
1	A	70	ARG	NE-CZ-NH1	5.84	123.22	120.30
1	B	96	ARG	CD-NE-CZ	5.70	131.57	123.60
1	A	70	ARG	NE-CZ-NH2	-5.58	117.51	120.30
1	B	24	ASP	CB-CG-OD1	5.41	123.17	118.30
1	A	135	ASP	CB-CG-OD1	5.40	123.16	118.30
1	A	96	ARG	NE-CZ-NH2	-5.31	117.64	120.30
1	A	428	ASP	CB-CG-OD2	-5.31	113.53	118.30
1	B	131	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	B	25	LYS	CD-CE-NZ	-5.22	99.70	111.70
1	A	167	LEU	CB-CG-CD1	-5.21	102.15	111.00
1	A	200	ASP	CB-CG-OD1	5.15	122.93	118.30
1	A	55	ASP	CB-CA-C	5.05	120.51	110.40
1	B	135	ASP	CB-CG-OD1	5.02	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3521	0	3510	38	0
1	B	3519	0	3515	35	0
2	A	53	0	31	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	53	0	31	1	0
3	A	2	0	0	0	0
4	A	2	0	0	1	0
4	B	2	0	0	1	0
5	A	479	0	0	14	1
5	B	460	0	0	7	0
All	All	8091	0	7087	69	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159[A]:VAL:HG21	5:A:980:HOH:O	1.27	1.29
1:A:355:GLN:OE1	1:A:432[B]:MET:SD	1.99	1.21
1:A:150:LYS:HE3	5:A:1032:HOH:O	1.66	0.95
1:A:43[A]:SER:OG	4:A:505:CL:CL	2.38	0.78
1:A:151[B]:VAL:HG12	1:A:232[B]:MET:HB3	1.64	0.78
1:B:97:LYS:HG3	5:B:946:HOH:O	1.87	0.74
1:A:2:PRO:N	1:A:27:SER:HG	1.86	0.72
1:A:57:ARG:O	1:A:57:ARG:HG2	1.88	0.72
1:A:129:THR:HG22	1:A:235[A]:GLU:HG3	1.73	0.71
1:A:151[B]:VAL:HG21	1:A:167:LEU:HD13	1.73	0.69
1:A:159[A]:VAL:HG23	5:A:979:HOH:O	1.94	0.68
1:B:43[A]:SER:OG	4:B:503:CL:CL	2.49	0.67
1:B:123:GLU:O	1:B:123:GLU:HG3	1.97	0.65
1:B:193:GLN:NE2	5:B:913:HOH:O	2.29	0.63
1:B:42:ASN:H	1:B:42:ASN:HD22	1.46	0.62
1:B:42:ASN:H	1:B:42:ASN:ND2	1.96	0.62
1:A:159[A]:VAL:CG2	5:A:979:HOH:O	2.48	0.61
1:A:327:LYS:HG3	5:B:970:HOH:O	2.01	0.61
1:A:55:ASP:O	1:A:58:TYR:HB2	2.01	0.60
1:A:291:SER:HB3	5:A:1035:HOH:O	2.00	0.60
1:B:5:VAL:HG22	1:B:30[A]:ILE:HD11	1.83	0.59
1:A:291:SER:CB	5:A:1035:HOH:O	2.49	0.59
1:B:341:ASN:ND2	1:B:342:GLU:HG3	2.18	0.59
1:A:361:TYR:HB3	1:B:58:TYR:CG	2.38	0.58
1:B:2:PRO:N	1:B:27:SER:HG	2.01	0.57
1:B:127:THR:HG21	1:B:235:GLU:OE1	2.07	0.55
1:B:379:ARG:HH22	1:B:438:LYS:H	1.53	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:985:HOH:O	1:B:404:MET:HG3	2.06	0.54
1:B:97:LYS:CE	5:B:998:HOH:O	2.57	0.53
1:B:97:LYS:HE3	5:B:998:HOH:O	2.09	0.52
1:B:353:VAL:CG1	1:B:436:LYS:HD3	2.40	0.52
1:A:355:GLN:HG3	5:A:647:HOH:O	2.10	0.52
1:B:353:VAL:HG11	1:B:432[B]:MET:HB3	1.93	0.51
1:B:362:TYR:CG	1:B:363:PRO:HD2	2.46	0.51
1:B:150:LYS:HG2	1:B:230:TYR:CD2	2.47	0.49
1:B:3:LYS:HE2	1:B:30[A]:ILE:HD11	1.95	0.49
1:A:169:GLU:OE1	5:A:830:HOH:O	2.20	0.49
1:A:203:GLU:O	5:A:1008:HOH:O	2.20	0.49
1:B:362:TYR:CD2	1:B:363:PRO:HD2	2.47	0.49
1:A:159[B]:VAL:CG1	2:A:501:FAD:HM73	2.44	0.48
1:B:393:ASP:OD2	5:B:959:HOH:O	2.20	0.48
1:A:181:ASP:OD2	5:A:820:HOH:O	2.20	0.48
1:A:42:ASN:H	1:A:42:ASN:ND2	2.11	0.47
1:A:255:ARG:NH1	1:A:256:LYS:HE3	2.29	0.47
1:A:159[B]:VAL:HG11	2:A:501:FAD:HM73	1.97	0.46
1:B:355:GLN:NE2	1:B:432[B]:MET:SD	2.89	0.46
1:A:361:TYR:HB3	1:B:58:TYR:CD2	2.51	0.46
1:A:226:LYS:HE2	1:A:226:LYS:HB2	1.82	0.45
1:B:253:LEU:HD23	1:B:259:ILE:HG12	1.99	0.45
1:A:341:ASN:ND2	5:A:722:HOH:O	2.20	0.44
1:A:429:LEU:H	1:A:429:LEU:HD12	1.82	0.44
1:B:123:GLU:O	1:B:123:GLU:CG	2.64	0.43
1:B:246:ILE:HD13	1:B:259:ILE:HD11	2.00	0.43
1:A:164:LEU:HD23	1:A:164:LEU:C	2.39	0.43
1:B:3:LYS:HE2	1:B:30[A]:ILE:CD1	2.49	0.43
1:B:353:VAL:HG12	1:B:436:LYS:HD3	2.01	0.42
1:A:44:ALA:HB2	1:B:361:TYR:CE1	2.54	0.42
1:A:4[B]:ILE:HD13	1:A:20:ILE:HD13	2.02	0.42
1:A:44:ALA:HB2	1:B:361:TYR:CZ	2.55	0.42
1:B:436:LYS:NZ	5:B:928:HOH:O	2.50	0.42
1:A:148:VAL:HG13	1:A:231:ASP:HB2	2.02	0.42
1:A:436:LYS:HB3	1:A:436:LYS:HE2	1.91	0.41
1:A:56:ARG:HG2	1:A:56:ARG:O	2.20	0.41
1:B:30[A]:ILE:HG23	1:B:30[A]:ILE:HD13	1.78	0.41
1:A:150:LYS:CE	5:A:1032:HOH:O	2.45	0.41
1:A:255:ARG:HG2	5:A:738:HOH:O	2.20	0.41
1:B:164:LEU:CD1	1:B:176:LEU:HB2	2.51	0.41
1:B:113:PRO:O	2:B:501:FAD:H8A	2.21	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:975:HOH:O	5:A:998:HOH:O[1_545]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	445/437 (102%)	430 (97%)	14 (3%)	1 (0%)	47 30
1	B	446/437 (102%)	436 (98%)	9 (2%)	1 (0%)	47 30
All	All	891/874 (102%)	866 (97%)	23 (3%)	2 (0%)	47 30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	10	VAL
1	B	10	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	386/376 (103%)	374 (97%)	12 (3%)	40 21
1	B	387/376 (103%)	379 (98%)	8 (2%)	53 36
All	All	773/752 (103%)	753 (97%)	20 (3%)	47 28

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	GLU
1	A	42	ASN
1	A	55	ASP
1	A	58	TYR
1	A	67	PHE
1	A	70	ARG
1	A	160	SER
1	A	181	ASP
1	A	244	LYS
1	A	323	ASN
1	A	340	PRO
1	A	421	PRO
1	B	30[A]	ILE
1	B	30[B]	ILE
1	B	42	ASN
1	B	88	ARG
1	B	216	ASN
1	B	323	ASN
1	B	341	ASN
1	B	421	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	ASN
1	A	141	GLN
1	A	242	ASN
1	A	250	ASN
1	A	308	GLN
1	A	323	ASN
1	A	324	ASN
1	A	355	GLN
1	A	380	GLN
1	B	42	ASN
1	B	85	ASN
1	B	99	ASN
1	B	209	ASN
1	B	216	ASN
1	B	218	ASN
1	B	242	ASN
1	B	250	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	308	GLN
1	B	323	ASN
1	B	324	ASN
1	B	341	ASN
1	B	355	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	501	-	51,58,58	1.59	11 (21%)	60,89,89	2.29	12 (20%)
2	FAD	B	501	-	51,58,58	1.68	9 (17%)	60,89,89	2.42	18 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	501	-	-	2/30/50/50	0/6/6/6
2	FAD	B	501	-	-	2/30/50/50	0/6/6/6

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	FAD	C2A-N3A	6.60	1.42	1.32
2	A	501	FAD	C5'-C4'	4.32	1.57	1.51
2	B	501	FAD	C5X-N5	3.87	1.41	1.35
2	B	501	FAD	C2A-N1A	3.70	1.40	1.33
2	B	501	FAD	C4X-N5	3.70	1.38	1.33
2	A	501	FAD	C2A-N3A	3.33	1.37	1.32
2	A	501	FAD	C4X-N5	3.32	1.38	1.33
2	A	501	FAD	C2A-N1A	3.12	1.39	1.33
2	B	501	FAD	C4A-N3A	3.02	1.39	1.35
2	B	501	FAD	C4-N3	2.82	1.37	1.33
2	A	501	FAD	O2'-C2'	-2.76	1.37	1.43
2	B	501	FAD	C1'-N10	2.72	1.51	1.48
2	A	501	FAD	C4-N3	2.56	1.37	1.33
2	A	501	FAD	C8M-C8	2.55	1.56	1.51
2	A	501	FAD	C5X-N5	2.51	1.39	1.35
2	A	501	FAD	C9-C8	2.37	1.43	1.37
2	B	501	FAD	C9A-N10	2.28	1.41	1.38
2	A	501	FAD	C8A-N7A	2.25	1.38	1.34
2	B	501	FAD	C9-C8	2.17	1.43	1.37
2	A	501	FAD	C4X-C10	2.10	1.40	1.38

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	FAD	C4-N3-C2	10.12	123.69	115.14
2	B	501	FAD	C4-N3-C2	9.29	122.99	115.14
2	B	501	FAD	C4X-C4-N3	-6.53	114.50	123.43
2	A	501	FAD	C10-C4X-N5	-6.27	116.92	121.26
2	A	501	FAD	C4X-C4-N3	-5.60	115.77	123.43
2	A	501	FAD	C4X-N5-C5X	5.35	122.12	116.77
2	B	501	FAD	C5X-C9A-N10	4.58	121.03	117.72
2	A	501	FAD	N3A-C2A-N1A	-4.45	121.72	128.68
2	B	501	FAD	C1'-N10-C9A	4.35	121.72	118.29
2	B	501	FAD	C9A-N10-C10	-4.33	116.24	121.91
2	B	501	FAD	N3A-C2A-N1A	-4.08	122.30	128.68
2	B	501	FAD	C1'-N10-C10	4.06	122.04	118.41
2	B	501	FAD	C6-C5X-N5	-3.82	114.84	119.05

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	FAD	O4B-C1B-C2B	-3.64	101.61	106.93
2	B	501	FAD	C6-C5X-C9A	3.62	123.79	119.05
2	A	501	FAD	O4B-C1B-C2B	-3.21	102.24	106.93
2	B	501	FAD	C2A-N1A-C6A	2.89	123.71	118.75
2	A	501	FAD	C4-C4X-N5	2.80	121.79	118.60
2	B	501	FAD	C9-C8-C7	2.63	124.35	119.91
2	A	501	FAD	C5A-C6A-N6A	2.53	124.20	120.35
2	A	501	FAD	C1'-N10-C9A	2.32	120.12	118.29
2	B	501	FAD	C5A-C6A-N1A	-2.24	115.28	120.35
2	B	501	FAD	O3B-C3B-C4B	-2.24	104.58	111.05
2	A	501	FAD	C1B-N9A-C4A	-2.23	122.72	126.64
2	B	501	FAD	P-O3P-PA	-2.13	125.50	132.83
2	B	501	FAD	N6A-C6A-N1A	2.08	122.89	118.57
2	B	501	FAD	O2A-PA-O1A	2.05	122.38	112.24
2	A	501	FAD	C7-C6-C5X	-2.05	118.32	121.22
2	B	501	FAD	C1B-N9A-C4A	-2.01	123.11	126.64
2	A	501	FAD	C4-C4X-C10	2.00	121.28	119.95

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	FAD	O4B-C4B-C5B-O5B
2	B	501	FAD	O4B-C4B-C5B-O5B
2	A	501	FAD	C3B-C4B-C5B-O5B
2	B	501	FAD	C3B-C4B-C5B-O5B


There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	FAD	2	0
2	B	501	FAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	437/437 (100%)	-0.13	7 (1%) 72 76	14, 19, 35, 62	4 (0%)
1	B	437/437 (100%)	-0.18	6 (1%) 75 79	13, 20, 32, 49	3 (0%)
All	All	874/874 (100%)	-0.16	13 (1%) 73 77	13, 20, 33, 62	7 (0%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	58	TYR	7.5
1	A	55	ASP	4.1
1	B	438	LYS	3.8
1	A	60	LEU	3.2
1	A	57	ARG	2.8
1	B	123	GLU	2.6
1	B	360	ASN	2.6
1	A	123	GLU	2.4
1	B	58	TYR	2.3
1	B	216	ASN	2.1
1	B	54	GLU	2.1
1	A	425	HIS	2.1
1	A	54	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	A	502	1/1	0.93	0.11	37,37,37,37	0
4	CL	B	502	1/1	0.96	0.09	30,30,30,30	0
4	CL	A	505	1/1	0.97	0.16	45,45,45,45	0
4	CL	A	504	1/1	0.97	0.05	27,27,27,27	0
2	FAD	B	501	53/53	0.97	0.07	12,15,18,21	0
4	CL	B	503	1/1	0.97	0.19	41,41,41,41	0
2	FAD	A	501	53/53	0.97	0.07	12,15,17,18	0
3	MG	A	503	1/1	0.99	0.09	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.