



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 12, 2022 – 06:12 PM JST

PDB ID : 7ER8
Title : Crystal structure of human Biliverdin IX-beta reductase B with Sulfasalazine (SAS)
Authors : Griesinger, C.; Lee, D.; Ryu, K.S.; Kim, M.; Ha, J.H.
Deposited on : 2021-05-06
Resolution : 1.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.25
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.25


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	<div> <div>3%</div> <div>87%</div> <div>9%</div> <div>.</div> </div>
1	B	216	<div> <div>5%</div> <div>85%</div> <div>9%</div> <div>5%</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3700 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Flavin reductase (NADPH).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	S	0	4	0
			1584	993	282	301	8			
1	B	206	Total	C	N	O	S	0	1	0
			1559	978	279	294	8			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	HIS	-	expression tag	UNP P30043
A	-8	HIS	-	expression tag	UNP P30043
A	-7	HIS	-	expression tag	UNP P30043
A	-6	HIS	-	expression tag	UNP P30043
A	-5	HIS	-	expression tag	UNP P30043
A	-4	HIS	-	expression tag	UNP P30043
A	-3	HIS	-	expression tag	UNP P30043
A	-2	HIS	-	expression tag	UNP P30043
A	-1	HIS	-	expression tag	UNP P30043
A	0	HIS	-	expression tag	UNP P30043
B	-9	HIS	-	expression tag	UNP P30043
B	-8	HIS	-	expression tag	UNP P30043
B	-7	HIS	-	expression tag	UNP P30043
B	-6	HIS	-	expression tag	UNP P30043
B	-5	HIS	-	expression tag	UNP P30043
B	-4	HIS	-	expression tag	UNP P30043
B	-3	HIS	-	expression tag	UNP P30043
B	-2	HIS	-	expression tag	UNP P30043
B	-1	HIS	-	expression tag	UNP P30043
B	0	HIS	-	expression tag	UNP P30043

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃) (labeled as "Ligand of Interest" by depositor).



- # SAS
- 
- The chemical structure represents the sodium salt of 4-((4-aminophenyl)diazenyl)-2-hydroxybenzoic acid. It features three aromatic rings: a top benzene ring with a carboxylate group (C18=O4, C18-O5, HO) and a hydroxyl group (C15-O3, OH); a middle benzene ring connected via an azo group (N4=N3); and a bottom benzene ring with an amino group (N1=N2, NH). A sodium cation (Na+) is shown as a yellow sphere bonded to the carboxylate oxygen (O5) and the amino nitrogen (N2). Atoms are color-coded: carbon (grey), oxygen (red), nitrogen (blue), and sodium (yellow). Labels C1 through C18 identify the carbon atoms, and O1 through O5 identify the oxygen atoms. The sodium atom is labeled Na.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			28	18	4	5	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	S	0	0
			28	18	4	5	1		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	213	Total	O	0	0
			213	213		
5	B	167	Total	O	0	0
			167	167		

i

- Molecule 1: Flavin reductase (NADPH)



- Molecule 1: Flavin reductase (NADPH)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	82.16Å 117.37Å 41.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.75 – 1.45 47.75 – 1.45	Depositor EDS
% Data completeness (in resolution range)	99.3 (47.75-1.45) 99.3 (47.75-1.45)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.49 (at 1.45Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.166 , 0.199 0.166 , 0.199	Depositor DCC
R_{free} test set	2000 reflections (2.76%)	wwPDB-VP
Wilson B-factor (Å ²)	14.0	Xtriage
Anisotropy	0.279	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 50.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3700	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, SAS, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/1628	0.87	3/2217 (0.1%)
1	B	0.55	0/1593	0.71	3/2169 (0.1%)
All	All	0.61	0/3221	0.79	6/4386 (0.1%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	135	MET	CG-SD-CE	-6.66	89.54	100.20
1	A	134	ARG	NE-CZ-NH1	6.15	123.38	120.30
1	B	134	ARG	NE-CZ-NH1	5.98	123.29	120.30
1	A	134	ARG	NE-CZ-NH2	-5.90	117.35	120.30
1	B	134	ARG	NE-CZ-NH2	-5.72	117.44	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1584	0	1597	18	0
1	B	1559	0	1578	19	0
2	A	48	0	25	1	0
2	B	48	0	25	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	28	0	13	6	0
3	B	28	0	13	2	0
4	A	10	0	0	0	0
4	B	15	0	0	0	0
5	A	213	0	0	6	4
5	B	167	0	0	9	4
All	All	3700	0	3251	39	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 39 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:46:ARG:NH2	5:B:501:HOH:O	1.76	1.14
1:A:170:ARG:NH2	5:A:505:HOH:O	2.26	0.68
1:B:126[A]:GLN:OE1	5:B:502:HOH:O	2.15	0.65
1:B:120:LYS:NZ	5:B:509:HOH:O	2.32	0.63
1:A:118:PRO:O	5:A:501:HOH:O	2.15	0.62

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:648:HOH:O	5:B:593:HOH:O[2_454]	1.88	0.32
5:A:569:HOH:O	5:B:568:HOH:O[2_455]	1.97	0.23
5:A:522:HOH:O	5:B:593:HOH:O[2_454]	2.02	0.18
5:A:622:HOH:O	5:B:638:HOH:O[2_455]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	209/216 (97%)	205 (98%)	3 (1%)	1 (0%)	29	9
1	B	205/216 (95%)	201 (98%)	3 (2%)	1 (0%)	29	9
All	All	414/432 (96%)	406 (98%)	6 (1%)	2 (0%)	29	9

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	152	PRO
1	B	152	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/179 (97%)	173 (99%)	1 (1%)	86	69
1	B	170/179 (95%)	168 (99%)	2 (1%)	71	43
All	All	344/358 (96%)	341 (99%)	3 (1%)	78	57

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	111	SER
1	B	111	SER
1	B	204	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	SO4	B	403	-	4,4,4	0.18	0	6,6,6	0.31	0
4	SO4	A	403	-	4,4,4	0.14	0	6,6,6	0.16	0
3	SAS	A	402	-	28,30,30	1.82	6 (21%)	37,42,42	4.32	16 (43%)
2	NAP	A	401	-	45,52,52	3.01	9 (20%)	56,80,80	2.16	16 (28%)
2	NAP	B	401	-	45,52,52	4.01	11 (24%)	56,80,80	2.09	11 (19%)
4	SO4	B	405	-	4,4,4	0.14	0	6,6,6	0.19	0
3	SAS	B	402	-	28,30,30	1.99	6 (21%)	37,42,42	2.61	6 (16%)
4	SO4	B	404	-	4,4,4	0.14	0	6,6,6	0.24	0
4	SO4	A	404	-	4,4,4	0.15	0	6,6,6	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	401	-	-	7/31/67/67	0/5/5/5
3	SAS	B	402	-	-	0/16/20/20	0/3/3/3
2	NAP	B	401	-	-	7/31/67/67	0/5/5/5
3	SAS	A	402	-	-	4/16/20/20	0/3/3/3

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	NAP	C2D-C1D	-13.76	1.32	1.53
2	B	401	NAP	O4D-C1D	13.08	1.59	1.41
2	B	401	NAP	O4B-C1B	12.86	1.59	1.41
2	A	401	NAP	O4D-C1D	10.93	1.56	1.41
2	A	401	NAP	C2D-C1D	-9.30	1.39	1.53

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	SAS	O2-S1-O1	-18.81	96.43	119.55
3	B	402	SAS	O2-S1-O1	-12.92	103.67	119.55
3	A	402	SAS	C6-S1-N2	9.36	118.61	106.83
3	A	402	SAS	C11-C6-S1	-7.88	111.20	119.77
2	A	401	NAP	O7N-C7N-C3N	6.62	127.55	119.63

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	NAP	O4D-C1D-N1N-C2N
2	A	401	NAP	O4D-C1D-N1N-C6N
2	A	401	NAP	C2D-C1D-N1N-C2N
2	A	401	NAP	C2D-C1D-N1N-C6N
2	B	401	NAP	O4D-C1D-N1N-C2N


There are no ring outliers.


4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	SAS	6	0
2	A	401	NAP	1	0
2	B	401	NAP	3	0
3	B	402	SAS	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.








Ligand SAS A 402





Ligand NAP A 401














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	207/216 (95%)	-0.14	6 (2%) 51 53	8, 15, 42, 68	0
1	B	206/216 (95%)	0.06	10 (4%) 29 32	13, 25, 47, 88	0
All	All	413/432 (95%)	-0.04	16 (3%) 39 42	8, 20, 47, 88	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	44	GLY	6.3
1	B	206	GLN	4.9
1	A	116	TRP	4.9
1	B	1	MET	4.7
1	B	172	PRO	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	SO4	B	405	5/5	0.49	0.27	110,111,114,114	0
4	SO4	A	404	5/5	0.62	0.26	94,96,98,99	0
3	SAS	B	402	28/28	0.67	0.18	45,55,78,82	0
4	SO4	B	403	5/5	0.72	0.22	57,61,68,76	0
4	SO4	A	403	5/5	0.85	0.32	101,101,102,104	0
3	SAS	A	402	28/28	0.86	0.18	24,36,86,87	0
4	SO4	B	404	5/5	0.93	0.18	60,67,73,73	0
2	NAP	B	401	48/48	0.94	0.11	17,20,25,31	0
2	NAP	A	401	48/48	0.98	0.07	9,11,13,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around SO4 B 405:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SO4 A 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SAS B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SO4 B 403:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SO4 A 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SAS A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SO4 B 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around NAP B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAP A 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.