



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 14, 2020 – 10:22 pm BST

PDB ID : 3ET6  
Title : The crystal structure of the catalytic domain of a eukaryotic guanylate cyclase  
Authors : Winger, J.A.; Derbyshire, E.R.; Lamers, M.H.; Marletta, M.A.; Kuriyan, J.  
Deposited on : 2008-10-07  
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	190	<div> <div>7%</div> <div> <div></div> <div>58%</div> <div>33%</div> <div>6%</div> <div>••</div> </div> </div>
2	B	190	<div> <div>%</div> <div> <div></div> <div>76%</div> <div>17%</div> <div>••</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	B	7	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2999 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Soluble guanylyl cyclase beta.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	184	Total	As	C	N	O	S	0	0	0
			1430	2	894	250	271	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	467	SER	-	EXPRESSION TAG	UNP Q5YLC2
A	656	THR	-	EXPRESSION TAG	UNP Q5YLC2

- Molecule 2 is a protein called Soluble guanylyl cyclase beta.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	183	Total	As	C	N	O	S	0	0	0
			1430	3	895	249	270	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	467	SER	-	EXPRESSION TAG	UNP Q5YLC2
B	656	THR	-	EXPRESSION TAG	UNP Q5YLC2

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	25	Total	O	0	0
			25	25		
4	B	74	Total	O	0	0
			74	74		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Soluble guanylyl cyclase beta



#### • Molecule 2: Soluble guanylyl cyclase beta



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	123.68 Å 123.68 Å 62.82 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.00 – 2.55 28.01 – 2.55	Depositor EDS
% Data completeness (in resolution range)	(Not available) (28.00-2.55) 94.6 (28.01-2.55)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.73 (at 2.54 Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.172 , 0.215 0.162 , 0.206	Depositor DCC
$R_{free}$ test set	883 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.8	Xtriage
Anisotropy	0.676	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 89.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2999	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CAS, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	0/1436	0.68	0/1937
2	B	0.79	0/1426	0.76	0/1923
All	All	0.70	0/2862	0.72	0/3860

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1430	0	1402	68	0
2	B	1430	0	1396	30	0
3	A	15	0	0	1	0
3	B	25	0	0	0	0
4	A	25	0	0	0	0
4	B	74	0	0	0	0
All	All	2999	0	2798	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:638:THR:O	1:A:639:VAL:CG1	1.69	1.39
2:B:566:GLU:CB	2:B:567:PRO:HD3	1.73	1.18
1:A:628:ARG:HH21	1:A:628:ARG:HG3	1.05	1.16
1:A:638:THR:C	1:A:639:VAL:HG13	1.54	1.14
2:B:566:GLU:HB3	2:B:567:PRO:HD3	1.31	1.09
1:A:638:THR:C	1:A:639:VAL:CG1	2.11	1.06
2:B:566:GLU:CB	2:B:567:PRO:CD	2.41	0.99
1:A:638:THR:O	1:A:639:VAL:HG13	0.75	0.92
1:A:628:ARG:NH2	1:A:628:ARG:HG3	1.71	0.92
1:A:560:VAL:O	1:A:560:VAL:HG22	1.70	0.89
2:B:566:GLU:HB2	2:B:567:PRO:HD3	1.55	0.88
1:A:628:ARG:CG	1:A:628:ARG:HH21	1.87	0.87
2:B:566:GLU:HB2	2:B:567:PRO:CD	2.06	0.82
1:A:631:ILE:CG2	1:A:648:TYR:HB3	2.17	0.74
1:A:561:ALA:HA	1:A:567:PRO:HA	1.70	0.72
1:A:536:THR:HG22	1:A:537:VAL:N	2.04	0.72
1:A:638:THR:C	1:A:639:VAL:HG12	2.10	0.72
1:A:539:CYS:O	1:A:540:ASP:HB2	1.90	0.71
1:A:508:ARG:HB3	1:A:556:GLU:HG3	1.73	0.70
2:B:566:GLU:HB3	2:B:567:PRO:CD	2.15	0.67
1:A:490:ALA:HA	1:A:498:VAL:HG21	1.76	0.67
1:A:631:ILE:HG21	1:A:648:TYR:HB3	1.79	0.64
2:B:494:SER:HB3	2:B:497:GLU:HG3	1.79	0.64
1:A:639:VAL:O	1:A:640:LYS:HB2	1.97	0.63
2:B:625:SER:HB3	2:B:628:ARG:HD3	1.80	0.62
2:B:625:SER:CB	2:B:628:ARG:HD3	2.29	0.62
2:B:600:THR:OG1	2:B:642:LYS:HE3	2.00	0.61
1:A:547:LEU:HA	1:A:574:MET:HE1	1.83	0.61
1:A:609:GLU:HG3	1:A:649:LEU:HD21	1.83	0.61
1:A:560:VAL:O	1:A:560:VAL:CG2	2.42	0.61
1:A:592:CAS:CE2	2:B:525:ILE:HD13	2.31	0.60
1:A:639:VAL:HG23	1:A:639:VAL:O	2.02	0.60
1:A:550:ALA:HB2	1:A:574:MET:HE1	1.84	0.58
1:A:534:ASN:OD1	1:A:539:CYS:N	2.37	0.58
1:A:639:VAL:HG22	1:A:641:GLY:H	1.68	0.58
1:A:501:LEU:O	1:A:501:LEU:HD23	2.04	0.57
2:B:521:LYS:C	2:B:521:LYS:HD2	2.26	0.57
1:A:493:SER:HB2	1:A:498:VAL:HG23	1.86	0.56
1:A:550:ALA:HB2	1:A:574:MET:CE	2.36	0.56
1:A:630:GLU:OE2	1:A:632:ARG:NH2	2.41	0.54
1:A:539:CYS:SG	1:A:541:ASP:HB3	2.50	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:616:SER:HB2	1:A:645:MET:SD	2.50	0.52
1:A:550:ALA:CB	1:A:574:MET:HE1	2.39	0.52
2:B:653:LEU:C	2:B:653:LEU:HD23	2.30	0.52
2:B:473:HIS:O	2:B:578:PRO:HA	2.10	0.51
1:A:650:LEU:HD23	1:A:651:SER:O	2.09	0.51
2:B:494:SER:HB3	2:B:497:GLU:OE1	2.10	0.50
1:A:479:LEU:CD2	1:A:553:MET:HE3	2.42	0.50
1:A:653:LEU:HD12	1:A:653:LEU:H	1.77	0.50
2:B:488:GLU:O	2:B:492:ARG:HD2	2.12	0.50
1:A:638:THR:O	1:A:639:VAL:CB	2.50	0.49
1:A:627:GLU:O	1:A:628:ARG:HD2	2.11	0.49
2:B:652:PRO:O	2:B:653:LEU:HB3	2.12	0.48
1:A:571:ARG:NH1	1:A:605:GLU:OE2	2.46	0.48
1:A:477:THR:HG22	1:A:546:LEU:HD23	1.96	0.48
2:B:479:LEU:HD23	2:B:479:LEU:C	2.34	0.48
1:A:508:ARG:HB3	1:A:556:GLU:CG	2.43	0.48
1:A:584:VAL:CG2	2:B:523:GLU:HG3	2.44	0.47
1:A:562:SER:OG	1:A:563:SER:N	2.47	0.47
1:A:584:VAL:HG21	2:B:523:GLU:HG3	1.95	0.47
1:A:526:GLY:O	1:A:527:ASP:HB2	2.14	0.47
1:A:539:CYS:O	1:A:540:ASP:CB	2.59	0.47
1:A:546:LEU:HD12	1:A:546:LEU:O	2.14	0.47
1:A:549:PHE:CD1	1:A:553:MET:HE2	2.50	0.46
2:B:522:VAL:HG11	2:B:590:ARG:HH21	1.80	0.46
1:A:521:LYS:O	1:A:521:LYS:HD2	2.15	0.46
1:A:501:LEU:C	1:A:501:LEU:HD23	2.36	0.46
1:A:550:ALA:CB	1:A:574:MET:CE	2.93	0.46
1:A:646:ARG:HG3	1:A:647:THR:N	2.31	0.45
2:B:494:SER:HB3	2:B:497:GLU:CG	2.45	0.45
2:B:642:LYS:HZ3	2:B:642:LYS:HG3	1.58	0.45
2:B:526:GLY:O	2:B:527:ASP:CB	2.65	0.45
1:A:599:ASN:O	1:A:603:ARG:HD2	2.18	0.44
2:B:533:CYS:HB3	2:B:542:HIS:HB2	2.00	0.44
1:A:520:TYR:C	1:A:520:TYR:CD1	2.91	0.44
2:B:652:PRO:O	2:B:653:LEU:CB	2.66	0.44
2:B:566:GLU:HB2	2:B:567:PRO:HD2	1.92	0.43
1:A:582:GLY:O	1:A:592:CAS:N	2.51	0.43
1:A:645:MET:HG2	1:A:646:ARG:N	2.33	0.43
1:A:562:SER:O	1:A:563:SER:HB2	2.17	0.43
2:B:633:GLU:OE1	2:B:646:ARG:HD2	2.19	0.43
1:A:479:LEU:CD1	1:A:553:MET:HE3	2.49	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:526:GLY:O	2:B:527:ASP:CG	2.57	0.43
1:A:526:GLY:O	1:A:527:ASP:CB	2.67	0.43
1:A:478:VAL:HG22	1:A:601:ALA:CB	2.49	0.43
1:A:650:LEU:CD2	1:A:651:SER:O	2.67	0.42
1:A:605:GLU:HG3	1:A:606:SER:N	2.34	0.42
1:A:494:SER:HB3	1:A:497:GLU:CG	2.49	0.42
1:A:525:ILE:HD13	2:B:590:ARG:NH2	2.35	0.42
1:A:646:ARG:NH2	3:A:8:PO4:O1	2.53	0.42
1:A:627:GLU:C	1:A:628:ARG:HD2	2.40	0.41
1:A:515:GLU:O	1:A:517:PRO:HD3	2.21	0.41
1:A:476:ALA:HB1	1:A:533:CYS:O	2.21	0.40
1:A:613:ILE:HG22	1:A:615:ILE:HG22	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	178/190 (94%)	158 (89%)	17 (10%)	3 (2%)	9	11
2	B	176/190 (93%)	165 (94%)	11 (6%)	0	100	100
All	All	354/380 (93%)	323 (91%)	28 (8%)	3 (1%)	19	27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	639	VAL
1	A	652	PRO
1	A	541	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	155/160 (97%)	138 (89%)	17 (11%)	6	6
2	B	153/159 (96%)	135 (88%)	18 (12%)	5	5
All	All	308/319 (97%)	273 (89%)	35 (11%)	5	5

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	478	VAL
1	A	492	ARG
1	A	521	LYS
1	A	545	VAL
1	A	546	LEU
1	A	552	ARG
1	A	559	ARG
1	A	560	VAL
1	A	590	ARG
1	A	593	LEU
1	A	605	GLU
1	A	615	ILE
1	A	628	ARG
1	A	638	THR
1	A	639	VAL
1	A	646	ARG
1	A	653	LEU
2	B	467	SER
2	B	482	ASP
2	B	492	ARG
2	B	496	LEU
2	B	504	GLU
2	B	521	LYS
2	B	523	GLU
2	B	535	VAL
2	B	547	LEU
2	B	559	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	578	PRO
2	B	587	LYS
2	B	588	MET
2	B	590	ARG
2	B	603	ARG
2	B	615	ILE
2	B	638	THR
2	B	642	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	507	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	CAS	B	621	2	5,8,9	2.35	1 (20%)	1,9,11	0.58	0
2	CAS	B	592	2	5,8,9	1.79	1 (20%)	1,9,11	0.58	0
2	CAS	B	499	2	5,8,9	2.09	1 (20%)	1,9,11	1.28	0
1	CAS	A	499	1	5,8,9	2.26	1 (20%)	1,9,11	0.77	0
1	CAS	A	592	1	5,8,9	1.96	1 (20%)	1,9,11	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CAS	B	621	2	-	0/0/7/9	-
2	CAS	B	592	2	-	0/0/7/9	-
2	CAS	B	499	2	-	0/0/7/9	-
1	CAS	A	499	1	-	0/0/7/9	-
1	CAS	A	592	1	-	0/0/7/9	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	621	CAS	CA-N	-5.21	1.32	1.48
1	A	499	CAS	CA-N	-4.75	1.33	1.48
2	B	499	CAS	CA-N	-4.47	1.34	1.48
1	A	592	CAS	CA-N	-4.00	1.36	1.48
2	B	592	CAS	CA-N	-3.43	1.37	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	592	CAS	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PO4	B	3	-	4,4,4	0.78	0	6,6,6	0.66	0
3	PO4	B	7	-	4,4,4	0.83	0	6,6,6	0.46	0
3	PO4	A	5	-	4,4,4	0.72	0	6,6,6	1.08	0
3	PO4	A	2	-	4,4,4	0.83	0	6,6,6	0.42	0
3	PO4	B	4	-	4,4,4	0.75	0	6,6,6	0.62	0
3	PO4	A	8	-	4,4,4	0.60	0	6,6,6	0.70	0
3	PO4	B	6	-	4,4,4	0.80	0	6,6,6	0.63	0
3	PO4	B	1	-	4,4,4	0.89	0	6,6,6	0.40	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	8	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	182/190 (95%)	0.24	13 (7%) 16 19	52, 83, 131, 147	0
2	B	180/190 (94%)	-0.25	2 (1%) 80 85	24, 51, 107, 140	0
All	All	362/380 (95%)	-0.01	15 (4%) 37 44	24, 69, 124, 147	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	559	ARG	5.2
1	A	639	VAL	5.2
1	A	560	VAL	5.2
1	A	561	ALA	4.0
2	B	561	ALA	3.7
1	A	562	SER	3.6
1	A	563	SER	3.6
1	A	558	SER	3.2
1	A	653	LEU	3.0
1	A	492	ARG	2.9
1	A	474	PRO	2.6
1	A	567	PRO	2.5
2	B	566	GLU	2.3
1	A	628	ARG	2.1
1	A	557	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CAS	B	499	9/10	0.98	0.10	81,101,126,148	0
1	CAS	A	499	9/10	0.98	0.08	44,72,79,91	0
1	CAS	A	592	9/10	0.98	0.10	59,64,90,150	0
2	CAS	B	592	9/10	0.99	0.09	40,48,71,75	0
2	CAS	B	621	9/10	0.99	0.09	17,41,65,171	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PO4	B	7	5/5	0.56	0.48	153,155,157,161	0
3	PO4	B	3	5/5	0.85	0.18	118,118,119,121	0
3	PO4	B	6	5/5	0.85	0.23	110,118,122,123	0
3	PO4	A	5	5/5	0.87	0.17	95,118,123,125	0
3	PO4	B	4	5/5	0.88	0.19	104,115,118,123	0
3	PO4	A	8	5/5	0.90	0.32	101,103,109,150	0
3	PO4	A	2	5/5	0.91	0.14	121,122,127,127	0
3	PO4	B	1	5/5	0.93	0.12	86,91,108,108	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.