



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 14, 2020 – 10:38 pm BST

PDB ID : 3ETS  
Title : Crystal structure of a bacterial arylsulfate sulfotransferase catalytic intermediate with 4-methylumbelliferone bound in the active site  
Authors : Malojcic, G.; Owen, R.L.; Grimshaw, J.P.; Glockshuber, R.  
Deposited on : 2008-10-08  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	571	<div> <div>4%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div>.</div> </div> </div>
1	B	571	<div> <div>2%</div> <div> <div></div> <div>90%</div> <div>8%</div> <div>..</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	4MU	A	572	-	X	-	-

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 9152 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Arylsulfate sulfotransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	563	Total	C	N	O	S	0	3	0
			4456	2830	761	854	11			
1	B	564	Total	C	N	O	S	0	2	0
			4459	2832	762	854	11			

- Molecule 2 is 7-hydroxy-4-methyl-2H-chromen-2-one (three-letter code: 4MU) (formula: C<sub>10</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	10	3		
2	B	1	Total	C	O	0	0
			13	10	3		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	88	Total	O	0	0
			88	88		
4	B	93	Total	O	0	0
			93	93		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Arylsulfate sulfotransferase



- Molecule 1: Arylsulfate sulfotransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 1 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	181.83Å 181.83Å 100.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.43 – 2.40 38.43 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (38.43-2.40) 99.7 (38.43-2.40)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	0.13	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.55 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.3	Depositor
R, $R_{free}$	0.175 , 0.216 0.180 , 0.221	Depositor DCC
$R_{free}$ test set	1202 reflections (1.63%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.3	Xtriage
Anisotropy	0.376	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 31.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.029 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9152	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HS8, SO4, 4MU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.76	1/4559 (0.0%)	0.72	2/6185 (0.0%)
1	B	0.75	0/4559	0.72	0/6186
All	All	0.76	1/9118 (0.0%)	0.72	2/12371 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	418	CYS	CB-SG	6.59	1.93	1.82

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	193	ASP	CB-CG-OD1	5.36	123.13	118.30
1	A	172	ASP	CB-CG-OD1	5.36	123.12	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4456	0	4299	31	0
1	B	4459	0	4310	27	0
2	A	13	0	7	5	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	13	0	7	1	0
3	A	15	0	0	1	0
3	B	15	0	0	0	0
4	A	88	0	0	0	0
4	B	93	0	0	3	0
All	All	9152	0	8623	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:436:HS8:S	1:A:436:HS8:NE2	2.07	1.26
1:A:59:GLN:HG2	3:A:576:SO4:O4	1.93	0.69
1:B:159:ILE:H	1:B:159:ILE:HD12	1.58	0.68
1:A:282:ILE:HG22	1:A:283:ARG:HG3	1.81	0.62
1:A:48:GLU:CD	1:A:48:GLU:H	2.04	0.61
1:A:406:LEU:CD1	1:A:475:ILE:HD13	2.33	0.59
1:B:282:ILE:HG22	1:B:283:ARG:HG3	1.85	0.58
1:B:169:LEU:N	1:B:170:PRO:CD	2.69	0.55
1:A:359:SER:OG	1:A:372:SER:HB3	2.08	0.53
1:A:406:LEU:HD11	1:A:475:ILE:HD13	1.91	0.53
1:A:169:LEU:N	1:A:170:PRO:CD	2.72	0.53
1:A:223:VAL:HG13	1:A:254:SER:HB3	1.92	0.51
1:B:567:GLN:CG	4:B:586:HOH:O	2.59	0.49
1:B:145:SER:HB3	1:B:170:PRO:HB3	1.94	0.49
1:B:138:HIS:CE1	1:B:208:TYR:HE2	2.32	0.48
1:A:474:LYS:HB2	1:A:485:VAL:HG11	1.95	0.48
1:A:138:HIS:CE1	1:A:208:TYR:HE2	2.32	0.48
1:B:406:LEU:CD1	1:B:475:ILE:HD13	2.44	0.48
1:A:31:LEU:HA	1:A:31:LEU:HD12	1.67	0.47
1:A:380:LYS:HG2	1:A:388:LYS:HB2	1.95	0.47
1:B:416:ILE:HD13	1:B:427:SER:OG	2.14	0.47
1:B:470:PHE:CD1	1:B:470:PHE:N	2.83	0.47
1:B:286:ILE:HB	1:B:299:TRP:HB2	1.97	0.47
1:A:356:HIS:CE1	2:A:572:4MU:H6	2.50	0.46
1:B:155:LYS:HG3	1:B:156:ASN:OD1	2.15	0.46
1:A:470:PHE:CD1	1:A:470:PHE:N	2.82	0.45
1:A:230:GLU:OE1	1:A:239:GLU:OE2	2.34	0.45
1:A:252:HIS:NE2	2:A:572:4MU:O1'	2.36	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:THR:HG23	1:A:176:PHE:HB3	1.99	0.45
2:B:572:4MU:HM4	2:B:572:4MU:H5	1.76	0.45
1:A:410:ASP:OD2	1:A:414:LYS:HB2	2.17	0.45
1:A:407:LYS:HA	1:A:408:PRO:HD3	1.87	0.44
1:A:28:LEU:HD11	1:A:65:TYR:HB3	2.00	0.44
1:B:327:LEU:HA	1:B:328:ALA:HA	1.67	0.44
1:B:500:ILE:HG13	1:B:501:THR:HG23	2.00	0.44
1:B:48:GLU:H	1:B:48:GLU:CD	2.21	0.44
1:B:436:HS8:HE1	4:B:619:HOH:O	2.18	0.44
2:A:572:4MU:HM4	2:A:572:4MU:H5	1.66	0.43
1:A:212:ILE:HA	1:A:221:THR:O	2.17	0.43
1:A:435:GLN:C	1:A:436:HS8:HD1	2.22	0.43
1:A:31:LEU:HD13	1:A:67:GLY:HA2	2.00	0.43
1:B:159:ILE:HD12	1:B:159:ILE:N	2.29	0.43
1:B:154:ASP:OD1	1:B:155:LYS:N	2.52	0.43
1:A:356:HIS:CE1	2:A:572:4MU:C6	3.02	0.42
1:A:58:GLY:HA3	1:A:571:LYS:O	2.19	0.42
1:A:316:LEU:HA	1:A:316:LEU:HD23	1.88	0.42
1:B:223:VAL:HG13	1:B:254:SER:HB3	2.02	0.42
1:A:356:HIS:NE2	2:A:572:4MU:H6	2.34	0.42
1:B:228:TRP:CZ2	1:B:241:HIS:CD2	3.08	0.42
1:B:470:PHE:O	1:B:487:GLU:HA	2.20	0.42
1:B:29:VAL:HG23	1:B:70:ILE:HD11	2.02	0.42
1:B:380:LYS:HG2	1:B:388:LYS:HB2	2.01	0.41
1:A:70:ILE:HD13	1:A:81:VAL:HG11	2.03	0.41
1:B:306:ASP:HA	1:B:307:PRO:HD3	1.89	0.41
1:A:307:PRO:HA	1:A:354:TRP:CE3	2.56	0.41
1:B:483:GLN:O	1:B:485:VAL:HG13	2.20	0.40
1:A:301:LEU:CD1	1:A:357:VAL:HG21	2.52	0.40
1:B:301:LEU:CD1	1:B:357:VAL:HG21	2.52	0.40
1:B:470:PHE:HD1	1:B:470:PHE:N	2.19	0.40
1:B:477:GLU:HG3	4:B:612:HOH:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	561/571 (98%)	535 (95%)	26 (5%)	0	100	100
1	B	561/571 (98%)	535 (95%)	24 (4%)	2 (0%)	34	48
All	All	1122/1142 (98%)	1070 (95%)	50 (4%)	2 (0%)	47	62

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	171	PHE
1	B	502	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	470/482 (98%)	461 (98%)	9 (2%)	57	75
1	B	470/482 (98%)	460 (98%)	10 (2%)	53	72
All	All	940/964 (98%)	921 (98%)	19 (2%)	55	74

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	143	GLN
1	A	154	ASP
1	A	208	TYR
1	A	265	ARG
1	A	269	SER
1	A	403	SER
1	A	427	SER
1	A	485	VAL
1	A	503	ILE
1	B	31	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	143	GLN
1	B	154	ASP
1	B	160	LEU
1	B	208	TYR
1	B	265	ARG
1	B	308	LYS
1	B	327	LEU
1	B	427	SER
1	B	503	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	204	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	HS8	B	436	1	10,14,15	1.35	1 (10%)	4,20,22	2.85	2 (50%)
1	HS8	A	436	1	10,14,15	1.84	2 (20%)	4,20,22	1.99	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HS8	B	436	1	-	0/5/12/14	0/1/1/1
1	HS8	A	436	1	-	0/5/12/14	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	436	HS8	O3-S	4.05	1.49	1.42
1	B	436	HS8	O3-S	2.91	1.47	1.42
1	A	436	HS8	O1-S	2.64	1.47	1.42

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	436	HS8	O1-S-O3	-4.71	106.96	119.22
1	A	436	HS8	CB-CA-C	-2.98	105.89	111.47
1	B	436	HS8	CB-CA-C	-2.75	106.32	111.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	436	HS8	1	0
1	A	436	HS8	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	B	575	-	4,4,4	0.11	0	6,6,6	0.64	0
3	SO4	A	575	-	4,4,4	0.21	0	6,6,6	0.85	0
3	SO4	A	576	-	4,4,4	0.21	0	6,6,6	0.38	0
2	4MU	A	572	-	12,14,14	2.44	4 (33%)	16,20,20	2.94	11 (68%)
3	SO4	B	574	-	4,4,4	0.22	0	6,6,6	0.72	0
3	SO4	A	574	-	4,4,4	0.34	0	6,6,6	0.54	0
2	4MU	B	572	-	12,14,14	2.08	4 (33%)	16,20,20	2.51	8 (50%)
3	SO4	B	576	-	4,4,4	0.26	0	6,6,6	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	4MU	B	572	-	-	-	0/2/2/2
2	4MU	A	572	-	-	-	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	572	4MU	C4A-C8A	6.42	1.50	1.41
2	B	572	4MU	C4A-C8A	5.36	1.48	1.41
2	A	572	4MU	C3-C2	2.80	1.43	1.37
2	B	572	4MU	C4-C4A	2.76	1.48	1.42
2	A	572	4MU	O1'-C7	-2.59	1.31	1.37
2	B	572	4MU	C3-C2	2.45	1.42	1.37
2	A	572	4MU	C4-C4A	2.22	1.47	1.42
2	B	572	4MU	O1'-C7	-2.15	1.32	1.37

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	572	4MU	CM4-C4-C3	5.21	128.35	120.74
2	A	572	4MU	C5-C4A-C8A	4.67	121.90	116.50
2	B	572	4MU	C5-C4A-C8A	4.27	121.44	116.50
2	A	572	4MU	C8-C8A-C4A	-4.03	118.59	123.05
2	B	572	4MU	C8-C8A-C4A	-3.75	118.90	123.05
2	B	572	4MU	O1-C8A-C8	3.50	120.17	116.03
2	B	572	4MU	CM4-C4-C3	3.49	125.84	120.74
2	A	572	4MU	O1-C2-C3	3.42	123.63	119.27
2	A	572	4MU	CM4-C4-C4A	-3.23	116.08	121.08

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	572	4MU	C6-C5-C4A	-3.21	116.67	121.13
2	B	572	4MU	CM4-C4-C4A	-3.19	116.14	121.08
2	B	572	4MU	C6-C5-C4A	-3.13	116.77	121.13
2	A	572	4MU	C6-C7-C8	3.00	124.22	120.39
2	A	572	4MU	O1-C8A-C8	2.99	119.57	116.03
2	B	572	4MU	C6-C7-C8	2.60	123.72	120.39
2	A	572	4MU	C5-C4A-C4	-2.48	118.93	123.66
2	A	572	4MU	C3-C4-C4A	-2.24	115.57	117.78
2	A	572	4MU	C7-C8-C8A	-2.19	117.35	120.42
2	B	572	4MU	O1-C2-C3	2.06	121.89	119.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	576	SO4	1	0
2	A	572	4MU	5	0
2	B	572	4MU	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	562/571 (98%)	-0.29	22 (3%) 39 38	33, 47, 79, 141	0
1	B	563/571 (98%)	-0.42	13 (2%) 60 58	33, 47, 79, 139	0
All	All	1125/1142 (98%)	-0.36	35 (3%) 49 47	33, 47, 79, 141	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	331	GLY	7.5
1	A	329	HIS	6.4
1	A	1	ALA	4.6
1	B	329	HIS	4.3
1	A	338	PRO	4.3
1	A	333	GLN	3.7
1	B	1	ALA	3.6
1	A	334	ALA	3.1
1	B	155	LYS	2.9
1	A	335	LYS	2.8
1	A	317	ASP	2.8
1	B	411	ALA	2.7
1	B	327	LEU	2.7
1	A	330	ALA	2.7
1	B	425	GLU	2.6
1	B	156	ASN	2.6
1	B	332	GLN	2.6
1	B	412	ASN	2.6
1	B	154	ASP	2.4
1	A	332	GLN	2.4
1	A	328	ALA	2.4
1	A	425	GLU	2.4
1	B	414	LYS	2.4
1	A	357	VAL	2.4

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	331	GLY	2.3
1	B	159	ILE	2.3
1	A	156[A]	ASN	2.2
1	A	426	ASN	2.2
1	A	155	LYS	2.1
1	A	462	LEU	2.1
1	A	372	SER	2.1
1	A	318	ALA	2.1
1	A	461	ALA	2.0
1	A	414	LYS	2.0
1	A	478	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	HS8	A	436	14/15	0.98	0.20	39,43,47,50	0
1	HS8	B	436	14/15	0.99	0.19	37,42,46,47	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	4MU	B	572	13/13	0.90	0.29	45,50,55,56	10
2	4MU	A	572	13/13	0.91	0.25	48,53,60,65	7
3	SO4	A	575	5/5	0.94	0.11	91,95,97,99	0
3	SO4	B	574	5/5	0.96	0.13	63,75,78,79	0
3	SO4	A	576	5/5	0.96	0.10	69,74,79,79	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	A	574	5/5	0.97	0.11	70,74,80,82	0
3	SO4	B	575	5/5	0.97	0.16	94,95,97,97	0
3	SO4	B	576	5/5	0.98	0.06	61,75,76,78	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.