



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 07:24 pm BST

PDB ID : 6ETU
Title : Crystal structure of KDM4D with tetrazolhydrazide compound 7
Authors : Malecki, P.H.; Link, A.; Weiss, M.S.; Heinemann, U.
Deposited on : 2017-10-27
Resolution : 1.33 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1385 (1.36-1.32)
Clashscore	141614	1417 (1.36-1.32)
Ramachandran outliers	138981	1397 (1.36-1.32)
Sidechain outliers	138945	1397 (1.36-1.32)
RSRZ outliers	127900	1369 (1.36-1.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	<div> <div>3%</div> <div> <div></div> <div>80%</div> <div>15%</div> <div>• •</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	A	408	-	-	X	-
8	BWZ	A	411[B]	-	X	-	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 6695 atoms, of which 3112 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysine-specific demethylase 4D.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	330	Total	C	H	N	O	S	1	67	0
			6116	1976	3073	511	537	19			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			10	2	6	2		
5	A	1	Total	C	H	O	0	0
			10	2	6	2		
5	A	1	Total	C	H	O	0	0
			10	2	6	2		
5	A	1	Total	C	H	O	0	0
			10	2	6	2		
5	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	1
			5	4	1		

- Molecule 7 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Ni	0	0
			1	1		

- Molecule 8 is [[3-(2 {H}-1,2,3,4-tetrazol-5-yl)phenyl]carbonylamino]azanium (three-letter code: BWZ) (formula: C₈H₉N₆O).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	A	1	Total	C	H	N	O	0	1
			24	8	9	6	1		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	478	Total	O	0	42
			496	496		

- Molecule 1: Lysine-specific demethylase 4D



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	72.00 Å 72.00 Å 150.87 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.24 – 1.33 48.24 – 1.33	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.24-1.33) 99.8 (48.24-1.33)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.45 (at 1.33 Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.123 , 0.154 0.126 , 0.152	Depositor DCC
R_{free} test set	1107 reflections (1.20%)	wwPDB-VP
Wilson B-factor (Å ²)	13.2	Xtriage
Anisotropy	0.184	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 55.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	6695	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, ZN, CL, NA, BWZ, EDO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.24	9/3315 (0.3%)	1.15	11/4483 (0.2%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	68	GLU	CD-OE1	-7.15	1.17	1.25
1	A	208	GLU	CD-OE1	-6.82	1.18	1.25
1	A	207[A]	GLY	C-O	-6.37	1.13	1.23
1	A	207[B]	GLY	C-O	-6.37	1.13	1.23
1	A	194	GLU	CD-OE1	-6.28	1.18	1.25
1	A	196[A]	MET	CA-C	5.61	1.67	1.52
1	A	196[B]	MET	CA-C	5.61	1.67	1.52
1	A	302	ASP	CG-OD2	-5.47	1.12	1.25
1	A	213	TYR	CE1-CZ	-5.07	1.31	1.38

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	32[A]	ASP	CB-CG-OD2	-7.67	111.40	118.30
1	A	32[B]	ASP	CB-CG-OD2	-7.67	111.40	118.30
1	A	322	ASP	CB-CG-OD1	7.12	124.71	118.30
1	A	195	ASP	CB-CG-OD2	-6.24	112.68	118.30
1	A	32[A]	ASP	CB-CG-OD1	6.13	123.82	118.30
1	A	32[B]	ASP	CB-CG-OD1	6.13	123.82	118.30
1	A	322	ASP	CB-CG-OD2	-5.91	112.98	118.30
1	A	326	ARG	NE-CZ-NH2	-5.89	117.36	120.30
1	A	102	ARG	NE-CZ-NH2	-5.63	117.48	120.30
1	A	67[A]	SER	CA-C-O	5.41	131.46	120.10
1	A	67[B]	SER	CA-C-O	5.41	131.46	120.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3043	3073	3007	57	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	20	30	29	9	0
6	A	5	0	0	0	0
7	A	1	0	0	0	0
8	A	15	9	0	0	0
9	A	496	0	0	16	0
All	All	3583	3112	3036	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:48[B]:LEU:HD11	1:A:272:MET:CE	1.79	1.11
1:A:225[A]:ARG:HG2	1:A:228[A]:ARG:NH2	1.78	0.99
1:A:48[B]:LEU:HD11	1:A:272:MET:HE2	1.43	0.96
1:A:225[A]:ARG:HG2	1:A:228[A]:ARG:HH22	1.30	0.92
1:A:164[B]:LEU:CD1	1:A:324[B]:PHE:CE2	2.53	0.91
1:A:164[B]:LEU:HD11	1:A:324[B]:PHE:CE2	2.09	0.87
1:A:175[B]:VAL:HG13	9:A:808:HOH:O	1.77	0.85
1:A:164[B]:LEU:HD11	1:A:324[B]:PHE:CD2	2.13	0.84
1:A:48[B]:LEU:HD11	1:A:272:MET:HE3	1.61	0.81
1:A:164[B]:LEU:HD12	1:A:324[B]:PHE:CE2	2.16	0.80
1:A:206[A]:LEU:HD21	1:A:289[A]:CYS:SG	2.24	0.77
1:A:255[B]:LYS:HD2	5:A:408:EDO:H21	1.67	0.76
1:A:167:GLU:HG3	1:A:301[B]:ILE:HD11	1.67	0.74
1:A:196[B]:MET:CE	1:A:309:GLN:OE1	2.35	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:170[A]:VAL:HG22	1:A:171[A]:VAL:H	1.57	0.69
1:A:176[B]:ASN:HB3	9:A:515:HOH:O	1.92	0.69
1:A:196[B]:MET:HE1	1:A:309:GLN:OE1	1.93	0.68
1:A:48[B]:LEU:HD13	1:A:49:ALA:N	2.09	0.67
5:A:408:EDO:H11	9:A:576:HOH:O	1.97	0.65
1:A:156:HIS:HD2	9:A:850:HOH:O	1.79	0.64
1:A:255[B]:LYS:HD2	5:A:408:EDO:C2	2.28	0.62
1:A:206[A]:LEU:HD11	1:A:289[A]:CYS:SG	2.40	0.61
1:A:160[B]:ILE:HG12	9:A:533[B]:HOH:O	2.03	0.59
1:A:329[B]:GLN:OE1	1:A:332:ARG:HD2	2.03	0.58
1:A:204:LEU:HD13	1:A:210[A]:LYS:HZ1	1.68	0.58
1:A:255[B]:LYS:CD	5:A:408:EDO:H21	2.35	0.57
1:A:164[B]:LEU:CD1	1:A:324[B]:PHE:HE2	2.11	0.56
1:A:136:TYR:HB3	1:A:187[B]:THR:HG21	1.87	0.56
1:A:317:VAL:HG23	1:A:317:VAL:O	2.06	0.55
1:A:127:LYS:NZ	5:A:404:EDO:H22	2.22	0.55
1:A:170[B]:VAL:HG12	9:A:671[B]:HOH:O	2.05	0.55
5:A:408:EDO:H22	9:A:877[B]:HOH:O	2.07	0.54
1:A:255[B]:LYS:HE2	5:A:406:EDO:H22	1.89	0.54
1:A:136:TYR:CE2	1:A:138:ALA:HB2	2.43	0.53
1:A:196[B]:MET:HE3	1:A:309:GLN:OE1	2.07	0.52
1:A:165:GLU:HA	1:A:170[A]:VAL:O	2.10	0.52
1:A:123[A]:ARG:HH12	5:A:404:EDO:H12	1.75	0.52
1:A:320:SER:HB3	9:A:659:HOH:O	2.11	0.51
1:A:61[A]:GLU:HG3	1:A:62[A]:THR:HG23	1.94	0.50
1:A:167:GLU:CG	1:A:301[B]:ILE:HD11	2.40	0.50
1:A:176[A]:ASN:HB3	9:A:509:HOH:O	2.09	0.50
1:A:48[B]:LEU:CD1	1:A:272:MET:HE3	2.40	0.47
1:A:255[B]:LYS:HE3	9:A:743:HOH:O	2.14	0.47
1:A:160[B]:ILE:HD11	1:A:176[B]:ASN:OD1	2.15	0.46
1:A:118[A]:PHE:HE1	1:A:211:THR:HG1	1.59	0.46
1:A:176[A]:ASN:N	9:A:509:HOH:O	2.50	0.45
1:A:315:ALA:O	1:A:316:ARG:CB	2.65	0.45
1:A:130:ILE:HG12	9:A:765[A]:HOH:O	2.17	0.44
1:A:164[B]:LEU:HD12	1:A:324[B]:PHE:HE2	1.74	0.44
1:A:301[B]:ILE:HD12	9:A:563:HOH:O	2.18	0.44
1:A:255[A]:LYS:HE3	9:A:887:HOH:O	2.18	0.43
1:A:187[A]:THR:HB	1:A:284:ASN:HD22	1.85	0.41
1:A:170[A]:VAL:HG22	1:A:171[A]:VAL:N	2.29	0.41
1:A:317:VAL:HA	1:A:340:GLN:OE1	2.20	0.41
1:A:319:PHE:HB3	9:A:882:HOH:O	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:LYS:HZ2	5:A:404:EDO:H22	1.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	395/342 (116%)	391 (99%)	3 (1%)	1 (0%)	41 19

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	316	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	342/292 (117%)	331 (97%)	11 (3%)	39 7

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	PHE
1	A	118[A]	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	118[B]	PHE
1	A	136	TYR
1	A	170[A]	VAL
1	A	170[B]	VAL
1	A	210[A]	LYS
1	A	210[B]	LYS
1	A	316	ARG
1	A	318	THR
1	A	319	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	EDO	A	405	-	3,3,3	0.64	0	2,2,2	0.28	0
5	EDO	A	406	-	3,3,3	0.22	0	2,2,2	0.40	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	A	408	-	3,3,3	0.76	0	2,2,2	0.79	0
5	EDO	A	407	-	3,3,3	0.44	0	2,2,2	0.90	0
6	SO4	A	409[A]	-	4,4,4	0.42	0	6,6,6	0.56	0
5	EDO	A	404	-	3,3,3	0.56	0	2,2,2	1.19	0
8	BWZ	A	411[B]	7	16,16,16	2.72	6 (37%)	21,21,21	2.50	12 (57%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	405	-	-	0/1/1/1	-
5	EDO	A	406	-	-	1/1/1/1	-
5	EDO	A	408	-	-	1/1/1/1	-
8	BWZ	A	411[B]	7	-	4/10/10/10	0/2/2/2
5	EDO	A	404	-	-	1/1/1/1	-
5	EDO	A	407	-	-	1/1/1/1	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	411[B]	BWZ	NAO-NAN	-5.77	1.33	1.41
8	A	411[B]	BWZ	CAF-CAE	-5.61	1.33	1.48
8	A	411[B]	BWZ	NAD-NAB	-4.24	1.28	1.34
8	A	411[B]	BWZ	CAJ-CAL	-3.74	1.42	1.50
8	A	411[B]	BWZ	CAE-NAC	-2.96	1.30	1.33
8	A	411[B]	BWZ	NAC-NAA	-2.82	1.30	1.34

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	411[B]	BWZ	CAE-NAC-NAA	4.98	109.30	104.87
8	A	411[B]	BWZ	NAC-NAA-NAB	-4.53	106.57	109.53
8	A	411[B]	BWZ	NAC-CAE-NAD	-4.18	106.72	111.39
8	A	411[B]	BWZ	CAG-CAI-CAF	-3.22	116.51	120.56
8	A	411[B]	BWZ	CAH-CAJ-CAK	-3.09	115.58	119.24
8	A	411[B]	BWZ	CAL-NAN-NAO	-2.72	117.21	121.59
8	A	411[B]	BWZ	CAH-CAG-CAI	2.68	124.05	120.25
8	A	411[B]	BWZ	CAE-NAD-NAB	2.68	107.25	104.87
8	A	411[B]	BWZ	CAF-CAK-CAJ	2.55	123.85	121.09
8	A	411[B]	BWZ	CAK-CAF-CAE	2.28	123.40	120.05

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
8	A	411[B]	BWZ	CAF-CAE-NAC	2.21	127.86	124.12
8	A	411[B]	BWZ	CAI-CAF-CAE	-2.19	117.06	120.79

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	411[B]	BWZ	NAD-CAE-CAF-CAK
5	A	408	EDO	O1-C1-C2-O2
5	A	404	EDO	O1-C1-C2-O2
8	A	411[B]	BWZ	NAC-CAE-CAF-CAK
8	A	411[B]	BWZ	NAD-CAE-CAF-CAI
8	A	411[B]	BWZ	NAC-CAE-CAF-CAI
5	A	406	EDO	O1-C1-C2-O2
5	A	407	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	406	EDO	1	0
5	A	408	EDO	5	0
5	A	404	EDO	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	330/342 (96%)	-0.21	10 (3%)	50 57	9, 14, 31, 84	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	319	PHE	8.5
1	A	318	THR	6.5
1	A	317	VAL	5.3
1	A	316	ARG	4.4
1	A	335	LEU	4.2
1	A	313	GLY	4.1
1	A	315	ALA	3.5
1	A	338	ARG	2.9
1	A	170[A]	VAL	2.8
1	A	334[A]	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	EDO	A	406	4/4	0.84	0.11	37,44,47,47	0
5	EDO	A	407	4/4	0.88	0.14	41,49,52,54	0
5	EDO	A	408	4/4	0.93	0.16	18,26,32,33	0
8	BWZ	A	411[B]	15/15	0.94	0.09	12,15,20,22	24
5	EDO	A	404	4/4	0.95	0.14	36,44,44,45	0
4	NA	A	403	1/1	0.96	0.21	44,44,44,44	1
5	EDO	A	405	4/4	0.96	0.11	22,26,31,31	0
6	SO4	A	409[A]	5/5	0.96	0.06	20,21,21,22	5
3	CL	A	402	1/1	0.96	0.05	41,41,41,41	0
2	ZN	A	401	1/1	1.00	0.03	21,21,21,21	0
7	NI	A	410	1/1	1.00	0.10	10,10,10,10	1

6.5 Other polymers [i](#)

There are no such residues in this entry.