



wwPDB X-ray Structure Validation Summary Report ⓘ

May 21, 2020 – 02:59 am BST

PDB ID : 3EYB
Title : Structural and functional insights into the ligand binding domain of a non-duplicated RXR from the invertebrate chordate amphioxus
Authors : Tocchini-Valentini, G.D.; Rochel, N.; Moras, D.; Structural Proteomics in Europe (SPINE)
Deposited on : 2008-10-20
Resolution : 2.79 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	219	<div> <div>66%</div> <div>20%</div> <div>5%</div> <div>9%</div> </div>
1	B	219	<div> <div>59%</div> <div>27%</div> <div>5%</div> <div>9%</div> </div>
1	C	219	<div> <div>61%</div> <div>26%</div> <div>•</div> <div>9%</div> </div>
1	D	219	<div> <div>67%</div> <div>21%</div> <div>•</div> <div>9%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6196 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nuclear hormone receptor RXR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	199	Total	C	N	O	S	0	0	0
			1537	989	266	273	9			
1	B	199	Total	C	N	O	S	0	0	0
			1537	989	266	273	9			
1	C	199	Total	C	N	O	S	0	0	0
			1537	989	266	273	9			
1	D	199	Total	C	N	O	S	0	0	0
			1536	989	266	272	9			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	12	Total	O	0	0
			12	12		
2	B	13	Total	O	0	0
			13	13		
2	C	10	Total	O	0	0
			10	10		
2	D	14	Total	O	0	0
			14	14		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Nuclear hormone receptor RXR



• Molecule 1: Nuclear hormone receptor RXR



• Molecule 1: Nuclear hormone receptor RXR



• Molecule 1: Nuclear hormone receptor RXR


Chain D:

67%

21%

•

9%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.72Å 96.12Å 131.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.79 14.97 – 2.79	Depositor EDS
% Data completeness (in resolution range)	(Not available) (15.00-2.79) 90.3 (14.97-2.79)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.56 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.1	Depositor
R, R_{free}	0.202 , 0.257 0.253 , 0.307	Depositor DCC
R_{free} test set	1192 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	50.7	Xtriage
Anisotropy	1.085	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 35.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6196	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/1566	0.88	2/2116 (0.1%)
1	B	0.70	0/1566	0.82	2/2116 (0.1%)
1	C	0.76	2/1566 (0.1%)	0.82	4/2116 (0.2%)
1	D	0.74	0/1565	0.85	0/2116
All	All	0.74	2/6263 (0.0%)	0.85	8/8464 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	303	LYS	CE-NZ	14.87	1.86	1.49
1	C	303	LYS	CD-CE	5.48	1.65	1.51

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	303	LYS	CD-CE-NZ	-7.70	93.99	111.70
1	B	281	PRO	N-CA-CB	6.27	110.82	103.30
1	C	281	PRO	N-CA-CB	6.10	110.62	103.30
1	A	281	PRO	N-CA-CB	5.85	110.32	103.30
1	A	392	ASP	CB-CA-C	-5.46	99.49	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1537	0	1543	36	0
1	B	1537	0	1543	50	0
1	C	1537	0	1543	47	0
1	D	1536	0	1543	33	0
2	A	12	0	0	4	0
2	B	13	0	0	3	0
2	C	10	0	0	2	0
2	D	14	0	0	4	0
All	All	6196	0	6172	165	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 165 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:303:LYS:CE	1:C:303:LYS:NZ	1.86	1.37
1:C:476:ILE:H	1:C:476:ILE:HD13	1.05	1.14
1:B:476:ILE:HD13	1:B:476:ILE:H	1.15	1.06
1:B:422:ARG:HG3	1:B:422:ARG:HH11	1.15	1.04
1:C:407:PRO:HD2	1:C:422:ARG:NH1	1.74	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	195/219 (89%)	181 (93%)	12 (6%)	2 (1%)	15	44
1	B	195/219 (89%)	181 (93%)	11 (6%)	3 (2%)	10	33
1	C	195/219 (89%)	180 (92%)	12 (6%)	3 (2%)	10	33
1	D	195/219 (89%)	175 (90%)	19 (10%)	1 (0%)	29	61

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	780/876 (89%)	717 (92%)	54 (7%)	9 (1%)	13	39

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	362	HIS
1	C	362	HIS
1	C	477	ASP
1	D	362	HIS
1	B	351	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/191 (84%)	137 (86%)	23 (14%)	3	10
1	B	160/191 (84%)	137 (86%)	23 (14%)	3	10
1	C	160/191 (84%)	142 (89%)	18 (11%)	6	18
1	D	160/191 (84%)	145 (91%)	15 (9%)	8	26
All	All	640/764 (84%)	561 (88%)	79 (12%)	4	15

5 of 79 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	422	ARG
1	B	476	ILE
1	D	450	ARG
1	B	429	LEU
1	B	457	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	317	HIS
1	C	326	GLN
1	D	317	HIS
1	B	464	HIS
1	D	344	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.