



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 25, 2020 – 06:19 am BST

PDB ID : 5F2M  
Title : Tagatose-1,6-bisphosphate aldolase from *Streptococcus pyogenes* in complex with competitive inhibitor hexitol-1,6-bisphosphate  
Authors : LowKam, C.  
Deposited on : 2015-12-02  
Resolution : 1.92 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)
RSRZ outliers	127900	7793 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	<div> <div>9%</div> <div>96%</div> <div>..</div> </div>
1	B	327	<div> <div>%</div> <div>96%</div> <div>..</div> </div>
1	C	327	<div> <div>17%</div> <div>94%</div> <div>6%</div> </div>
1	D	327	<div> <div>11%</div> <div>93%</div> <div>6%</div> </div>

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 22167 atoms, of which 10303 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tagatose 1,6-diphosphate aldolase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	324	Total	C	H	N	O	S	0	5	0
			5123	1631	2552	424	507	9			
1	B	324	Total	C	H	N	O	S	0	6	0
			5123	1631	2552	424	508	8			
1	C	326	Total	C	H	N	O	S	0	6	0
			5205	1659	2593	431	513	9			
1	D	326	Total	C	H	N	O	S	0	12	0
			5237	1667	2606	436	518	10			

- Molecule 2 is D-MANNITOL-1,6-DIPHOSPHATE (three-letter code: M2P) (formula:  $C_6H_{16}O_{12}P_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	1
			40	12	24	4		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	4	Total Ca 4 4	0	0
3	A	3	Total Ca 3 3	0	0
3	D	2	Total Ca 2 2	0	0
3	C	1	Total Ca 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	395	Total O 395 395	0	0
4	B	495	Total O 495 495	0	0
4	C	248	Total O 248 248	0	0
4	D	291	Total O 291 291	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tagatose 1,6-diphosphate aldolase 2




- Molecule 1: Tagatose 1,6-diphosphate aldolase 2




- Molecule 1: Tagatose 1,6-diphosphate aldolase 2



- Molecule 1: Tagatose 1,6-diphosphate aldolase 2





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.98Å 107.84Å 237.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.20 – 1.92 45.20 – 1.92	Depositor EDS
% Data completeness (in resolution range)	91.6 (45.20-1.92) 91.2 (45.20-1.92)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.54 (at 1.92Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.159 , 0.207 0.163 , 0.208	Depositor DCC
$R_{free}$ test set	2000 reflections (1.67%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.8	Xtriage
Anisotropy	0.820	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 65.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	22167	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 34.49 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.7716e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, M2P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.47	0/2629	0.57	0/3550
1	B	0.48	0/2638	0.59	0/3561
1	C	0.38	0/2663	0.52	0/3594
1	D	0.40	0/2700	0.53	0/3643
All	All	0.44	0/10630	0.55	0/14348

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	25[B]	ALA	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2571	2552	2536	5	0
1	B	2571	2552	2522	6	0
1	C	2612	2593	2591	9	0
1	D	2631	2606	2582	14	0
2	A	40	0	24	0	0
3	A	3	0	0	0	0
3	B	4	0	0	0	0
3	C	1	0	0	0	0
3	D	2	0	0	0	0
4	A	395	0	0	3	1
4	B	495	0	0	3	2
4	C	248	0	0	1	1
4	D	291	0	0	5	2
All	All	11864	10303	10255	34	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 34 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:GLU:OE2	4:A:3101:HOH:O	2.03	0.75
1:C:156:GLU:OE2	4:C:501:HOH:O	2.05	0.73
1:D:276[B]:CYS:SG	4:D:733:HOH:O	2.47	0.72
1:D:84:GLU:O	4:D:501:HOH:O	2.09	0.70
1:B:7[B]:ASN:OD1	4:B:501:HOH:O	2.10	0.69

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:3220:HOH:O	4:B:614:HOH:O[1_655]	1.94	0.26
4:C:650:HOH:O	4:D:695:HOH:O[2_564]	2.01	0.19
4:B:880:HOH:O	4:D:538:HOH:O[3_645]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/327 (100%)	318 (97%)	9 (3%)	0	100	100
1	B	328/327 (100%)	317 (97%)	9 (3%)	2 (1%)	25	14
1	C	330/327 (101%)	314 (95%)	12 (4%)	4 (1%)	13	4
1	D	336/327 (103%)	327 (97%)	9 (3%)	0	100	100
All	All	1321/1308 (101%)	1276 (97%)	39 (3%)	6 (0%)	34	18

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	4	LEU
1	B	277	GLY
1	C	26[A]	PHE
1	C	26[B]	PHE
1	C	277	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/274 (100%)	271 (98%)	4 (2%)	65	61
1	B	275/274 (100%)	273 (99%)	2 (1%)	84	83
1	C	278/274 (102%)	273 (98%)	5 (2%)	59	53
1	D	281/274 (103%)	275 (98%)	6 (2%)	53	46
All	All	1109/1096 (101%)	1092 (98%)	17 (2%)	71	61

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	27[B]	ASP
1	C	29[A]	ARG
1	D	29[A]	ARG
1	C	27[A]	ASP
1	D	29[B]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 10 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	M2P	A	3001[B]	-	19,19,19	0.99	1 (5%)	28,28,28	0.76	0
2	M2P	A	3001[A]	-	19,19,19	0.99	2 (10%)	28,28,28	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	M2P	A	3001[B]	-	-	7/24/24/24	-
2	M2P	A	3001[A]	-	-	4/24/24/24	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	3001[B]	M2P	O5-C5	-2.19	1.38	1.43
2	A	3001[A]	M2P	O5-C5	-2.15	1.38	1.43
2	A	3001[A]	M2P	C3-C4	-2.05	1.49	1.53

There are no bond angle outliers.

There are no chirality outliers.

5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	3001[B]	M2P	O5-C5-C6-O6
2	A	3001[A]	M2P	C4-C5-C6-O6
2	A	3001[A]	M2P	O5-C5-C6-O6
2	A	3001[B]	M2P	O4-C4-C5-C6
2	A	3001[B]	M2P	C3-C4-C5-O5


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

## Ligand M2P A 3001 (B)





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	324/327 (99%)	0.35	30 (9%) 8 10	16, 28, 76, 115	2 (0%)
1	B	324/327 (99%)	0.04	2 (0%) 89 90	16, 26, 47, 96	1 (0%)
1	C	326/327 (99%)	0.91	57 (17%) 1 1	22, 45, 108, 338	4 (1%)
1	D	326/327 (99%)	0.51	35 (10%) 6 7	18, 42, 85, 121	1 (0%)
All	All	1300/1308 (99%)	0.45	124 (9%) 8 9	16, 35, 84, 338	8 (0%)

The worst 5 of 124 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1	THR	14.7
1	C	40	GLN	8.9
1	C	2	ILE	8.7
1	C	36	MET	8.6
1	C	44	PRO	8.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	B	403	1/1	0.59	0.14	86,86,86,86	0
3	CA	D	402	1/1	0.68	0.10	82,82,82,82	0
3	CA	A	3004	1/1	0.88	0.15	62,62,62,62	0
2	M2P	A	3001[B]	20/20	0.89	0.26	4,44,55,56	20
2	M2P	A	3001[A]	20/20	0.89	0.26	40,44,57,58	20
3	CA	C	401	1/1	0.94	0.23	62,62,62,62	0
3	CA	B	404	1/1	0.96	0.08	60,60,60,60	0
3	CA	D	401	1/1	0.96	0.07	32,32,32,32	0
3	CA	B	401	1/1	0.98	0.08	26,26,26,26	0
3	CA	B	402	1/1	0.99	0.09	28,28,28,28	0
3	CA	A	3003	1/1	1.00	0.07	25,25,25,25	0
3	CA	A	3002	1/1	1.00	0.10	27,27,27,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around M2P A 3001 (B):**




2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around M2P A 3001 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.