



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 20, 2022 – 06:04 PM JST

PDB ID : 7F5H  
Title : The crystal structure of RBD-Nanobody complex, DL28 (SC4)  
Authors : Luo, Z.P.; Li, T.; Lai, Y.; Zhou, Y.; Tan, J.; Li, D.  
Deposited on : 2021-06-22  
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.29
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.29


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	213	 84% 8% 8%
1	B	213	 83% 9% 8%
2	C	143	 71% 8% 20%
2	D	143	 76% 8% 16%
3	E	4	 25% 75%
3	F	4	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BMA	E	3	-	-	-	X
3	FUC	E	4	-	-	-	X
3	NAG	F	2	-	-	-	X
3	BMA	F	3	-	-	-	X
3	FUC	F	4	-	-	-	X
4	GOL	A	601	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5026 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SARS-CoV-2 Spike Receptor-Binding Domain (RBD).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	196	Total	C	N	O	S	0	0	0
			1550	994	259	289	8			
1	B	196	Total	C	N	O	S	0	0	0
			1550	994	259	289	8			


There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	327	ALA	-	expression tag	UNP P0DTC2
A	328	GLY	-	expression tag	UNP P0DTC2
A	329	SER	-	expression tag	UNP P0DTC2
A	532	GLY	-	expression tag	UNP P0DTC2
A	533	THR	-	expression tag	UNP P0DTC2
A	534	LEU	-	expression tag	UNP P0DTC2
A	535	GLU	-	expression tag	UNP P0DTC2
A	536	VAL	-	expression tag	UNP P0DTC2
A	537	LEU	-	expression tag	UNP P0DTC2
A	538	PHE	-	expression tag	UNP P0DTC2
A	539	GLN	-	expression tag	UNP P0DTC2
B	327	ALA	-	expression tag	UNP P0DTC2
B	328	GLY	-	expression tag	UNP P0DTC2
B	329	SER	-	expression tag	UNP P0DTC2
B	532	GLY	-	expression tag	UNP P0DTC2
B	533	THR	-	expression tag	UNP P0DTC2
B	534	LEU	-	expression tag	UNP P0DTC2
B	535	GLU	-	expression tag	UNP P0DTC2
B	536	VAL	-	expression tag	UNP P0DTC2
B	537	LEU	-	expression tag	UNP P0DTC2
B	538	PHE	-	expression tag	UNP P0DTC2
B	539	GLN	-	expression tag	UNP P0DTC2

- Molecule 2 is a protein called Nanobody DL28.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	114	Total	C	N	O	S	0	0	0
			868	536	151	177	4			
2	D	120	Total	C	N	O	S	0	0	0
			910	559	161	186	4			

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	4	Total	C	N	O	0	0	0
			49	28	2	19			
3	F	4	Total	C	N	O	0	0	0
			49	28	2	19			

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	9	Total	O	0	0
			9	9		
6	B	6	Total	O	0	0
			6	6		
6	C	2	Total	O	0	0
			2	2		
6	D	4	Total	O	0	0
			4	4		

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SARS-CoV-2 Spike Receptor-Binding Domain (RBD)

Chain A: 




- Molecule 1: SARS-CoV-2 Spike Receptor-Binding Domain (RBD)

Chain B: 



- Molecule 2: Nanobody DL28

Chain C: 



- Molecule 2: Nanobody DL28

Chain D: 



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:



HA01  
HA02  
BOA3  
FO04



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	177.46Å 177.46Å 133.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.81 – 3.00 47.81 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.81-3.00) 99.8 (47.81-3.00)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.54 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.226 , 0.248 0.228 , 0.254	Depositor DCC
$R_{free}$ test set	1307 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	83.5	Xtriage
Anisotropy	0.239	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 66.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5026	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	97.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, FUC, NAG, GOL, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	0/1594	0.76	0/2169
1	B	0.61	0/1594	0.77	0/2169
2	C	0.63	0/885	0.79	0/1198
2	D	0.64	0/927	0.82	0/1253
All	All	0.62	0/5000	0.78	0/6789

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1550	0	1467	8	0
1	B	1550	0	1467	13	0
2	C	868	0	816	6	0
2	D	910	0	854	10	0
3	E	49	0	43	0	0
3	F	49	0	43	0	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
4	C	6	0	8	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	6	0	8	1	0
5	B	5	0	0	0	0
6	A	9	0	0	0	0
6	B	6	0	0	0	0
6	C	2	0	0	0	0
6	D	4	0	0	0	0
All	All	5026	0	4722	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:483:VAL:HG21	2:D:63:LYS:HD3	1.46	0.96
1:B:483:VAL:CG2	2:D:63:LYS:HD3	2.08	0.83
1:A:335:LEU:HD23	1:A:362:VAL:HG23	1.60	0.82
1:B:483:VAL:HG21	2:D:63:LYS:CD	2.17	0.74
1:B:379:CYS:HB2	1:B:384:PRO:HG3	1.72	0.71
1:A:358:ILE:HB	1:A:395:VAL:HG13	1.78	0.63
1:B:412:PRO:HG3	1:B:429:PHE:HB3	1.81	0.62
1:B:350:VAL:HG11	1:B:402:ILE:HG23	1.84	0.59
1:B:483:VAL:HG21	2:D:63:LYS:CE	2.33	0.58
1:A:350:VAL:HG22	1:A:422:ASN:HB3	1.89	0.54
2:D:5:GLN:HB2	2:D:23:ALA:HB3	1.90	0.52
2:D:11:LEU:H	4:D:201:GOL:H31	1.75	0.51
2:C:27:SER:OG	2:C:28:ASP:N	2.42	0.51
1:A:412:PRO:HG3	1:A:429:PHE:HB3	1.92	0.50
2:C:7:SER:OG	2:C:21:SER:OG	2.30	0.50
2:D:35:GLY:HA3	2:D:50:ILE:HA	1.95	0.48
2:C:2:VAL:HA	2:C:26:GLY:HA3	1.95	0.48
2:D:86:PRO:HA	2:D:112:VAL:HB	1.95	0.47
1:B:418:ILE:HA	1:B:422:ASN:HB2	1.97	0.46
1:B:353:TRP:O	1:B:466:ARG:NH1	2.48	0.46
2:D:34:MET:HG2	2:D:96:VAL:HG12	1.97	0.46
1:B:379:CYS:HA	1:B:432:CYS:HA	1.99	0.45
1:A:341:VAL:HG23	1:A:342:PHE:HD1	1.81	0.45
1:B:428:ASP:OD1	1:B:428:ASP:N	2.49	0.44
2:C:31:SER:HB2	2:C:53:GLU:HG3	1.99	0.43
1:B:383:SER:HA	1:B:384:PRO:HD3	1.92	0.43
1:A:497:PHE:CG	1:A:507:PRO:HG3	2.54	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:89:THR:HG23	2:D:111:THR:HA	2.02	0.42
1:A:366:SER:OG	1:A:388:ASN:OD1	2.34	0.42
2:C:34:MET:HG2	2:C:96:VAL:HG22	2.01	0.42
1:A:401:VAL:HG22	1:A:509:ARG:HG2	2.02	0.40
1:B:358:ILE:HG23	1:B:395:VAL:HB	2.03	0.40
2:C:35:GLY:HA3	2:C:50:ILE:HA	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/213 (91%)	187 (96%)	7 (4%)	0	100	100
1	B	194/213 (91%)	186 (96%)	8 (4%)	0	100	100
2	C	112/143 (78%)	110 (98%)	2 (2%)	0	100	100
2	D	118/143 (82%)	117 (99%)	1 (1%)	0	100	100
All	All	618/712 (87%)	600 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/183 (92%)	166 (99%)	2 (1%)	71	90
1	B	168/183 (92%)	165 (98%)	3 (2%)	59	85
2	C	92/116 (79%)	92 (100%)	0	100	100
2	D	95/116 (82%)	95 (100%)	0	100	100
All	All	523/598 (88%)	518 (99%)	5 (1%)	76	91

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	347	PHE
1	A	377	PHE
1	B	357	ARG
1	B	377	PHE
1	B	524	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	E	1	3,1	14,14,15	0.46	0	17,19,21	2.10	2 (11%)
3	NAG	E	2	3	14,14,15	0.46	0	17,19,21	0.89	1 (5%)
3	BMA	E	3	3	11,11,12	0.27	0	15,15,17	1.22	1 (6%)
3	FUC	E	4	3	10,10,11	0.44	0	14,14,16	0.74	0
3	NAG	F	1	3,1	14,14,15	0.67	0	17,19,21	1.85	5 (29%)
3	NAG	F	2	3	14,14,15	0.32	0	17,19,21	0.90	0
3	BMA	F	3	3	11,11,12	0.30	0	15,15,17	0.88	1 (6%)
3	FUC	F	4	3	10,10,11	0.33	0	14,14,16	0.90	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	E	1	3,1	-	1/6/23/26	0/1/1/1
3	NAG	E	2	3	-	1/6/23/26	0/1/1/1
3	BMA	E	3	3	-	2/2/19/22	0/1/1/1
3	FUC	E	4	3	-	-	0/1/1/1
3	NAG	F	1	3,1	-	4/6/23/26	0/1/1/1
3	NAG	F	2	3	-	1/6/23/26	0/1/1/1
3	BMA	F	3	3	-	0/2/19/22	0/1/1/1
3	FUC	F	4	3	-	-	0/1/1/1

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	1	NAG	C1-O5-C5	6.77	121.37	112.19
3	E	1	NAG	O5-C5-C6	-3.97	100.99	107.20
3	F	1	NAG	C1-C2-N2	-3.85	103.90	110.49
3	F	1	NAG	C1-O5-C5	3.62	117.09	112.19
3	E	3	BMA	C1-O5-C5	3.52	116.97	112.19
3	F	1	NAG	O5-C5-C6	-2.97	102.55	107.20
3	F	1	NAG	C8-C7-N2	-2.92	111.16	116.10
3	F	1	NAG	O7-C7-N2	2.50	126.54	121.95
3	E	2	NAG	C4-C3-C2	-2.49	107.37	111.02
3	F	3	BMA	C1-O5-C5	2.39	115.43	112.19

There are no chirality outliers.


All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	E	3	BMA	O5-C5-C6-O6
3	F	1	NAG	C8-C7-N2-C2
3	E	3	BMA	C4-C5-C6-O6
3	F	1	NAG	O7-C7-N2-C2
3	E	1	NAG	C4-C5-C6-O6
3	F	1	NAG	C4-C5-C6-O6
3	F	2	NAG	C8-C7-N2-C2
3	F	1	NAG	C1-C2-N2-C7
3	E	2	NAG	C8-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	PO4	B	602	-	4,4,4	1.77	1 (25%)	6,6,6	0.32	0
4	GOL	C	201	-	5,5,5	0.81	0	5,5,5	0.40	0
4	GOL	D	201	-	5,5,5	0.54	0	5,5,5	0.87	0
4	GOL	B	601	-	5,5,5	0.11	0	5,5,5	0.37	0
4	GOL	A	601	-	5,5,5	0.32	0	5,5,5	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	C	201	-	-	0/4/4/4	-
4	GOL	D	201	-	-	0/4/4/4	-
4	GOL	B	601	-	-	0/4/4/4	-
4	GOL	A	601	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	602	PO4	P-O1	3.16	1.58	1.50

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	601	GOL	C1-C2-C3-O3
4	A	601	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	201	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	196/213 (92%)	-0.09	0 <b>100</b> <b>100</b>	59, 89, 148, 170	0
1	B	196/213 (92%)	-0.05	2 (1%) 82 59	64, 105, 154, 175	0
2	C	114/143 (79%)	-0.22	0 <b>100</b> <b>100</b>	57, 85, 114, 141	0
2	D	120/143 (83%)	-0.33	0 <b>100</b> <b>100</b>	61, 82, 127, 154	0
All	All	626/712 (87%)	-0.15	2 (0%) <b>94</b> <b>84</b>	57, 90, 146, 175	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	373	SER	2.1
1	B	333	THR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	BMA	E	3	11/12	<b>0.29</b>	<b>0.46</b>	138,186,202,206	0
3	BMA	F	3	11/12	<b>0.50</b>	<b>0.44</b>	133,176,192,202	0
3	FUC	F	4	10/11	<b>0.59</b>	<b>0.47</b>	149,181,195,200	0
3	FUC	E	4	10/11	<b>0.66</b>	<b>0.59</b>	174,194,216,221	0
3	NAG	E	1	14/15	0.72	0.33	135,171,210,230	0


*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	F	2	14/15	0.75	0.45	139,186,214,223	0
3	NAG	F	1	14/15	0.80	0.32	126,169,200,230	0
3	NAG	E	2	14/15	0.83	0.43	139,168,196,205	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	D	201	6/6	0.67	0.37	83,101,108,110	0
4	GOL	A	601	6/6	0.70	0.44	97,122,127,131	0
5	PO4	B	602	5/5	0.83	0.47	70,70,70,70	5
4	GOL	B	601	6/6	0.85	0.29	86,100,105,108	0
4	GOL	C	201	6/6	0.87	0.58	58,60,60,60	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.