



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 02:02 am BST

PDB ID : 5F7S  
Title : Cycloalternan-degrading enzyme from *Trueperella pyogenes*  
Authors : Light, S.H.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)  
Deposited on : 2015-12-08  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	733	<div> <div>2%</div> <div> <div></div> <div>91%</div> <div>5%</div> <div>.</div> </div> </div>
1	B	733	<div> <div>2%</div> <div> <div></div> <div>91%</div> <div>5%</div> <div>.</div> </div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 11919 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycoside hydrolase family 31.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	701	Total	C	N	O	S	0	0	0
			5608	3559	985	1045	19			
1	B	702	Total	C	N	O	S	0	0	0
			5618	3566	986	1047	19			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	730	GLY	-	expression tag	UNP X4QP62
A	731	ALA	-	expression tag	UNP X4QP62
A	732	ALA	-	expression tag	UNP X4QP62
A	733	LEU	-	expression tag	UNP X4QP62
B	730	GLY	-	expression tag	UNP X4QP62
B	731	ALA	-	expression tag	UNP X4QP62
B	732	ALA	-	expression tag	UNP X4QP62
B	733	LEU	-	expression tag	UNP X4QP62

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	3	Total	Ca	0	0
			3	3		
2	A	6	Total	Ca	0	0
			6	6		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula:  $C_6H_{14}O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			10	6	4		
4	B	1	Total	C	O	0	0
			10	6	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	320	Total 320	O 320	0	0
5	B	337	Total 337	O 337	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycoside hydrolase family 31



- Molecule 1: Glycoside hydrolase family 31



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	197.13Å 104.66Å 89.03Å 90.00° 90.74° 90.00°	Depositor
Resolution (Å)	30.00 – 2.30 29.67 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.00-2.30) 99.8 (29.67-2.30)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.49 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.170 , 0.211 0.175 , 0.213	Depositor DCC
$R_{free}$ test set	4135 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.2	Xtriage
Anisotropy	0.119	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 34.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.029 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	11919	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 56.20 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.8331e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, PEG, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.55	0/5788	0.75	2/7902 (0.0%)
1	B	0.57	0/5799	0.77	3/7917 (0.0%)
All	All	0.56	0/11587	0.76	5/15819 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	445	ARG	NE-CZ-NH2	-7.49	116.55	120.30
1	B	445	ARG	NE-CZ-NH1	6.60	123.60	120.30
1	A	445	ARG	NE-CZ-NH2	-6.04	117.28	120.30
1	B	541	ARG	NE-CZ-NH2	5.54	123.07	120.30
1	A	445	ARG	NE-CZ-NH1	5.04	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5608	0	5269	14	0
1	B	5618	0	5283	10	0
2	A	6	0	0	0	0
2	B	3	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	7	0	10	0	0
4	B	20	0	28	0	0
5	A	320	0	0	0	0
5	B	337	0	0	1	0
All	All	11919	0	10590	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:239:ASP:HB3	1:B:249:LEU:HB3	1.90	0.54
1:A:239:ASP:HB3	1:A:249:LEU:HB2	1.93	0.49
1:B:50:TRP:CD1	1:B:105:ALA:HB2	2.48	0.48
1:A:409:LEU:HG	1:A:415:VAL:CG1	2.45	0.47
1:A:50:TRP:CD1	1:A:105:ALA:HB2	2.50	0.47
1:A:352:GLN:HB3	1:A:353:PRO:HD2	1.99	0.44
1:A:515:SER:O	1:A:528:TYR:HA	2.18	0.44
1:A:150:SER:N	1:A:151:PRO:CD	2.82	0.43
1:A:436:MET:HE2	1:A:492:TYR:CZ	2.54	0.43
1:B:436:MET:HE2	1:B:492:TYR:CZ	2.54	0.42
1:A:129:SER:HA	1:A:155:VAL:O	2.19	0.42
1:A:514:PHE:CZ	1:A:529:TRP:HB3	2.55	0.42
1:B:330:GLU:HA	1:B:387:TRP:HB2	2.02	0.42
1:B:515:SER:O	1:B:528:TYR:HA	2.20	0.42
1:B:178:PHE:HA	1:B:213:MET:O	2.20	0.41
1:B:529:TRP:HA	1:B:559:GLY:O	2.21	0.41
1:A:455:LEU:HD23	1:A:459:LEU:HD12	2.03	0.41
1:A:35:ALA:HB3	1:A:44:MET:CE	2.51	0.41
1:B:117:ARG:NH1	5:B:901:HOH:O	2.40	0.41
1:A:431:PHE:N	1:A:432:PRO:HD3	2.36	0.40
1:B:563:ALA:HB1	1:B:570:PRO:HG3	2.02	0.40
1:A:151:PRO:HA	1:A:164:LYS:O	2.22	0.40
1:A:595:HIS:O	1:A:595:HIS:CG	2.74	0.40
1:B:150:SER:N	1:B:151:PRO:CD	2.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	695/733 (95%)	681 (98%)	12 (2%)	2 (0%)	41	50
1	B	698/733 (95%)	682 (98%)	15 (2%)	1 (0%)	51	64
All	All	1393/1466 (95%)	1363 (98%)	27 (2%)	3 (0%)	47	58

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	395	ASP
1	B	395	ASP
1	A	396	ASP

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	580/597 (97%)	575 (99%)	5 (1%)	78	89
1	B	581/597 (97%)	570 (98%)	11 (2%)	57	73
All	All	1161/1194 (97%)	1145 (99%)	16 (1%)	67	81

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	ARG
1	A	217	GLN
1	A	429	TRP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	565	PHE
1	A	589	GLN
1	B	15	ARG
1	B	192	LEU
1	B	217	GLN
1	B	224	ARG
1	B	260	LYS
1	B	349	ASN
1	B	429	TRP
1	B	527	LEU
1	B	565	PHE
1	B	589	GLN
1	B	644	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	319	ASN
1	B	349	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 9 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PGE	B	804	-	9,9,9	0.47	0	8,8,8	0.15	0
3	PEG	A	807	-	6,6,6	0.47	0	5,5,5	0.29	0
4	PGE	B	805	-	9,9,9	0.52	0	8,8,8	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PGE	B	804	-	-	2/7/7/7	-
3	PEG	A	807	-	-	1/4/4/4	-
4	PGE	B	805	-	-	5/7/7/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	805	PGE	O2-C3-C4-O3
4	B	804	PGE	O2-C3-C4-O3
4	B	804	PGE	O1-C1-C2-O2
4	B	805	PGE	O1-C1-C2-O2
4	B	805	PGE	O3-C5-C6-O4
4	B	805	PGE	C1-C2-O2-C3
3	A	807	PEG	O1-C1-C2-O2
4	B	805	PGE	C3-C4-O3-C5

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	701/733 (95%)	-0.17	17 (2%) 59 66	20, 34, 58, 100	0
1	B	702/733 (95%)	-0.14	16 (2%) 60 67	22, 34, 58, 95	0
All	All	1403/1466 (95%)	-0.16	33 (2%) 59 66	20, 34, 59, 100	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	429	TRP	4.9
1	B	429	TRP	4.6
1	B	144	ASP	3.6
1	B	397	ASP	3.5
1	B	51	PRO	3.3
1	A	256	PHE	3.2
1	B	395	ASP	3.2
1	A	397	ASP	3.1
1	B	52	GLU	3.1
1	A	52	GLU	2.9
1	B	222	ALA	2.8
1	A	350	GLN	2.8
1	B	386	LEU	2.8
1	A	430	TRP	2.7
1	B	53	HIS	2.7
1	A	99	HIS	2.6
1	B	328	VAL	2.5
1	A	386	LEU	2.5
1	B	396	ASP	2.4
1	B	54	GLU	2.4
1	A	98	PRO	2.3
1	B	385	ILE	2.3
1	A	257	GLU	2.3
1	B	394	THR	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	2.2
1	A	329	ILE	2.2
1	A	395	ASP	2.2
1	A	328	VAL	2.2
1	A	385	ILE	2.1
1	A	433	ASN	2.1
1	B	327	VAL	2.0
1	A	51	PRO	2.0
1	A	54	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CA	A	806	1/1	0.75	0.10	77,77,77,77	0
4	PGE	B	805	10/10	0.84	0.19	59,66,68,68	0
2	CA	B	802	1/1	0.86	0.10	74,74,74,74	0
3	PEG	A	807	7/7	0.90	0.17	51,55,57,58	0
4	PGE	B	804	10/10	0.92	0.16	46,47,51,52	0
2	CA	B	801	1/1	0.93	0.15	73,73,73,73	0
2	CA	A	805	1/1	0.96	0.10	55,55,55,55	0
2	CA	A	801	1/1	0.96	0.04	53,53,53,53	0
2	CA	A	803	1/1	0.96	0.06	49,49,49,49	0
2	CA	A	802	1/1	0.97	0.06	58,58,58,58	0
2	CA	B	803	1/1	0.98	0.10	59,59,59,59	0
2	CA	A	804	1/1	0.99	0.04	42,42,42,42	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.