



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2021 – 08:41 PM EDT

PDB ID : 3F9Z  
Title : Structural Insights into Lysine Multiple Methylation by SET Domain Methyltransferases, SET8-Y245F / H4-Lys20 / AdoHcy  
Authors : Couture, J.-F.; Dirk, L.M.A.; Brunzelle, J.S.; Houtz, R.L.; Trievel, R.C.  
Deposited on : 2008-11-14  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	166	<div> <div>7%</div> <div> <div></div> <div>83%</div> <div>11%</div> <div>6%</div> </div> </div>
1	B	166	<div> <div>5%</div> <div> <div></div> <div>89%</div> <div>7%</div> <div>...</div> </div> </div>
1	C	166	<div> <div>4%</div> <div> <div></div> <div>87%</div> <div>9%</div> <div>...</div> </div> </div>
1	D	166	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>12%</div> <div>.</div> </div> </div>
2	E	10	<div> <div></div> <div> <div></div> <div>70%</div> <div>20%</div> <div>10%</div> </div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	10	
2	G	10	
2	H	10	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6115 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-lysine N-methyltransferase SETD8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	156	Total	C	N	O	S	0	7	0
			1267	792	220	248	7			
1	B	161	Total	C	N	O	S	0	7	0
			1297	809	228	254	6			
1	C	160	Total	C	N	O	S	0	4	0
			1285	802	225	252	6			
1	D	161	Total	C	N	O	S	0	3	0
			1289	804	227	251	7			


There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	187	GLY	-	expression tag	UNP Q9NQR1
A	188	ALA	-	expression tag	UNP Q9NQR1
A	189	MET	-	expression tag	UNP Q9NQR1
A	190	GLY	-	expression tag	UNP Q9NQR1
A	245	PHE	TYR	engineered mutation	UNP Q9NQR1
B	187	GLY	-	expression tag	UNP Q9NQR1
B	188	ALA	-	expression tag	UNP Q9NQR1
B	189	MET	-	expression tag	UNP Q9NQR1
B	190	GLY	-	expression tag	UNP Q9NQR1
B	245	PHE	TYR	engineered mutation	UNP Q9NQR1
C	187	GLY	-	expression tag	UNP Q9NQR1
C	188	ALA	-	expression tag	UNP Q9NQR1
C	189	MET	-	expression tag	UNP Q9NQR1
C	190	GLY	-	expression tag	UNP Q9NQR1
C	245	PHE	TYR	engineered mutation	UNP Q9NQR1
D	187	GLY	-	expression tag	UNP Q9NQR1
D	188	ALA	-	expression tag	UNP Q9NQR1
D	189	MET	-	expression tag	UNP Q9NQR1
D	190	GLY	-	expression tag	UNP Q9NQR1
D	245	PHE	TYR	engineered mutation	UNP Q9NQR1

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	9	Total	C	N	O	0	0	0
			81	48	21	12			
2	F	8	Total	C	N	O	0	0	0
			72	44	20	8			
2	G	7	Total	C	N	O	0	0	0
			55	35	13	7			
2	H	8	Total	C	N	O	0	0	0
			76	47	21	8			

- Molecule 3 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula:  $C_{14}H_{20}N_6O_5S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
3	B	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
3	C	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
3	D	1	Total	C	N	O	S	0	0
			26	14	6	5	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	126	Total 126	O 126	0	0
4	B	147	Total 147	O 147	0	0
4	C	131	Total 131	O 131	0	0
4	D	156	Total 156	O 156	0	0
4	E	8	Total 8	O 8	0	0
4	F	3	Total 3	O 3	0	0
4	G	7	Total 7	O 7	0	0
4	H	11	Total 11	O 11	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone-lysine N-methyltransferase SETD8




- Molecule 1: Histone-lysine N-methyltransferase SETD8




- Molecule 1: Histone-lysine N-methyltransferase SETD8




- Molecule 1: Histone-lysine N-methyltransferase SETD8



- Molecule 2: Histone H4




- Molecule 2: Histone H4



● Molecule 2: Histone H4



● Molecule 2: Histone H4





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	43.99Å 45.55Å 94.41Å 89.37° 87.09° 90.50°	Depositor
Resolution (Å)	38.90 – 1.60 30.26 – 1.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (38.90-1.60) 95.5 (30.26-1.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.37 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.216 , 0.257 0.237 , 0.272	Depositor DCC
$R_{free}$ test set	4612 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.3	Xtriage
Anisotropy	0.013	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 41.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.017 for h,-k,-l 0.023 for -h,k,-l 0.013 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6115	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	8.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.63 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.6585e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.74	0/1325	0.81	2/1774 (0.1%)
1	B	0.77	0/1358	0.82	1/1819 (0.1%)
1	C	0.76	0/1328	0.86	1/1781 (0.1%)
1	D	0.83	0/1326	0.85	1/1779 (0.1%)
2	E	0.73	0/81	0.85	0/104
2	F	0.60	0/72	0.89	0/93
2	G	0.72	0/55	0.96	0/72
2	H	0.78	0/76	1.04	0/97
All	All	0.77	0/5621	0.84	5/7519 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	339	ARG	NE-CZ-NH2	-5.95	117.33	120.30
1	A	339	ARG	NE-CZ-NH2	-5.82	117.39	120.30
1	C	324	ARG	NE-CZ-NH2	-5.65	117.48	120.30
1	A	335	ASP	CB-CG-OD2	-5.26	113.57	118.30
1	D	292	ARG	NE-CZ-NH1	5.23	122.92	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1267	0	1243	15	0
1	B	1297	0	1275	8	0
1	C	1285	0	1264	17	0
1	D	1289	0	1267	10	0
2	E	81	0	84	3	0
2	F	72	0	80	1	0
2	G	55	0	56	1	0
2	H	76	0	91	1	0
3	A	26	0	19	0	0
3	B	26	0	19	0	0
3	C	26	0	19	0	0
3	D	26	0	19	0	0
4	A	126	0	0	11	0
4	B	147	0	0	1	0
4	C	131	0	0	8	0
4	D	156	0	0	2	0
4	E	8	0	0	2	0
4	F	3	0	0	0	0
4	G	7	0	0	0	0
4	H	11	0	0	0	0
All	All	6115	0	5436	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:238:ARG:HG2	1:B:238:ARG:HH11	1.33	0.93
4:A:396:HOH:O	1:C:276:GLN:HG3	1.71	0.89
1:C:195:LYS:HD3	4:C:362:HOH:O	1.76	0.84
1:C:223:ILE:HD11	1:C:231:ILE:HD11	1.59	0.83
1:D:207:ASP:OD1	1:D:310:HIS:NE2	2.17	0.78
1:C:235:GLN:HG2	4:C:411:HOH:O	1.83	0.77
1:B:238:ARG:HG2	1:B:238:ARG:NH1	2.00	0.73
1:C:235:GLN:CG	4:C:411:HOH:O	2.36	0.73
1:D:256:LYS:HG3	4:D:514:HOH:O	1.87	0.73
1:A:309[A]:LEU:HD11	4:A:364:HOH:O	1.89	0.69
1:C:207:ASP:O	1:C:211:GLU:HG3	1.98	0.64
1:A:309[A]:LEU:CD1	4:A:364:HOH:O	2.47	0.61
1:A:277:TYR:CE1	4:A:364:HOH:O	2.51	0.60
1:A:311:ASP:HB2	4:A:364:HOH:O	2.01	0.60

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:276:GLN:NE2	4:C:357:HOH:O	2.34	0.60
1:B:312:ILE:HD13	4:B:387:HOH:O	2.04	0.58
1:C:223:ILE:HD11	1:C:231:ILE:CD1	2.33	0.58
2:E:18:HIS:HE1	4:E:38:HOH:O	1.88	0.57
1:A:278:LEU:HB2	4:A:424:HOH:O	2.05	0.57
1:D:265:ASP:O	1:D:268[A]:THR:HG22	2.08	0.54
1:D:193:LYS:HD3	1:D:201:GLU:OE1	2.08	0.53
1:B:241:PHE:HA	1:B:321:ILE:HD13	1.92	0.52
1:D:262:TYR:HB3	1:D:268[A]:THR:HG21	1.91	0.52
1:D:319:ILE:HD12	1:D:321:ILE:HD11	1.91	0.52
1:A:312:ILE:O	1:A:315:VAL:HG12	2.11	0.51
1:A:277:TYR:HE1	4:A:364:HOH:O	1.94	0.49
1:A:315:VAL:HG11	4:A:379:HOH:O	2.11	0.49
1:C:324:ARG:CG	4:C:376:HOH:O	2.61	0.49
1:A:251[A]:GLU:HG2	1:A:282:TYR:CE1	2.48	0.49
1:C:277:TYR:HD1	1:C:309:LEU:HD21	1.78	0.48
1:C:304:ASN:HB2	4:C:376:HOH:O	2.12	0.48
1:A:245:PHE:CE1	2:F:20:LYS:HE3	2.49	0.48
4:A:366:HOH:O	2:G:19:ARG:CB	2.63	0.47
1:C:324:ARG:HG2	4:C:376:HOH:O	2.15	0.46
1:C:217:GLY:HA3	1:C:234:LYS:HG2	1.98	0.46
1:C:351:LYS:HE3	4:C:533:HOH:O	2.17	0.45
1:C:223:ILE:CD1	1:C:231:ILE:HD11	2.40	0.45
1:A:292:ARG:HD3	4:A:513:HOH:O	2.16	0.45
1:A:197:GLU:O	1:A:201:GLU:HG3	2.18	0.44
1:A:210:ILE:HD12	1:A:312:ILE:HD11	2.00	0.44
1:D:207:ASP:O	1:D:211:GLU:HG3	2.18	0.44
1:D:268[B]:THR:HG23	1:D:287:THR:CG2	2.48	0.43
4:A:396:HOH:O	1:C:276:GLN:CG	2.47	0.43
1:B:202:GLU:O	1:B:206:ILE:HG12	2.20	0.42
1:C:277:TYR:CD1	1:C:309:LEU:HD21	2.53	0.42
1:B:308:LYS:HG2	1:B:319:ILE:CG1	2.50	0.42
1:D:341:LYS:O	1:D:345:GLU:HG3	2.20	0.41
1:B:308:LYS:HG2	1:B:319:ILE:HG13	2.02	0.41
4:D:18:HOH:O	2:H:18:HIS:HE1	2.02	0.41
1:D:268[B]:THR:HG22	1:D:269:GLY:O	2.21	0.41
1:A:309[A]:LEU:HD13	1:A:310:HIS:N	2.36	0.41
1:B:245:PHE:CE1	2:E:20:LYS:HE3	2.56	0.40
2:E:18:HIS:CE1	4:E:38:HOH:O	2.68	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/166 (96%)	154 (97%)	5 (3%)	0	100	100
1	B	166/166 (100%)	163 (98%)	3 (2%)	0	100	100
1	C	162/166 (98%)	160 (99%)	2 (1%)	0	100	100
1	D	162/166 (98%)	158 (98%)	4 (2%)	0	100	100
2	E	7/10 (70%)	7 (100%)	0	0	100	100
2	F	6/10 (60%)	6 (100%)	0	0	100	100
2	G	5/10 (50%)	5 (100%)	0	0	100	100
2	H	6/10 (60%)	6 (100%)	0	0	100	100
All	All	673/704 (96%)	659 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	139/139 (100%)	138 (99%)	1 (1%)	84	73
1	B	143/139 (103%)	139 (97%)	4 (3%)	43	18
1	C	140/139 (101%)	138 (99%)	2 (1%)	67	47
1	D	139/139 (100%)	135 (97%)	4 (3%)	42	18
2	E	8/9 (89%)	8 (100%)	0	100	100
2	F	7/9 (78%)	7 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	G	5/9 (56%)	5 (100%)	0	100	100
2	H	8/9 (89%)	8 (100%)	0	100	100
All	All	589/592 (100%)	578 (98%)	11 (2%)	57	34

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	273	TYR
1	B	194	SER
1	B	203	ARG
1	B	273	TYR
1	B	308	LYS
1	C	273	TYR
1	C	308	LYS
1	D	199	GLN
1	D	251	GLU
1	D	273	TYR
1	D	300	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	276	GLN
1	A	291	ASN
1	B	276	GLN
1	C	235	GLN
1	C	276	GLN
1	D	276	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	SAH	C	801	-	21,28,28	1.25	1 (4%)	20,40,40	1.63	4 (20%)
3	SAH	B	801	-	21,28,28	1.15	1 (4%)	20,40,40	1.41	2 (10%)
3	SAH	D	801	-	21,28,28	1.42	3 (14%)	20,40,40	1.33	2 (10%)
3	SAH	A	801	-	21,28,28	1.39	2 (9%)	20,40,40	1.51	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SAH	C	801	-	-	1/7/31/31	0/3/3/3
3	SAH	B	801	-	-	1/7/31/31	0/3/3/3
3	SAH	D	801	-	-	1/7/31/31	0/3/3/3
3	SAH	A	801	-	-	2/7/31/31	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	801	SAH	O4'-C1'	4.39	1.47	1.41
3	C	801	SAH	O4'-C4'	4.26	1.54	1.45
3	D	801	SAH	O4'-C1'	2.95	1.45	1.41
3	D	801	SAH	O2'-C2'	2.84	1.49	1.43
3	B	801	SAH	O2'-C2'	2.70	1.49	1.43
3	D	801	SAH	C2'-C1'	-2.58	1.49	1.53

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	801	SAH	O2'-C2'	2.28	1.48	1.43

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	801	SAH	N3-C2-N1	-4.83	121.13	128.68
3	A	801	SAH	N3-C2-N1	-4.41	121.79	128.68
3	C	801	SAH	N3-C2-N1	-4.25	122.03	128.68
3	D	801	SAH	N3-C2-N1	-3.87	122.64	128.68
3	C	801	SAH	O4'-C4'-C5'	3.14	116.91	108.83
3	A	801	SAH	C1'-N9-C4	-2.68	121.93	126.64
3	D	801	SAH	C1'-N9-C4	-2.34	122.53	126.64
3	C	801	SAH	C2-N1-C6	2.21	122.53	118.75
3	C	801	SAH	C1'-N9-C4	-2.13	122.89	126.64
3	B	801	SAH	C1'-N9-C4	-2.13	122.90	126.64

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	801	SAH	C-CA-CB-CG
3	D	801	SAH	CB-CG-SD-C5'
3	A	801	SAH	CB-CG-SD-C5'
3	C	801	SAH	CB-CG-SD-C5'
3	B	801	SAH	CB-CG-SD-C5'


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




## Ligand SAH C 801



## Ligand SAH B 801





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	156/166 (93%)	0.46	12 (7%) 13 11	3, 7, 13, 16	0
1	B	161/166 (96%)	0.43	8 (4%) 28 26	2, 5, 14, 17	0
1	C	160/166 (96%)	0.30	7 (4%) 34 31	2, 5, 15, 22	0
1	D	161/166 (96%)	0.24	4 (2%) 57 55	2, 5, 17, 23	1 (0%)
2	E	9/10 (90%)	0.34	0 100 100	6, 9, 15, 20	0
2	F	8/10 (80%)	0.86	1 (12%) 3 3	12, 15, 26, 33	0
2	G	7/10 (70%)	0.18	0 100 100	11, 13, 19, 26	0
2	H	8/10 (80%)	0.83	0 100 100	9, 14, 20, 27	0
All	All	670/704 (95%)	0.37	32 (4%) 30 28	2, 6, 16, 33	1 (0%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	261	LEU	6.1
1	A	263	ALA	4.8
1	A	260	ALA	4.7
1	A	342	ALA	3.2
1	A	262	TYR	3.1
1	B	195	LYS	3.0
1	A	268	THR	3.0
1	D	192	ARG	2.9
1	A	196	ALA	2.9
2	F	16	LYS	2.8
1	A	199	GLN	2.7
1	C	297	ILE	2.7
1	D	243	VAL	2.6
1	C	214	LYS	2.6
1	D	195	LYS	2.6
1	B	297	ILE	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	261	LEU	2.6
1	B	264	GLN	2.5
1	B	345	GLU	2.5
1	A	194	SER	2.4
1	A	257	LYS	2.4
1	B	196	ALA	2.4
1	C	341	LYS	2.3
1	A	195	LYS	2.3
1	C	195	LYS	2.2
1	B	320[A]	LEU	2.2
1	C	264	GLN	2.1
1	C	346	ALA	2.1
1	C	345	GLU	2.0
1	B	334	TYR	2.0
1	A	269	GLY	2.0
1	D	266	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SAH	A	801	26/26	0.96	0.10	5,8,10,11	0
3	SAH	C	801	26/26	0.96	0.12	4,7,9,10	0
3	SAH	D	801	26/26	0.96	0.13	2,4,6,7	0
3	SAH	B	801	26/26	0.97	0.13	2,5,7,8	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around SAH C 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around SAH D 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers ⓘ

There are no such residues in this entry.