



Full wwPDB NMR Structure Validation Report ⓘ

May 31, 2020 – 10:12 pm BST

PDB ID : 6F9A
Title : Solution structure of the MRH domain of Yos9 complexed with alpha3,alpha6-Man5
Authors : Kniss, A.; Kazemi, S.; Lohr, F.; Guntert, P.; Dotsch, V.
Deposited on : 2017-12-14

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 89%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	162	79% 7% • 14%

2 Ensemble composition and analysis ⓘ

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:95-A:142, A:148-A:149, A:156-A:245 (140)	0.23	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 5 single-model clusters were found.

Cluster number	Models
1	1, 10, 12, 14, 17, 19, 20
2	2, 3, 5, 15, 18
3	6, 8, 11
Single-model clusters	4; 7; 9; 13; 16

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2534 atoms, of which 1248 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called ER quality-control lectin.

Mol	Chain	Residues	Atoms						Trace
1	A	162	Total	C	H	N	O	S	0
			2534	814	1248	211	254	7	

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	88	GLY	-	expression tag	UNP A0A250W9K2
A	89	ALA	-	expression tag	UNP A0A250W9K2

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: ER quality-control lectin




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1 (medoid)


- Molecule 1: ER quality-control lectin



4.2.2 Score per residue for model 2


- Molecule 1: ER quality-control lectin






4.2.8 Score per residue for model 8

- Molecule 1: ER quality-control lectin




4.2.9 Score per residue for model 9

- Molecule 1: ER quality-control lectin



4.2.10 Score per residue for model 10

- Molecule 1: ER quality-control lectin



4.2.11 Score per residue for model 11


- Molecule 1: ER quality-control lectin






4.2.12 Score per residue for model 12

- Molecule 1: ER quality-control lectin




4.2.13 Score per residue for model 13

- Molecule 1: ER quality-control lectin




4.2.14 Score per residue for model 14


- Molecule 1: ER quality-control lectin



4.2.15 Score per residue for model 15

- Molecule 1: ER quality-control lectin






4.2.16 Score per residue for model 16

- Molecule 1: ER quality-control lectin

Chain A: 75% 10% 14%



4.2.17 Score per residue for model 17

- Molecule 1: ER quality-control lectin


Chain A: 77% 8% 14%



4.2.18 Score per residue for model 18

- Molecule 1: ER quality-control lectin


Chain A: 75% 10% 14%



4.2.19 Score per residue for model 19


- Molecule 1: ER quality-control lectin

Chain A: 79% 7% 14%



4.2.20 Score per residue for model 20

- Molecule 1: ER quality-control lectin



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	3.9
OPAL	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 6 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	1891
Number of shifts mapped to atoms	1891
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	89%

No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

5.1 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1121	1097	1097	2±1
All	All	22420	21940	21940	41

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:221:VAL:HG13	1:A:229:TYR:CE1	0.75	2.16	20	20
1:A:181:TYR:CE2	1:A:243:LEU:HD22	0.60	2.30	7	4
1:A:181:TYR:CE2	1:A:243:LEU:HD13	0.49	2.42	1	2
1:A:181:TYR:CZ	1:A:243:LEU:HD22	0.48	2.44	1	2
1:A:102:LEU:O	1:A:106:VAL:HG23	0.46	2.10	1	4
1:A:106:VAL:HG22	1:A:172:PHE:CD2	0.44	2.48	16	1
1:A:109:ILE:HD11	1:A:183:ILE:HD12	0.44	1.89	12	1
1:A:220:TRP:CE3	1:A:222:ARG:HD2	0.42	2.50	17	1
1:A:226:ILE:HD12	1:A:226:ILE:H	0.42	1.73	3	1
1:A:127:TYR:CD1	1:A:221:VAL:HG11	0.42	2.50	9	3
1:A:216:SER:O	1:A:217:THR:HG23	0.41	2.14	16	1
1:A:220:TRP:CE3	1:A:222:ARG:CD	0.40	3.04	9	1

5.2 Torsion angles ⓘ

5.2.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	140/162 (86%)	128±2 (91±1%)	11±2 (8±1%)	1±1 (1±1%)	29	74
All	All	2800/3240 (86%)	2558 (91%)	224 (8%)	18 (1%)	29	74

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	212	ASN	7
1	A	131	PRO	4
1	A	189	SER	3
1	A	210	GLY	3
1	A	200	ARG	1

5.2.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation

was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	122/141 (87%)	109±2 (89±2%)	13±2 (11±2%)	10	54
All	All	2440/2820 (87%)	2180 (89%)	260 (11%)	10	54

All 48 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	222	ARG	20
1	A	110	PHE	20
1	A	107	LYS	20
1	A	119	PHE	20
1	A	135	PHE	20
1	A	141	ARG	16
1	A	217	THR	16
1	A	234	THR	14
1	A	112	LYS	14
1	A	229	TYR	14
1	A	227	CYS	11
1	A	208	CYS	10
1	A	128	GLU	5
1	A	242	GLU	4
1	A	189	SER	3
1	A	200	ARG	3
1	A	175	LEU	3
1	A	170	ARG	3
1	A	103	ASN	3
1	A	185	GLU	3
1	A	195	VAL	2
1	A	174	LEU	2
1	A	237	GLU	2
1	A	178	ASP	2
1	A	177	ASP	2
1	A	100	LYS	2
1	A	194	ASP	2
1	A	230	GLU	2
1	A	159	ARG	2
1	A	213	SER	2
1	A	108	THR	1
1	A	193	CYS	1
1	A	138	PHE	1
1	A	216	SER	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	199	GLU	1
1	A	142	VAL	1
1	A	95	THR	1
1	A	171	GLU	1
1	A	137	GLN	1
1	A	184	SER	1
1	A	173	GLU	1
1	A	116	ARG	1
1	A	176	TYR	1
1	A	160	LEU	1
1	A	211	SER	1
1	A	162	LYS	1
1	A	201	MET	1
1	A	104	GLN	1

5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

5.3 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.4 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.5 Ligand geometry [i](#)

There are no ligands in this entry.

5.6 Other polymers [i](#)

There are no such molecules in this entry.

5.7 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 89% for the well-defined parts and 85% for the entire structure.

6.1 Chemical shift list 1

File name: input_cs.cif

Chemical shift list name: *minimized.str*

6.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1891
Number of shifts mapped to atoms	1891
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

6.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	155	0.09 ± 0.13	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	142	-0.13 ± 0.26	None needed (< 0.5 ppm)
$^{13}\text{C}'$	146	0.28 ± 0.12	None needed (< 0.5 ppm)
^{15}N	146	0.09 ± 0.29	None needed (< 0.5 ppm)

6.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 89%, i.e. 1538 atoms were assigned a chemical shift out of a possible 1727. 0 out of 24 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	681/692 (98%)	273/276 (99%)	274/280 (98%)	134/136 (99%)
Sidechain	713/878 (81%)	431/510 (85%)	282/333 (85%)	0/35 (0%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	144/157 (92%)	76/82 (93%)	66/72 (92%)	2/3 (67%)
Overall	1538/1727 (89%)	780/868 (90%)	622/685 (91%)	136/174 (78%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 85%, i.e. 1681 atoms were assigned a chemical shift out of a possible 1976. 0 out of 25 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	747/802 (93%)	300/320 (94%)	301/324 (93%)	146/158 (92%)
Sidechain	790/1017 (78%)	482/591 (82%)	308/381 (81%)	0/45 (0%)
Aromatic	144/157 (92%)	76/82 (93%)	66/72 (92%)	2/3 (67%)
Overall	1681/1976 (85%)	858/993 (86%)	675/777 (87%)	148/206 (72%)

6.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	232	GLN	HB3	0.06	3.37 – 0.67	-7.3
1	A	232	GLN	HB2	0.25	3.30 – 0.80	-7.2

6.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

