



# Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 05:20 am BST

PDB ID : 4FAY  
Title : Crystal structure of a trimeric bacterial microcompartment shell protein PduB with glycerol metabolites  
Authors : Pang, A.H.; Prentice, M.B.; Pickersgill, R.W.  
Deposited on : 2012-05-22  
Resolution : 1.56 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	<div> <div>4%</div> <div> <div></div> <div>82%</div> <div>6%</div> <div>12%</div> </div> </div>
1	B	258	<div> <div>4%</div> <div> <div></div> <div>81%</div> <div>7%</div> <div>12%</div> </div> </div>
1	C	258	<div> <div>4%</div> <div> <div></div> <div>81%</div> <div>7%</div> <div>12%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	C	303	-	X	-	-
3	ACT	B	307	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5639 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Microcompartments protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	227	Total	C	N	O	S	8	7	0
			1712	1068	300	331	13			
1	B	227	Total	C	N	O	S	0	5	0
			1697	1058	299	327	13			
1	C	227	Total	C	N	O	S	0	8	0
			1723	1075	305	330	13			

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	EXPRESSION TAG	UNP A5VMB3
A	-18	GLY	-	EXPRESSION TAG	UNP A5VMB3
A	-17	SER	-	EXPRESSION TAG	UNP A5VMB3
A	-16	SER	-	EXPRESSION TAG	UNP A5VMB3
A	-15	HIS	-	EXPRESSION TAG	UNP A5VMB3
A	-14	HIS	-	EXPRESSION TAG	UNP A5VMB3
A	-13	HIS	-	EXPRESSION TAG	UNP A5VMB3
A	-12	PRO	-	EXPRESSION TAG	UNP A5VMB3
A	-11	PRO	-	EXPRESSION TAG	UNP A5VMB3
A	-10	LYS	-	EXPRESSION TAG	UNP A5VMB3
A	-9	SER	-	EXPRESSION TAG	UNP A5VMB3
A	-8	SER	-	EXPRESSION TAG	UNP A5VMB3
A	-7	GLY	-	EXPRESSION TAG	UNP A5VMB3
A	-6	LEU	-	EXPRESSION TAG	UNP A5VMB3
A	-5	VAL	-	EXPRESSION TAG	UNP A5VMB3
A	-4	PRO	-	EXPRESSION TAG	UNP A5VMB3
A	-3	ARG	-	EXPRESSION TAG	UNP A5VMB3
A	-2	GLY	-	EXPRESSION TAG	UNP A5VMB3
A	-1	SER	-	EXPRESSION TAG	UNP A5VMB3
A	0	HIS	-	EXPRESSION TAG	UNP A5VMB3
B	-19	MET	-	EXPRESSION TAG	UNP A5VMB3
B	-18	GLY	-	EXPRESSION TAG	UNP A5VMB3
B	-17	SER	-	EXPRESSION TAG	UNP A5VMB3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	SER	-	EXPRESSION TAG	UNP A5VMB3
B	-15	HIS	-	EXPRESSION TAG	UNP A5VMB3
B	-14	HIS	-	EXPRESSION TAG	UNP A5VMB3
B	-13	HIS	-	EXPRESSION TAG	UNP A5VMB3
B	-12	PRO	-	EXPRESSION TAG	UNP A5VMB3
B	-11	PRO	-	EXPRESSION TAG	UNP A5VMB3
B	-10	LYS	-	EXPRESSION TAG	UNP A5VMB3
B	-9	SER	-	EXPRESSION TAG	UNP A5VMB3
B	-8	SER	-	EXPRESSION TAG	UNP A5VMB3
B	-7	GLY	-	EXPRESSION TAG	UNP A5VMB3
B	-6	LEU	-	EXPRESSION TAG	UNP A5VMB3
B	-5	VAL	-	EXPRESSION TAG	UNP A5VMB3
B	-4	PRO	-	EXPRESSION TAG	UNP A5VMB3
B	-3	ARG	-	EXPRESSION TAG	UNP A5VMB3
B	-2	GLY	-	EXPRESSION TAG	UNP A5VMB3
B	-1	SER	-	EXPRESSION TAG	UNP A5VMB3
B	0	HIS	-	EXPRESSION TAG	UNP A5VMB3
C	-19	MET	-	EXPRESSION TAG	UNP A5VMB3
C	-18	GLY	-	EXPRESSION TAG	UNP A5VMB3
C	-17	SER	-	EXPRESSION TAG	UNP A5VMB3
C	-16	SER	-	EXPRESSION TAG	UNP A5VMB3
C	-15	HIS	-	EXPRESSION TAG	UNP A5VMB3
C	-14	HIS	-	EXPRESSION TAG	UNP A5VMB3
C	-13	HIS	-	EXPRESSION TAG	UNP A5VMB3
C	-12	PRO	-	EXPRESSION TAG	UNP A5VMB3
C	-11	PRO	-	EXPRESSION TAG	UNP A5VMB3
C	-10	LYS	-	EXPRESSION TAG	UNP A5VMB3
C	-9	SER	-	EXPRESSION TAG	UNP A5VMB3
C	-8	SER	-	EXPRESSION TAG	UNP A5VMB3
C	-7	GLY	-	EXPRESSION TAG	UNP A5VMB3
C	-6	LEU	-	EXPRESSION TAG	UNP A5VMB3
C	-5	VAL	-	EXPRESSION TAG	UNP A5VMB3
C	-4	PRO	-	EXPRESSION TAG	UNP A5VMB3
C	-3	ARG	-	EXPRESSION TAG	UNP A5VMB3
C	-2	GLY	-	EXPRESSION TAG	UNP A5VMB3
C	-1	SER	-	EXPRESSION TAG	UNP A5VMB3
C	0	HIS	-	EXPRESSION TAG	UNP A5VMB3

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	136	Total	O	0	0
			136	136		
4	B	131	Total	O	0	0
			131	131		
4	C	134	Total	O	0	0
			134	134		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Microcompartments protein



- Molecule 1: Microcompartments protein



- Molecule 1: Microcompartments protein





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.86Å 120.86Å 145.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.78 – 1.56 46.51 – 1.56	Depositor EDS
% Data completeness (in resolution range)	99.6 (72.78-1.56) 99.3 (46.51-1.56)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	127.27 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.173 , 0.202 0.172 , 0.199	Depositor DCC
$R_{free}$ test set	4338 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.4	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 51.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.010 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.034 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5639	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.97	0/1739	0.84	1/2347 (0.0%)
1	B	0.95	1/1724 (0.1%)	0.82	2/2328 (0.1%)
1	C	0.93	0/1752	0.80	1/2364 (0.0%)
All	All	0.95	1/5215 (0.0%)	0.82	4/7039 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	107	ALA	CA-CB	5.53	1.64	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	64	MET	CG-SD-CE	-5.90	90.76	100.20
1	A	209	GLN	CA-CB-CG	5.16	124.76	113.40
1	B	100	ASP	CB-CG-OD1	5.08	122.87	118.30
1	B	215	ARG	NE-CZ-NH1	5.05	122.83	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1712	0	1694	14	0
1	B	1697	0	1679	17	0
1	C	1723	0	1711	15	0
2	A	42	0	56	3	0
2	B	36	0	48	5	0
2	C	24	0	30	4	0
3	B	4	0	3	2	0
4	A	136	0	0	0	0
4	B	131	0	0	4	0
4	C	134	0	0	1	0
All	All	5639	0	5221	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:LEU:HD11	1:A:197[A]:CYS:SG	1.90	1.11
1:C:156[A]:LEU:HD11	1:C:197[A]:CYS:SG	1.98	1.03
1:A:156:LEU:CD1	1:A:197[A]:CYS:SG	2.55	0.93
1:B:156:LEU:HD11	1:B:197[A]:CYS:SG	2.11	0.91
1:A:124:ASN:OD1	2:A:304:GOL:H31	1.80	0.81
1:C:156[A]:LEU:CD1	1:C:197[A]:CYS:SG	2.69	0.80
1:C:158[A]:CYS:HB2	1:C:197[A]:CYS:SG	2.22	0.79
1:B:156:LEU:CD1	1:B:197[A]:CYS:SG	2.75	0.75
1:A:39:LEU:O	2:A:305:GOL:H11	1.88	0.74
1:B:124:ASN:ND2	2:B:304:GOL:H31	2.05	0.72
1:B:124:ASN:HD21	2:B:304:GOL:H31	1.57	0.68
1:C:56:GLY:HA2	2:C:303:GOL:H2	1.80	0.64
1:B:136:ALA:O	3:B:307:ACT:H1	1.98	0.63
1:B:156:LEU:HD11	1:B:197[A]:CYS:HG	1.63	0.62
1:B:171:LYS:HE2	4:B:486:HOH:O	2.01	0.60
1:A:156:LEU:HD12	1:A:197[A]:CYS:SG	2.41	0.59
1:B:231:ASN:ND2	4:B:493:HOH:O	2.37	0.57
1:B:174:LYS:O	2:B:306:GOL:H12	2.05	0.56
1:A:156:LEU:HD11	1:A:197[A]:CYS:HG	1.69	0.56
1:C:131:GLU:OE1	2:C:303:GOL:C3	2.57	0.53
1:A:65:ASP:OD1	1:A:139:ALA:HB3	2.09	0.52
1:B:185:SER:HB2	1:B:186:PRO:CD	2.41	0.50
1:B:158[A]:CYS:HB2	1:B:197[A]:CYS:SG	2.51	0.49
1:A:124:ASN:O	1:A:230:VAL:HG12	2.13	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:185:SER:HB2	1:A:186:PRO:CD	2.43	0.48
1:A:160:CYS:HA	1:A:161:PRO:C	2.34	0.47
1:B:185:SER:HB2	1:B:186:PRO:HD2	1.96	0.47
1:C:185:SER:HB2	1:C:186:PRO:CD	2.45	0.46
1:A:191:SER:HB3	1:C:192:PHE:CG	2.51	0.45
1:B:171:LYS:HE3	4:B:477:HOH:O	2.16	0.45
3:B:307:ACT:H2	4:B:483:HOH:O	2.17	0.44
1:A:39:LEU:O	2:A:305:GOL:C1	2.63	0.44
1:C:158[A]:CYS:SG	1:C:195:GLU:HB3	2.57	0.44
1:C:160:CYS:HA	1:C:161:PRO:C	2.37	0.44
1:C:131:GLU:OE1	2:C:303:GOL:H31	2.19	0.42
1:C:109[A]:ARG:NH2	4:C:437:HOH:O	2.43	0.42
1:B:174:LYS:O	2:B:306:GOL:C1	2.66	0.42
1:C:156[B]:LEU:C	1:C:156[B]:LEU:HD23	2.40	0.42
1:A:161:PRO:HB2	1:B:52:SER:OG	2.20	0.42
1:B:40:HIS:CE1	2:B:305:GOL:H12	2.54	0.42
1:A:58:GLY:HA3	1:A:131:GLU:OE2	2.20	0.41
1:B:180:PRO:HB2	1:C:38:LYS:HE2	2.02	0.41
1:C:100:ASP:HA	1:C:101:PRO:HD3	1.93	0.41
1:C:80:TRP:CG	2:C:301:GOL:H11	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	231/258 (90%)	223 (96%)	7 (3%)	1 (0%)	34	14
1	B	229/258 (89%)	221 (96%)	7 (3%)	1 (0%)	34	14
1	C	232/258 (90%)	227 (98%)	4 (2%)	1 (0%)	34	14
All	All	692/774 (89%)	671 (97%)	18 (3%)	3 (0%)	34	14

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	83	ASP
1	C	83	ASP
1	A	83	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	179/200 (90%)	179 (100%)	0	100	100
1	B	177/200 (88%)	176 (99%)	1 (1%)	86	73
1	C	180/200 (90%)	180 (100%)	0	100	100
All	All	536/600 (89%)	535 (100%)	1 (0%)	93	86

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	65	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	124	ASN
1	B	209	GLN
1	C	40	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	C	301	-	5,5,5	0.41	0	5,5,5	0.70	0
2	GOL	A	305	-	5,5,5	0.32	0	5,5,5	0.39	0
2	GOL	B	301	-	5,5,5	0.43	0	5,5,5	0.27	0
2	GOL	B	302	-	5,5,5	0.30	0	5,5,5	0.71	0
2	GOL	B	305	-	5,5,5	0.88	0	5,5,5	1.23	0
2	GOL	B	306	-	5,5,5	0.35	0	5,5,5	0.52	0
2	GOL	A	304	-	5,5,5	0.37	0	5,5,5	0.28	0
2	GOL	C	303	-	5,5,5	0.67	0	5,5,5	1.59	2 (40%)
2	GOL	C	304	-	5,5,5	1.36	1 (20%)	5,5,5	0.76	0
2	GOL	A	303	-	5,5,5	0.33	0	5,5,5	0.51	0
2	GOL	B	303	-	5,5,5	0.37	0	5,5,5	0.27	0
2	GOL	A	307	-	5,5,5	0.33	0	5,5,5	0.38	0
2	GOL	C	302	-	5,5,5	0.82	0	5,5,5	0.61	0
3	ACT	B	307	-	1,3,3	2.30	1 (100%)	0,3,3	0.00	-
2	GOL	B	304	-	5,5,5	0.36	0	5,5,5	0.27	0
2	GOL	A	302	-	5,5,5	0.37	0	5,5,5	0.66	0
2	GOL	A	301	-	5,5,5	0.37	0	5,5,5	0.28	0
2	GOL	A	306	-	5,5,5	0.30	0	5,5,5	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	301	-	-	4/4/4/4	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	305	-	-	2/4/4/4	-
2	GOL	B	301	-	-	0/4/4/4	-
2	GOL	B	302	-	-	0/4/4/4	-
2	GOL	B	305	-	-	0/4/4/4	-
2	GOL	B	306	-	-	3/4/4/4	-
2	GOL	A	304	-	-	1/4/4/4	-
2	GOL	C	303	-	-	4/4/4/4	-
2	GOL	C	304	-	-	0/4/4/4	-
2	GOL	A	303	-	-	0/4/4/4	-
2	GOL	B	303	-	-	0/4/4/4	-
2	GOL	A	307	-	-	4/4/4/4	-
2	GOL	C	302	-	-	0/4/4/4	-
2	GOL	A	302	-	-	0/4/4/4	-
2	GOL	B	304	-	-	0/4/4/4	-
2	GOL	A	301	-	-	0/4/4/4	-
2	GOL	A	306	-	-	3/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	307	ACT	CH3-C	2.30	1.51	1.48
2	C	304	GOL	O2-C2	-2.20	1.36	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	303	GOL	O2-C2-C1	2.27	119.11	109.12
2	C	303	GOL	C3-C2-C1	-2.06	103.68	111.70

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	301	GOL	O1-C1-C2-C3
2	C	301	GOL	C1-C2-C3-O3
2	A	307	GOL	C1-C2-C3-O3
2	A	305	GOL	C1-C2-C3-O3
2	B	306	GOL	C1-C2-C3-O3
2	C	303	GOL	O1-C1-C2-C3
2	C	303	GOL	C1-C2-C3-O3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	A	307	GOL	O1-C1-C2-C3
2	A	306	GOL	C1-C2-C3-O3
2	C	303	GOL	O2-C2-C3-O3
2	A	307	GOL	O2-C2-C3-O3
2	C	301	GOL	O1-C1-C2-O2
2	C	301	GOL	O2-C2-C3-O3
2	A	306	GOL	O2-C2-C3-O3
2	B	306	GOL	O1-C1-C2-C3
2	C	303	GOL	O1-C1-C2-O2
2	A	306	GOL	O1-C1-C2-C3
2	A	305	GOL	O2-C2-C3-O3
2	A	307	GOL	O1-C1-C2-O2
2	B	306	GOL	O2-C2-C3-O3
2	A	304	GOL	C1-C2-C3-O3

There are no ring outliers.

8 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	301	GOL	1	0
2	A	305	GOL	2	0
2	B	305	GOL	1	0
2	B	306	GOL	2	0
2	A	304	GOL	1	0
2	C	303	GOL	3	0
3	B	307	ACT	2	0
2	B	304	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	227/258 (87%)	0.33	11 (4%) 30 35	10, 15, 27, 35	0
1	B	227/258 (87%)	0.30	11 (4%) 30 35	10, 15, 27, 36	0
1	C	227/258 (87%)	0.31	11 (4%) 30 35	11, 15, 27, 36	0
All	All	681/774 (87%)	0.31	33 (4%) 30 35	10, 15, 27, 36	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	236	TYR	5.0
1	C	236	TYR	5.0
1	B	237[A]	ILE	4.4
1	A	236	TYR	4.0
1	C	30	ARG	4.0
1	C	158[A]	CYS	3.9
1	A	233	PHE	3.5
1	B	11	VAL	3.4
1	C	11	VAL	2.9
1	B	230	VAL	2.9
1	A	84	THR	2.8
1	B	231	ASN	2.7
1	B	116	HIS	2.6
1	C	237[A]	ILE	2.6
1	C	233	PHE	2.5
1	B	113	ASP	2.4
1	C	19	GLU	2.4
1	A	117	ARG	2.4
1	C	113	ASP	2.4
1	B	39	LEU	2.4
1	A	139	ALA	2.3
1	A	237[A]	ILE	2.3
1	C	235	SER	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	11	VAL	2.3
1	C	232	ASP	2.3
1	A	197[A]	CYS	2.2
1	A	230	VAL	2.2
1	A	30	ARG	2.2
1	B	197[A]	CYS	2.2
1	A	228	GLU	2.1
1	B	232	ASP	2.1
1	C	228	GLU	2.1
1	B	233	PHE	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ACT	B	307	4/4	0.02	0.31	40,40,41,41	0
2	GOL	A	306	6/6	0.44	0.24	51,52,53,55	0
2	GOL	A	301	6/6	0.59	0.28	28,38,39,41	0
2	GOL	B	306	6/6	0.66	0.21	34,40,40,42	0
2	GOL	B	303	6/6	0.73	0.20	16,22,23,26	0
2	GOL	A	304	6/6	0.74	0.24	43,44,44,44	0
2	GOL	C	304	6/6	0.75	0.23	28,37,39,40	0
2	GOL	B	301	6/6	0.78	0.16	26,33,36,37	0
2	GOL	B	304	6/6	0.79	0.20	34,39,42,42	0
2	GOL	A	307	6/6	0.79	0.13	32,40,41,44	0
2	GOL	B	305	6/6	0.79	0.21	56,57,57,58	0
2	GOL	B	302	6/6	0.80	0.13	19,23,24,25	0
2	GOL	A	305	6/6	0.81	0.12	41,42,43,45	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GOL	C	303	6/6	0.83	0.17	16,29,34,37	0
2	GOL	A	302	6/6	0.84	0.12	20,22,26,27	0
2	GOL	C	302	6/6	0.84	0.12	19,22,24,24	0
2	GOL	C	301	6/6	0.86	0.22	24,38,42,43	0
2	GOL	A	303	6/6	0.87	0.12	18,24,27,31	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.