



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 8, 2020 – 07:30 AM BST

PDB ID : 2FD6
Title : Structure of Human Urokinase Plasminogen Activator in Complex with Urokinase Receptor and an anti-upar antibody at 1.9 Å
Authors : Huang, M.; Huai, Q.; Li, Y.
Deposited on : 2005-12-13
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	122	<div> <div>25%</div> <div>54%</div> <div>41%</div> <div>• •</div> </div>
2	L	214	<div> <div>3%</div> <div>66%</div> <div>31%</div> <div>•</div> </div>
3	H	213	<div> <div>4%</div> <div>73%</div> <div>24%</div> <div>•</div> </div>
4	U	276	<div> <div>7%</div> <div>52%</div> <div>35%</div> <div>• 10%</div> </div>
5	B	2	<div> <div>50%</div> <div>50%</div> </div>
6	C	2	<div> <div>100%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
12	PG4	U	402	-	-	X	-
5	NAG	B	1	X	-	-	-
5	FUC	B	2	X	-	-	-
6	NAG	C	1	-	-	X	-
6	NAG	C	2	X	-	-	X
8	ETX	L	401	-	-	X	-

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 6600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Urokinase-type plasminogen activator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	122	Total	C	N	O	S	0	0	0
			967	596	183	174	14			

- Molecule 2 is a protein called L chain of Fab of ATN-615 anti-uPAR antibody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	214	Total	C	N	O	S	0	0	0
			1648	1032	272	337	7			

- Molecule 3 is a protein called H chain of Fab of ATN-615 anti-uPAR antibody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	212	Total	C	N	O	S	0	0	0
			1618	1035	262	315	6			

- Molecule 4 is a protein called Urokinase plasminogen activator surface receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	U	249	Total	C	N	O	S	0	0	0
			1906	1141	349	382	34			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
U	1A	SER	-	cloning artifact	UNP Q03405

- Molecule 5 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	B	2	Total	C	N	O	0	0	0
			26	14	1	11			

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	C	2	Total	C	N	O	0	0	0
			30	16	2	12			

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 8 is 2-ETHOXYETHANOL (three-letter code: ETX) (formula: C₄H₁₀O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	L	1	Total	C	O	0	0
			6	4	2		
8	L	1	Total	C	O	0	0
			6	4	2		
8	H	1	Total	C	O	0	0
			6	4	2		

- Molecule 9 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	L	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	H	1	Total	C	O	0	0
			4	2	2		

- Molecule 10 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	H	1	Total	C	O	0	0
			10	6	4		

- Molecule 11 is 2-acetamido-2-deoxy- α -D-glucopyranose (three-letter code: NDG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	U	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 12 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	U	1	Total	C	O	0	0
			13	8	5		


- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	22	Total	O	0	0
			22	22		
13	L	121	Total	O	0	0
			121	121		
13	H	121	Total	O	0	0
			121	121		
13	U	72	Total	O	0	0
			72	72		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Urokinase-type plasminogen activator



- Molecule 2: L chain of Fab of ATN-615 anti-uPAR antibody




- Molecule 3: H chain of Fab of ATN-615 anti-uPAR antibody



- Molecule 4: Urokinase plasminogen activator surface receptor





- Molecule 5: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:



- Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.79Å 86.81Å 124.69Å 90.00° 94.54° 90.00°	Depositor
Resolution (Å)	41.36 – 1.90 41.36 – 1.85	Depositor EDS
% Data completeness (in resolution range)	95.8 (41.36-1.90) 92.4 (41.36-1.85)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 1.86Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.239 , 0.276 0.240 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	29.1	Xtriage
Anisotropy	0.350	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 58.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6600	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, NAG, ETX, EDO, NDG, PG4, SO4, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/994	0.59	0/1342
2	L	0.40	0/1688	0.70	1/2292 (0.0%)
3	H	0.50	2/1668 (0.1%)	0.77	3/2282 (0.1%)
4	U	0.35	0/1933	0.66	0/2599
All	All	0.41	2/6283 (0.0%)	0.69	4/8515 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	134	GLY	C-N	-8.55	1.14	1.34
3	H	134	GLY	N-CA	-5.83	1.37	1.46

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	134	GLY	CA-C-N	-7.82	100.00	117.20
3	H	133	LEU	C-N-CA	-7.25	107.08	122.30
3	H	134	GLY	O-C-N	6.50	133.11	122.70
2	L	93	TYR	N-CA-C	5.40	125.59	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	967	0	902	61	0
2	L	1648	0	1577	70	0
3	H	1618	0	1557	62	0
4	U	1906	0	1773	124	0
5	B	26	0	24	6	0
6	C	30	0	27	9	0
7	A	5	0	0	0	0
8	H	6	0	10	3	0
8	L	12	0	20	8	0
9	H	4	0	6	0	0
9	L	4	0	6	0	0
10	H	10	0	14	1	0
11	U	15	0	12	2	0
12	U	13	0	18	11	0
13	A	22	0	0	2	0
13	H	121	0	0	3	0
13	L	121	0	0	5	0
13	U	72	0	0	14	0
All	All	6600	0	5946	319	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

The worst 5 of 319 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:U:230:GLU:HB3	4:U:231:PRO:HD2	1.36	1.06
6:C:1:NAG:H61	6:C:2:NAG:H82	1.37	1.05
13:U:530:HOH:O	5:B:1:NAG:H83	1.57	1.02
4:U:221:GLN:HE21	12:U:402:PG4:H32	1.25	1.01
6:C:1:NAG:C6	6:C:2:NAG:H82	1.91	1.00

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	120/122 (98%)	98 (82%)	18 (15%)	4 (3%)	4	0
2	L	212/214 (99%)	204 (96%)	4 (2%)	4 (2%)	8	1
3	H	208/213 (98%)	199 (96%)	7 (3%)	2 (1%)	15	6
4	U	239/276 (87%)	215 (90%)	20 (8%)	4 (2%)	9	2
All	All	779/825 (94%)	716 (92%)	49 (6%)	14 (2%)	8	2

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	52	GLU
1	A	62	ALA
1	A	109	ARG
2	L	93	TYR
4	U	230	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	106/106 (100%)	102 (96%)	4 (4%)	33	24
2	L	186/186 (100%)	181 (97%)	5 (3%)	44	38
3	H	182/182 (100%)	176 (97%)	6 (3%)	38	29
4	U	223/244 (91%)	212 (95%)	11 (5%)	25	15
All	All	697/718 (97%)	671 (96%)	26 (4%)	34	25

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	H	62	LYS
3	H	201	VAL
4	U	212	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	H	144	PRO
3	H	172	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
3	H	31	ASN
3	H	166	GLN
4	U	229	HIS
3	H	128	ASN
4	U	78	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	B	1	5,4	15,15,15	0.46	0	21,21,21	0.79	1 (4%)
5	FUC	B	2	5	11,11,11	0.39	0	15,16,16	0.31	0
6	NAG	C	1	4,6	15,15,15	0.42	0	21,21,21	0.61	0
6	NAG	C	2	6	15,15,15	0.46	0	21,21,21	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	1	5,4	1/1/6/7	5/6/26/26	0/1/1/1
5	FUC	B	2	5	1/1/5/5	-	0/1/1/1
6	NAG	C	1	4,6	-	5/6/26/26	0/1/1/1
6	NAG	C	2	6	1/1/6/7	4/6/26/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1	NAG	O5-C1-C2	2.06	111.58	109.52

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	C	2	NAG	C1
5	B	2	FUC	C1
5	B	1	NAG	C1

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	2	NAG	C8-C7-N2-C2
6	C	2	NAG	O7-C7-N2-C2
6	C	1	NAG	C1-C2-N2-C7
6	C	1	NAG	C8-C7-N2-C2
6	C	1	NAG	O7-C7-N2-C2

There are no ring outliers.

3 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	2	NAG	5	0
6	C	1	NAG	9	0
5	B	1	NAG	6	0

5.6 Ligand geometry

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	EDO	L	405	-	3,3,3	0.56	0	2,2,2	0.61	0
8	ETX	H	403	-	5,5,5	1.12	0	4,4,4	2.58	1 (25%)
9	EDO	H	404	-	3,3,3	0.48	0	2,2,2	0.54	0
8	ETX	L	407	-	5,5,5	1.13	0	4,4,4	2.52	1 (25%)
7	SO4	A	501	-	4,4,4	0.20	0	6,6,6	0.11	0
12	PG4	U	402	-	12,12,12	1.11	1 (8%)	11,11,11	0.82	0
11	NDG	U	400	-	15,15,15	0.43	0	21,21,21	0.63	0
10	PGE	H	406	-	9,9,9	1.16	1 (11%)	8,8,8	1.91	2 (25%)
8	ETX	L	401	-	5,5,5	1.17	0	4,4,4	2.58	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	EDO	L	405	-	-	1/1/1/1	-
8	ETX	H	403	-	-	1/3/3/3	-
11	NDG	U	400	-	-	5/6/26/26	0/1/1/1
8	ETX	L	407	-	-	1/3/3/3	-
12	PG4	U	402	-	-	7/10/10/10	-
9	EDO	H	404	-	-	1/1/1/1	-
10	PGE	H	406	-	-	4/7/7/7	-
8	ETX	L	401	-	-	2/3/3/3	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	U	402	PG4	O2-C3	2.20	1.51	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	H	406	PGE	C4-C3	2.04	1.59	1.49

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	H	403	ETX	C3-O2-C2	4.85	130.26	112.90
8	L	401	ETX	C3-O2-C2	4.78	130.01	112.90
8	L	407	ETX	C3-O2-C2	4.76	129.94	112.90
10	H	406	PGE	C5-O3-C4	3.99	130.56	113.29
10	H	406	PGE	O2-C2-C1	2.70	121.92	110.07

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
11	U	400	NDG	C1-C2-N2-C7
11	U	400	NDG	C8-C7-N2-C2
11	U	400	NDG	O7-C7-N2-C2
10	H	406	PGE	O1-C1-C2-O2
11	U	400	NDG	O5-C5-C6-O6

There are no ring outliers.

6 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	H	403	ETX	3	0
8	L	407	ETX	1	0
12	U	402	PG4	11	0
11	U	400	NDG	2	0
10	H	406	PGE	1	0
8	L	401	ETX	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	H	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	H	134:GLY	C	135:CYS	N	1.14

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	122/122 (100%)	1.32	31 (25%) 0 0	33, 56, 82, 86	0
2	L	214/214 (100%)	0.24	7 (3%) 46 49	21, 33, 54, 91	0
3	H	212/213 (99%)	0.22	8 (3%) 40 43	17, 33, 52, 64	0
4	U	249/276 (90%)	0.61	19 (7%) 13 15	22, 43, 70, 99	0
All	All	797/825 (96%)	0.52	65 (8%) 11 13	17, 39, 72, 99	0

The worst 5 of 65 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	215	ALA	6.4
1	A	53	GLY	6.1
2	L	213	ALA	6.0
1	A	131	CYS	5.9
4	U	230	GLU	5.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NAG	C	2	15/15	0.58	0.52	98,102,104,105	0
6	NAG	C	1	15/15	0.64	0.26	71,76,79,80	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	NAG	B	1	15/15	0.70	0.25	78,81,84,86	0
5	FUC	B	2	11/11	0.90	0.29	98,99,101,102	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
8	ETX	L	401	6/6	0.51	0.35	41,48,52,61	0
8	ETX	H	403	6/6	0.52	0.38	43,48,49,52	0
8	ETX	L	407	6/6	0.54	0.25	52,56,59,59	0
10	PGE	H	406	10/10	0.68	0.23	38,61,63,64	0
11	NDG	U	400	15/15	0.70	0.20	71,81,83,85	0
12	PG4	U	402	13/13	0.73	0.29	44,48,54,58	0
9	EDO	L	405	4/4	0.85	0.12	47,49,49,50	0
9	EDO	H	404	4/4	0.89	0.14	42,46,50,53	0
7	SO4	A	501	5/5	0.98	0.13	47,47,47,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.