



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 19, 2022 – 08:14 AM JST

PDB ID : 7FF8  
Title : Pseudomonas aeruginosa Virulence Factor Regulator with cAMP ligand and Cl(triethylphosphine)gold(I)  
Authors : Chew, B.L.A.; Luo, D.  
Deposited on : 2021-07-22  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	?? (??), CSD ??CSD?? (????)
Xtriage (Phenix)	:	1.13
EDS	:	2.29
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.29


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	210	
1	B	210	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 2924 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called cAMP-activated global transcriptional regulator Vfr.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	197	Total	C	N	O	S	0	0	0
			1462	924	254	274	10			
1	B	188	Total	C	N	O	S	0	0	0
			1375	869	243	255	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	85	GLU	GLN	conflict	UNP P55222
B	85	GLU	GLN	conflict	UNP P55222

- Molecule 2 is ADENOSINE-3',5'-CYCLIC-MONOPHOSPHATE (three-letter code: CMP) (formula: C<sub>10</sub>H<sub>12</sub>N<sub>5</sub>O<sub>6</sub>P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			22	10	5	6	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			22	10	5	6	1		

- Molecule 3 is triethylphosphanuidylgold(1+) (three-letter code: AUF) (formula:  $C_6H_{15}AuP$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	Au	C	P	0	0
			8	1	6	1		
3	B	1	Total	Au	C	P	0	0
			8	1	6	1		


- Molecule 4 is GOLD ION (three-letter code: AU) (formula: Au) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Au	0	0
			2	2		
4	B	3	Total	Au	0	0
			3	3		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		
5	B	1	Total	Cl	0	0
			1	1		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: cAMP-activated global transcriptional regulator Vfr

Chain A: 



- Molecule 1: cAMP-activated global transcriptional regulator Vfr

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	140.26Å 49.93Å 62.38Å 90.00° 100.14° 90.00°	Depositor
Resolution (Å)	34.52 – 2.80 38.43 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (34.52-2.80) 92.8 (38.43-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.08 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, $R_{free}$	0.269 , 0.327 0.269 , 0.326	Depositor DCC
$R_{free}$ test set	560 reflections (5.24%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.3	Xtriage
Anisotropy	0.418	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 58.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.85	EDS
Total number of atoms	2924	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CMP, AU, CL, AUF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.57	5/1479 (0.3%)	0.93	5/1997 (0.3%)
1	B	0.43	1/1391 (0.1%)	0.70	3/1879 (0.2%)
All	All	0.50	6/2870 (0.2%)	0.83	8/3876 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	190	ARG	CB-CG	-9.43	1.27	1.52
1	A	101	GLU	CD-OE1	8.20	1.34	1.25
1	A	162	MET	CB-CG	5.67	1.69	1.51
1	A	202	HIS	CG-CD2	5.22	1.44	1.35
1	B	158	GLN	C-N	-5.12	1.24	1.34
1	A	190	ARG	CZ-NH2	-5.06	1.26	1.33

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	190	ARG	NE-CZ-NH2	-18.22	111.19	120.30
1	A	190	ARG	NE-CZ-NH1	14.91	127.75	120.30
1	A	162	MET	CA-CB-CG	14.17	137.39	113.30
1	A	190	ARG	CG-CD-NE	-11.34	87.98	111.80
1	B	200	LEU	CA-CB-CG	9.99	138.29	115.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	101	GLU	CA-CB-CG	6.41	127.50	113.40
1	B	77	LEU	CA-CB-CG	6.03	129.17	115.30
1	A	162	MET	CB-CG-SD	5.22	128.07	112.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	101	GLU	Sidechain,Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1462	0	1414	29	0
1	B	1375	0	1313	27	0
2	A	22	0	11	1	0
2	B	22	0	11	1	0
3	A	8	0	0	0	0
3	B	8	0	0	1	0
4	A	2	0	0	0	0
4	B	3	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	10	0	0	0	0
6	B	10	0	0	0	0
All	All	2924	0	2749	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:301:CMF:H2	2:A:301:CMF:C2	0.97	1.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:302:CMP:H2	2:B:302:CMP:C2	0.97	1.48
1:A:162:MET:SD	1:A:162:MET:CE	2.03	1.46
1:A:190:ARG:HG3	1:A:190:ARG:NH1	1.91	0.82
1:B:21:HIS:HA	1:B:101:GLU:HG3	1.62	0.79
1:B:156:CYS:SG	3:B:303:AUF:C1	2.70	0.79
1:B:150:ARG:HG2	1:B:154:ASP:OD1	1.82	0.79
1:A:190:ARG:O	1:A:194:SER:N	2.14	0.77
1:A:71:PHE:O	1:A:128:ARG:NH1	2.18	0.77
1:B:74:GLU:HG3	1:B:121:LEU:HD21	1.73	0.71
1:A:148:VAL:HB	1:A:191:VAL:HG11	1.74	0.69
1:B:71:PHE:O	1:B:128:ARG:NH1	2.24	0.66
1:B:150:ARG:O	1:B:154:ASP:N	2.26	0.66
1:B:60:GLU:OE1	1:B:92:ARG:NH1	2.27	0.65
1:A:55:ASP:N	1:A:55:ASP:OD1	2.24	0.63
1:A:54:GLU:O	1:B:134:ARG:NH2	2.34	0.60
1:A:191:VAL:O	1:A:195:LEU:HG	2.05	0.56
1:A:150:ARG:HG3	1:A:154:ASP:OD1	2.06	0.56
1:A:114:ASP:OD2	1:A:116:GLU:N	2.28	0.54
1:A:139:LEU:HD23	1:A:147:ARG:HD2	1.89	0.54
1:B:139:LEU:HD22	1:B:147:ARG:HD3	1.89	0.54
1:B:95:VAL:HG12	1:B:96:GLU:H	1.72	0.53
1:B:54:GLU:HB2	1:B:90:TRP:CZ3	2.43	0.53
1:A:143:ASP:O	1:A:147:ARG:HG3	2.10	0.52
1:B:41:LEU:O	1:B:102:ILE:N	2.39	0.52
1:A:138:ASP:O	1:A:142:LEU:HB2	2.10	0.52
1:B:194:SER:O	1:B:198:GLN:HG3	2.10	0.52
1:A:136:VAL:HG22	1:B:136:VAL:HG22	1.92	0.51
1:A:119:TYR:CE1	1:B:118:LEU:HD21	2.45	0.51
1:B:36:ASP:O	1:B:87:ARG:HG2	2.13	0.49
1:A:192:LEU:HB3	1:A:203:VAL:HG21	1.94	0.48
1:B:140:ALA:O	1:B:141:PHE:HB2	2.14	0.48
1:B:132:THR:O	1:B:136:VAL:HG23	2.14	0.47
1:A:190:ARG:HG3	1:A:190:ARG:HH11	1.78	0.47
1:A:54:GLU:HA	1:A:59:ARG:O	2.15	0.47
1:A:101:GLU:HG3	1:A:102:ILE:N	2.30	0.46
1:B:69:GLY:O	1:B:124:GLN:NE2	2.48	0.46
1:A:190:ARG:NH1	1:A:190:ARG:CG	2.69	0.45
1:A:53:ILE:O	1:A:61:MET:N	2.50	0.45
1:B:19:HIS:O	1:B:21:HIS:N	2.50	0.45
1:A:129:LEU:HD23	1:B:75:LEU:HD11	1.99	0.45
1:B:188:VAL:O	1:B:192:LEU:HG	2.17	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:106:LYS:O	1:A:110:LEU:HG	2.19	0.42
1:B:162:MET:N	1:B:169:GLN:O	2.50	0.42
1:A:72:PHE:CE1	1:A:91:VAL:HG11	2.55	0.41
1:A:74:GLU:N	1:A:74:GLU:OE2	2.50	0.41
1:A:67:ASN:O	1:A:70:ASP:HB2	2.19	0.41
1:A:114:ASP:OD2	1:A:116:GLU:HG2	2.20	0.41
1:A:114:ASP:OD2	1:A:115:SER:N	2.54	0.41
1:A:159:PRO:O	1:A:160:ASP:HB2	2.20	0.41
1:B:117:ILE:HD12	1:B:117:ILE:HA	1.96	0.41
1:B:151:THR:HA	1:B:154:ASP:HB2	2.02	0.41
1:B:201:VAL:HA	1:B:209:VAL:O	2.20	0.41
1:B:38:CYS:SG	1:B:87:ARG:NH1	2.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	193/210 (92%)	180 (93%)	13 (7%)	0	100	100
1	B	182/210 (87%)	173 (95%)	6 (3%)	3 (2%)	9	31
All	All	375/420 (89%)	353 (94%)	19 (5%)	3 (1%)	19	49

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	21	HIS
1	B	39	GLU
1	B	20	CYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	145/181 (80%)	143 (99%)	2 (1%)	67	90
1	B	132/181 (73%)	131 (99%)	1 (1%)	81	94
All	All	277/362 (76%)	274 (99%)	3 (1%)	73	92

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	GLU
1	A	190	ARG
1	B	56	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	202	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 7 are monoatomic - leaving 8 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.



There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	197/210 (93%)	0.47	13 (6%) 18 11	19, 37, 64, 75	0
1	B	188/210 (89%)	0.57	19 (10%) 7 4	21, 41, 65, 76	0
All	All	385/420 (91%)	0.52	32 (8%) 11 6	19, 39, 64, 76	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	212	GLY	6.9
1	B	207	THR	5.0
1	A	162	MET	4.9
1	B	101	GLU	4.8
1	A	153	LEU	4.0
1	B	200	LEU	3.5
1	B	205	GLY	3.4
1	B	157	GLN	3.4
1	A	202	HIS	3.3
1	A	206	LYS	3.2
1	A	113	GLN	3.2
1	B	38	CYS	2.8
1	B	154	ASP	2.8
1	A	201	VAL	2.7
1	B	112	GLN	2.7
1	A	184	SER	2.6
1	A	56	ASP	2.6
1	B	206	LYS	2.5
1	A	160	ASP	2.5
1	B	163	THR	2.5
1	B	208	MET	2.5
1	A	57	ASP	2.5
1	B	19	HIS	2.3
1	B	176	GLU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	207	THR	2.2
1	B	117	ILE	2.2
1	B	54	GLU	2.1
1	B	20	CYS	2.1
1	B	21	HIS	2.0
1	B	48	SER	2.0
1	A	157	GLN	2.0
1	A	169	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ


There are no monosaccharides in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	AUF	B	303	8/8	0.84	0.31	15,40,64,95	0
3	AUF	A	302	8/8	0.88	0.23	33,45,64,106	0
4	AU	B	304	1/1	0.90	0.08	135,135,135,135	0
5	CL	A	305	1/1	0.91	0.13	70,70,70,70	0
6	SO4	A	306	5/5	0.92	0.20	36,39,46,47	0
2	CMP	A	301	22/22	0.94	0.18	12,23,30,34	0
6	SO4	B	308	5/5	0.95	0.11	35,36,43,46	0
2	CMP	B	302	22/22	0.96	0.14	15,24,33,35	0
6	SO4	A	307	5/5	0.96	0.14	46,53,59,65	0
6	SO4	B	307	5/5	0.96	0.18	32,44,48,51	0
4	AU	A	303	1/1	0.96	0.06	109,109,109,109	0
4	AU	B	306	1/1	0.97	0.05	85,85,85,85	0
4	AU	A	304	1/1	0.98	0.05	65,65,65,65	0
4	AU	B	305	1/1	0.99	0.03	71,71,71,71	0
5	CL	B	301	1/1	0.99	0.30	70,70,70,70	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around AUF A 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around AU B 304:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around CL A 305:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around CMP A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around CMP B 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around AU A 303:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around AU B 306:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around AU A 304:**


$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around AU B 305:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers ⓘ

There are no such residues in this entry.