



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 12, 2021 – 08:02 AM EDT

PDB ID : 2FGK  
Title : Crystal structure of the ABC-cassette E631Q mutant of HlyB with bound ATP  
Authors : Zaitseva, J.; Oswald, C.; Jumpertz, T.; Jenewein, S.; Holland, I.B.; Schmitt, L.  
Deposited on : 2005-12-22  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	
1	B	247	
1	C	247	
1	D	247	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7848 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-hemolysin translocation ATP-binding protein hlyB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	241	Total	C	N	O	S	0	0	0
			1893	1194	342	352	5			
1	B	241	Total	C	N	O	S	0	0	0
			1893	1194	342	352	5			
1	C	241	Total	C	N	O	S	0	0	0
			1893	1194	342	352	5			
1	D	241	Total	C	N	O	S	0	0	0
			1893	1194	342	352	5			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	461	HIS	-	expression tag	UNP P08716
A	462	HIS	-	expression tag	UNP P08716
A	463	HIS	-	expression tag	UNP P08716
A	464	HIS	-	expression tag	UNP P08716
A	465	HIS	-	expression tag	UNP P08716
A	466	HIS	-	expression tag	UNP P08716
A	631	GLN	GLU	engineered mutation	UNP P08716
B	461	HIS	-	expression tag	UNP P08716
B	462	HIS	-	expression tag	UNP P08716
B	463	HIS	-	expression tag	UNP P08716
B	464	HIS	-	expression tag	UNP P08716
B	465	HIS	-	expression tag	UNP P08716
B	466	HIS	-	expression tag	UNP P08716
B	631	GLN	GLU	engineered mutation	UNP P08716
C	461	HIS	-	expression tag	UNP P08716
C	462	HIS	-	expression tag	UNP P08716
C	463	HIS	-	expression tag	UNP P08716
C	464	HIS	-	expression tag	UNP P08716
C	465	HIS	-	expression tag	UNP P08716
C	466	HIS	-	expression tag	UNP P08716
C	631	GLN	GLU	engineered mutation	UNP P08716

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	461	HIS	-	expression tag	UNP P08716
D	462	HIS	-	expression tag	UNP P08716
D	463	HIS	-	expression tag	UNP P08716
D	464	HIS	-	expression tag	UNP P08716
D	465	HIS	-	expression tag	UNP P08716
D	466	HIS	-	expression tag	UNP P08716
D	631	GLN	GLU	engineered mutation	UNP P08716

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	36	Total	O	0	0
			36	36		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	44	Total 44	O 44	0	0
3	C	32	Total 32	O 32	0	0
3	D	40	Total 40	O 40	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Alpha-hemolysin translocation ATP-binding protein hlyB




- Molecule 1: Alpha-hemolysin translocation ATP-binding protein hlyB





- Molecule 1: Alpha-hemolysin translocation ATP-binding protein hlyB





● Molecule 1: Alpha-hemolysin translocation ATP-binding protein hlyB



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.14Å 189.26Å 63.48Å 90.00° 111.87° 90.00°	Depositor
Resolution (Å)	20.00 – 2.70 20.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	98.9 (20.00-2.70) 98.8 (20.00-2.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.53 (at 2.71Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.223 , 0.280 0.222 , 0.280	Depositor DCC
$R_{free}$ test set	1406 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	49.7	Xtriage
Anisotropy	0.062	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 47.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.406 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7848	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.27	8/1920 (0.4%)	0.68	4/2589 (0.2%)
1	B	0.41	0/1920	0.58	0/2589
1	C	0.73	8/1920 (0.4%)	0.67	4/2589 (0.2%)
1	D	0.41	0/1920	0.57	0/2589
All	All	0.79	16/7680 (0.2%)	0.63	8/10356 (0.1%)

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	593	GLU	CD-OE2	43.26	1.73	1.25
1	A	593	GLU	CD-OE1	18.30	1.45	1.25
1	C	558	SER	CB-OG	14.44	1.61	1.42
1	A	595	TYR	CE2-CZ	13.03	1.55	1.38
1	C	595	TYR	CE2-CZ	12.90	1.55	1.38

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	557	ARG	NE-CZ-NH2	-10.21	115.19	120.30
1	C	557	ARG	NE-CZ-NH2	-9.98	115.31	120.30
1	A	593	GLU	OE1-CD-OE2	9.53	134.74	123.30
1	C	595	TYR	CB-CG-CD2	-6.88	116.87	121.00
1	A	595	TYR	CB-CG-CD2	-6.80	116.92	121.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1893	0	1949	29	0
1	B	1893	0	1949	36	0
1	C	1893	0	1949	31	0
1	D	1893	0	1949	37	0
2	A	31	0	12	0	0
2	B	31	0	12	1	0
2	C	31	0	12	0	0
2	D	31	0	12	0	0
3	A	36	0	0	5	0
3	B	44	0	0	2	0
3	C	32	0	0	3	0
3	D	40	0	0	1	0
All	All	7848	0	7844	130	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 130 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:593:GLU:CD	1:A:593:GLU:OE2	1.73	1.25
1:C:571:SER:HB2	1:C:574:LYS:HB2	1.36	1.06
1:B:612:GLN:HE22	1:B:635:ALA:H	1.13	0.90
1:C:560:ILE:HB	3:C:139:HOH:O	1.73	0.88
1:C:612:GLN:HE22	1:C:635:ALA:H	1.27	0.83

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/247 (97%)	223 (93%)	10 (4%)	6 (2%)	5	14
1	B	239/247 (97%)	226 (95%)	10 (4%)	3 (1%)	12	30
1	C	239/247 (97%)	222 (93%)	11 (5%)	6 (2%)	5	14
1	D	239/247 (97%)	224 (94%)	13 (5%)	2 (1%)	19	43
All	All	956/988 (97%)	895 (94%)	44 (5%)	17 (2%)	8	21

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	603	GLY
1	C	601	GLU
1	C	571	SER
1	D	565	LEU
1	A	604	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	207/213 (97%)	192 (93%)	15 (7%)	14	34
1	B	207/213 (97%)	196 (95%)	11 (5%)	22	48
1	C	207/213 (97%)	194 (94%)	13 (6%)	18	40
1	D	207/213 (97%)	199 (96%)	8 (4%)	32	61
All	All	828/852 (97%)	781 (94%)	47 (6%)	20	44

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	556	ASN
1	C	664	LEU
1	C	586	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	613	ARG
1	C	682	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	539	ASN
1	D	642	HIS
1	C	623	ASN
1	D	602	GLN
1	C	612	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ATP	B	801	-	26,33,33	1.19	2 (7%)	31,52,52	1.53	4 (12%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ATP	A	800	-	26,33,33	1.14	2 (7%)	31,52,52	1.53	4 (12%)
2	ATP	C	802	-	26,33,33	0.95	1 (3%)	31,52,52	1.62	4 (12%)
2	ATP	D	803	-	26,33,33	1.20	2 (7%)	31,52,52	1.51	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	B	801	-	-	3/18/38/38	0/3/3/3
2	ATP	A	800	-	-	2/18/38/38	0/3/3/3
2	ATP	C	802	-	-	3/18/38/38	0/3/3/3
2	ATP	D	803	-	-	3/18/38/38	0/3/3/3

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	ATP	PG-O1G	3.52	1.61	1.50
2	D	803	ATP	PG-O1G	3.34	1.61	1.50
2	A	800	ATP	PG-O1G	3.14	1.60	1.50
2	D	803	ATP	O4'-C1'	2.95	1.45	1.41
2	A	800	ATP	O4'-C1'	2.84	1.45	1.41

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	802	ATP	N3-C2-N1	-4.55	121.57	128.68
2	A	800	ATP	N3-C2-N1	-4.33	121.91	128.68
2	D	803	ATP	N3-C2-N1	-4.19	122.13	128.68
2	B	801	ATP	N3-C2-N1	-4.15	122.20	128.68
2	A	800	ATP	O3G-PG-O3B	4.06	118.24	104.64

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	801	ATP	O4'-C4'-C5'-O5'
2	B	801	ATP	C3'-C4'-C5'-O5'
2	A	800	ATP	PA-O3A-PB-O1B

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	D	803	ATP	O4'-C4'-C5'-O5'
2	C	802	ATP	PA-O3A-PB-O1B


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	ATP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	241/247 (97%)	1.26	48 (19%) <b>1</b> <b>0</b>	39, 51, 62, 71	0
1	B	241/247 (97%)	1.10	40 (16%) <b>1</b> <b>1</b>	40, 51, 63, 70	0
1	C	241/247 (97%)	1.24	42 (17%) <b>1</b> <b>1</b>	37, 50, 61, 71	0
1	D	241/247 (97%)	1.14	43 (17%) <b>1</b> <b>1</b>	40, 51, 63, 71	0
All	All	964/988 (97%)	1.19	173 (17%) <b>1</b> <b>1</b>	37, 51, 63, 71	0

The worst 5 of 173 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	604	ALA	14.3
1	D	707	ASP	10.4
1	B	550	GLN	9.1
1	C	551	ASP	9.0
1	A	569	GLY	8.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ATP	A	800	31/31	0.91	0.19	32,41,44,44	0
2	ATP	C	802	31/31	0.92	0.17	28,40,43,44	0
2	ATP	D	803	31/31	0.92	0.24	38,46,47,50	0
2	ATP	B	801	31/31	0.93	0.16	37,42,44,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around ATP C 802:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around ATP D 803:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around ATP B 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.