



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2021 – 04:15 PM EDT

PDB ID : 3FIR  
Title : Crystal structure of Glycosylated K135E PEB3  
Authors : Min, T.; Matte, A.; Cygler, M.  
Deposited on : 2008-12-12  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	251	<div> <div>2%</div> <div> <div></div> <div>79%</div> <div>11%</div> <div>8%</div> </div> </div>
1	B	251	<div> <div>8%</div> <div> <div></div> <div>75%</div> <div>14%</div> <div>8%</div> </div> </div>
2	C	2	<div> <div>50%</div> <div>50%</div> </div>
2	D	2	<div> <div>50%</div> <div>50%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NGZ	C	2	X	-	-	-
2	NGZ	D	2	X	-	-	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4083 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Major antigenic peptide PEB3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	231	Total	C	N	O	S	0	0	0
			1811	1159	310	341	1			
1	B	231	Total	C	N	O	S	0	0	0
			1811	1159	310	341	1			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	135	GLU	LYS	engineered mutation	UNP Q0PBL7
A	251	HIS	-	expression tag	UNP Q0PBL7
B	135	GLU	LYS	engineered mutation	UNP Q0PBL7
B	251	HIS	-	expression tag	UNP Q0PBL7

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy- $\alpha$ -L-glucopyranose-(1-3)-2,4-bisacetamido-2,4,6-trideoxy- $\beta$ -D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			29	18	3	8			
2	D	2	Total	C	N	O	0	0	0
			29	18	3	8			

- Molecule 3 is CITRATE ANION (three-letter code: FLC) (formula: C<sub>6</sub>H<sub>5</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		
3	B	1	Total	C	O	0	0
			13	6	7		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	229	Total	O	0	0
			229	229		
4	B	148	Total	O	0	0
			148	148		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Major antigenic peptide PEB3



- Molecule 1: Major antigenic peptide PEB3



- Molecule 2: 2-acetamido-2-deoxy-alpha-L-glucopyranose-(1-3)-2,4-bisacetamido-2,4,6-trideoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-alpha-L-glucopyranose-(1-3)-2,4-bisacetamido-2,4,6-trideoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.58Å 101.96Å 56.81Å 90.00° 108.88° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 19.94 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.5 (20.00-2.00) 97.4 (19.94-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.54 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.4.0067	Depositor
R, $R_{free}$	0.216 , 0.235 0.214 , 0.230	Depositor DCC
$R_{free}$ test set	1756 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.6	Xtriage
Anisotropy	0.833	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 47.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4083	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: B6D, FLC, NGZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.31	0/1850	0.51	0/2499
1	B	0.31	0/1850	0.52	0/2499
All	All	0.31	0/3700	0.52	0/4998

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1811	0	1810	18	0
1	B	1811	0	1810	24	0
2	C	29	0	26	0	0
2	D	29	0	26	1	0
3	A	13	0	5	0	0
3	B	13	0	5	1	0
4	A	229	0	0	1	0
4	B	148	0	0	0	0
All	All	4083	0	3682	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.



All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23:ASN:ND2	1:B:66:ASP:OD2	2.12	0.81
1:B:90:ASN:OD1	1:B:92:SER:HB2	1.81	0.80
1:A:181:ASP:OD2	1:A:181:ASP:N	2.27	0.66
1:A:180:GLN:HA	1:A:186:TRP:CZ2	2.38	0.58
1:A:180:GLN:HA	1:A:186:TRP:HZ2	1.69	0.57
3:B:301:FLC:OA2	3:B:301:FLC:OB1	2.22	0.56
1:B:113:LYS:O	1:B:115:LYS:HE2	2.06	0.55
1:A:130:PRO:HA	1:A:168:VAL:O	2.08	0.54
1:B:35:LYS:HE3	1:B:53:PHE:HE1	1.72	0.54
1:B:222:GLY:O	1:B:223:ALA:C	2.47	0.53
1:B:28:GLY:HA3	1:B:173:SER:HB3	1.91	0.52
1:A:126:ARG:HB3	1:A:165:VAL:HG11	1.93	0.51
1:B:44:LYS:CD	1:B:45:THR:HG23	2.39	0.51
1:B:237:LYS:HA	1:B:237:LYS:HE3	1.93	0.49
1:A:161:ARG:HB2	4:A:411:HOH:O	2.12	0.48
1:B:235:SER:O	1:B:240:LYS:HE3	2.14	0.48
1:A:208:ASP:OD1	1:A:208:ASP:N	2.43	0.48
1:B:25:TYR:CD1	1:B:63:ALA:HB2	2.49	0.48
1:B:215:PHE:HB2	1:B:248:TRP:CZ3	2.50	0.47
1:A:177:LEU:HB2	1:A:180:GLN:HB2	1.97	0.47
1:A:26:GLY:O	1:A:53:PHE:HA	2.15	0.46
1:B:23:ASN:OD1	1:B:50:ASN:ND2	2.46	0.45
1:A:174:ALA:C	1:A:176:LYS:H	2.18	0.45
1:A:101:GLU:O	1:A:101:GLU:HG2	2.15	0.45
1:B:219:ALA:HB2	1:B:231:ILE:CD1	2.46	0.45
1:B:177:LEU:O	1:B:180:GLN:HB3	2.17	0.45
1:B:219:ALA:HB2	1:B:231:ILE:HD11	1.99	0.45
2:D:1:B6D:H8B	2:D:2:NGZ:H3	2.00	0.44
1:A:225:LYS:HD3	1:A:229:ASP:OD2	2.18	0.43
1:A:215:PHE:HB2	1:A:248:TRP:CZ3	2.53	0.43
1:B:225:LYS:O	1:B:229:ASP:OD1	2.37	0.42
1:A:236:SER:OG	1:A:238:GLU:HG2	2.19	0.42
1:B:24:LEU:HG	1:B:69:ILE:HB	2.02	0.42
1:B:130:PRO:HA	1:B:168:VAL:O	2.20	0.41
1:B:225:LYS:HD2	1:B:229:ASP:OD1	2.21	0.41
1:A:62:LYS:O	1:A:66:ASP:OD1	2.38	0.41
1:A:53:PHE:HE2	1:A:173:SER:HB2	1.86	0.41
1:A:188:THR:OG1	1:A:189:TRP:N	2.53	0.41
1:B:60:PHE:CE1	1:B:85:PHE:CZ	3.09	0.41
1:A:98:TYR:HB3	1:A:248:TRP:HB3	2.02	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:44:LYS:O	1:B:44:LYS:HG2	2.20	0.41
1:B:44:LYS:HD2	1:B:45:THR:HG23	2.02	0.40
1:B:105:LEU:HD23	1:B:177:LEU:HD21	2.04	0.40
1:B:70:LEU:O	1:B:217:VAL:HA	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/251 (91%)	222 (97%)	6 (3%)	1 (0%)	34	30
1	B	229/251 (91%)	226 (99%)	2 (1%)	1 (0%)	34	30
All	All	458/502 (91%)	448 (98%)	8 (2%)	2 (0%)	34	30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	223	ALA
1	A	172	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	190/204 (93%)	181 (95%)	9 (5%)	26	22
1	B	190/204 (93%)	180 (95%)	10 (5%)	22	18
All	All	380/408 (93%)	361 (95%)	19 (5%)	24	20

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	87	LYS
1	A	100	ARG
1	A	119	ASP
1	A	180	GLN
1	A	181	ASP
1	A	208	ASP
1	A	225	LYS
1	A	237	LYS
1	A	238	GLU
1	B	43	GLU
1	B	44	LYS
1	B	87	LYS
1	B	100	ARG
1	B	115	LYS
1	B	119	ASP
1	B	225	LYS
1	B	229	ASP
1	B	237	LYS
1	B	238	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	162	ASN
1	A	182	GLN
1	B	162	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	B6D	C	1	2,1	15,16,17	1.05	1 (6%)	16,22,24	2.15	4 (25%)
2	NGZ	C	2	2	13,13,15	0.43	0	14,17,21	0.91	1 (7%)
2	B6D	D	1	2,1	15,16,17	1.12	1 (6%)	16,22,24	2.66	6 (37%)
2	NGZ	D	2	2	13,13,15	0.54	0	14,17,21	1.14	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B6D	C	1	2,1	-	2/8/25/28	0/1/1/1
2	NGZ	C	2	2	4/4/4/7	2/6/19/26	0/1/1/1
2	B6D	D	1	2,1	-	4/8/25/28	0/1/1/1
2	NGZ	D	2	2	4/4/4/7	2/6/19/26	1/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	B6D	O3-C3	-3.87	1.33	1.43
2	D	1	B6D	O3-C3	-3.86	1.33	1.43

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	B6D	O3-C3-C4	6.85	123.49	109.66
2	D	1	B6D	O3-C3-C2	5.43	120.70	109.47
2	C	1	B6D	O3-C3-C2	4.65	119.09	109.47
2	C	1	B6D	O3-C3-C4	4.36	118.45	109.66
2	C	1	B6D	C1-C2-N2	-4.10	103.48	110.49
2	C	1	B6D	C3-C4-N4	-3.62	103.77	110.62
2	D	1	B6D	C1-C2-N2	-3.27	104.90	110.49
2	D	1	B6D	C4-N4-C10	3.03	130.54	123.18
2	D	2	NGZ	C1-O5-C5	2.60	119.17	113.51
2	D	1	B6D	C2-N2-C7	-2.22	119.74	122.90
2	C	2	NGZ	C1-O5-C5	2.06	117.98	113.51
2	D	2	NGZ	O5-C5-C4	2.06	113.98	110.65
2	D	1	B6D	O5-C1-C2	-2.05	108.05	111.29

All (8) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	2	NGZ	C2
2	C	2	NGZ	C1
2	C	2	NGZ	C3
2	C	2	NGZ	C5
2	D	2	NGZ	C2
2	D	2	NGZ	C1
2	D	2	NGZ	C3
2	D	2	NGZ	C5

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	2	NGZ	C8-C7-N2-C2
2	C	2	NGZ	O7-C7-N2-C2
2	D	1	B6D	C3-C4-N4-C10
2	D	2	NGZ	C8-C7-N2-C2
2	D	2	NGZ	O7-C7-N2-C2
2	C	1	B6D	C8-C7-N2-C2
2	D	1	B6D	C8-C7-N2-C2
2	C	1	B6D	O7-C7-N2-C2
2	D	1	B6D	O7-C7-N2-C2
2	D	1	B6D	C5-C4-N4-C10


All (1) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	2	NGZ	C1-C2-C3-C4-C5-O5

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	B6D	1	0
2	D	2	NGZ	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	FLC	B	301	-	3,12,12	0.80	0	3,17,17	1.68	0
3	FLC	A	301	-	3,12,12	0.65	0	3,17,17	0.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FLC	B	301	-	-	1/6/16/16	-
3	FLC	A	301	-	-	3/6/16/16	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	301	FLC	CAC-CA-CB-OHB
3	A	301	FLC	CAC-CA-CB-CG
3	A	301	FLC	CAC-CA-CB-CBC
3	B	301	FLC	CAC-CA-CB-OHB

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	301	FLC	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	231/251 (92%)	0.10	4 (1%) 70 68	20, 32, 42, 51	0
1	B	231/251 (92%)	0.43	20 (8%) 10 9	22, 35, 56, 60	0
All	All	462/502 (92%)	0.27	24 (5%) 27 26	20, 33, 53, 60	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	176	LYS	4.4
1	B	223	ALA	4.3
1	B	87	LYS	4.3
1	B	175	ARG	4.1
1	B	39	ASN	3.9
1	B	43	GLU	3.7
1	A	175	ARG	3.6
1	B	225	LYS	3.2
1	B	176	LYS	3.2
1	A	135	GLU	3.0
1	B	65	LYS	2.9
1	B	241	GLU	2.8
1	B	21	ASP	2.6
1	B	228	GLN	2.6
1	B	49	VAL	2.4
1	B	229	ASP	2.3
1	B	47	VAL	2.2
1	B	42	SER	2.2
1	B	41	TYR	2.2
1	B	135	GLU	2.1
1	A	251	HIS	2.1
1	B	251	HIS	2.1
1	B	185	ALA	2.0
1	B	30	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NGZ	D	2	13/15	0.70	0.21	57,57,58,58	13
2	NGZ	C	2	13/15	0.76	0.18	47,47,47,48	13
2	B6D	D	1	16/17	0.80	0.20	53,54,55,56	16
2	B6D	C	1	16/17	0.83	0.20	44,44,45,46	16

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	FLC	A	301	13/13	0.91	0.13	37,37,38,38	0
3	FLC	B	301	13/13	0.93	0.12	40,41,41,42	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.