



Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 11:00 am BST

PDB ID : 5FI1
Title : Crystal Structure of the P-Rex1 DH/PH tandem in complex with Cdc42
Authors : Cash, J.N.; Tesmer, J.J.G.
Deposited on : 2015-12-22
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	<div> <div>6%</div> <div> <div></div> <div>81%</div> <div>11%</div> <div>8%</div> </div> </div>
2	B	194	<div> <div></div> <div> <div>81%</div> <div>11%</div> <div>7%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4194 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 3,4,5-trisphosphate-dependent Rac exchanger 1 protein, Phosphatidylinositol 3,4,5-trisphosphate-dependent Rac exchanger 1 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	344	Total	C	N	O	S	0	0	0
			2788	1767	490	516	15			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q8TCU6
A	-1	GLU	-	expression tag	UNP Q8TCU6
A	0	PHE	-	expression tag	UNP Q8TCU6

- Molecule 2 is a protein called Cell division control protein 42 homolog.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	180	Total	C	N	O	S	0	0	0
			1406	907	223	269	7			


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	expression tag	UNP P60953
B	-1	GLU	-	expression tag	UNP P60953
B	0	PHE	-	expression tag	UNP P60953


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol 3,4,5-trisphosphate-dependent Rac exchanger 1 protein, Phosphatidylinositol 3,4,5-trisphosphate-dependent Rac exchanger 1 protein



- Molecule 2: Cell division control protein 42 homolog



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	67.31Å 95.91Å 277.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.12 – 3.20 43.12 – 3.20	Depositor EDS
% Data completeness (in resolution range)	81.2 (43.12-3.20) 77.2 (43.12-3.20)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.41 (at 3.19Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.205 , 0.198 0.211 , 0.201	Depositor DCC
R_{free} test set	607 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	79.5	Xtriage
Anisotropy	0.659	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 64.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4194	wwPDB-VP
Average B, all atoms (Å ²)	124.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/2837	0.40	0/3823
2	B	0.23	0/1438	0.43	0/1958
All	All	0.24	0/4275	0.41	0/5781

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2788	0	2814	23	0
2	B	1406	0	1421	12	0
All	All	4194	0	4235	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:44:VAL:HG21	2:B:173:ILE:HD13	1.69	0.75
1:A:284:GLY:HA2	1:A:351:ASN:HD21	1.51	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:ARG:NH1	1:A:213:THR:OG1	2.31	0.64
1:A:61:GLU:OE1	1:A:202:TYR:OH	2.15	0.63
2:B:53:LEU:HD22	2:B:173:ILE:HD11	1.86	0.56
1:A:56:GLU:OE2	2:B:32:TYR:OH	2.15	0.56
1:A:297:LEU:HA	1:A:333:THR:HG23	1.88	0.55
1:A:202:TYR:HA	1:A:205:LEU:HD12	1.87	0.54
1:A:254:GLN:HG3	1:A:265:LEU:HD12	1.91	0.53
2:B:83:SER:HB3	2:B:86:SER:HB3	1.94	0.49
1:A:162:LEU:HB3	1:A:168:VAL:HG11	1.94	0.49
1:A:51:LEU:O	1:A:55:ASN:ND2	2.41	0.47
2:B:62:GLU:O	2:B:68:ARG:NH1	2.48	0.47
1:A:362:HIS:CE1	1:A:364:THR:HG22	2.51	0.46
2:B:9:VAL:HG12	2:B:58:THR:HG21	1.97	0.46
1:A:245:GLU:OE1	2:B:66:ARG:NH1	2.41	0.45
2:B:98:VAL:HG21	2:B:149:LEU:HD13	1.97	0.45
1:A:289:ARG:HE	1:A:328:ARG:HG2	1.82	0.44
2:B:35:THR:HG23	2:B:36:VAL:HG23	1.99	0.44
1:A:131:LEU:O	1:A:134:VAL:HG22	2.18	0.44
1:A:169:ARG:O	1:A:173:LEU:HG	2.19	0.43
1:A:362:HIS:HB2	1:A:369:TRP:CZ3	2.55	0.42
2:B:8:VAL:HG22	2:B:79:LEU:HD12	2.00	0.42
1:A:196:ILE:HG13	1:A:197:GLN:N	2.35	0.42
1:A:57:ILE:HG12	1:A:205:LEU:HD22	2.01	0.42
1:A:268:ILE:O	1:A:296:ASN:ND2	2.50	0.42
1:A:68:LEU:HB2	1:A:107:LEU:HD13	2.02	0.42
1:A:338:VAL:HG21	1:A:387:ILE:HD13	2.02	0.41
2:B:116:GLN:HB3	2:B:119:LEU:HD12	2.02	0.41
1:A:131:LEU:HA	1:A:131:LEU:HD23	1.91	0.41
1:A:360:LYS:HE2	1:A:369:TRP:CE3	2.56	0.41
2:B:65:ASP:N	2:B:65:ASP:OD1	2.54	0.41
1:A:344:GLY:N	1:A:355:VAL:O	2.50	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/374 (91%)	329 (97%)	11 (3%)	0	100	100
2	B	178/194 (92%)	173 (97%)	5 (3%)	0	100	100
All	All	518/568 (91%)	502 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	306/332 (92%)	303 (99%)	3 (1%)	76	90
2	B	160/173 (92%)	157 (98%)	3 (2%)	57	81
All	All	466/505 (92%)	460 (99%)	6 (1%)	69	87

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	HIS
1	A	94	GLU
1	A	188	LEU
2	B	11	ASP
2	B	31	GLU
2	B	66	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	351	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	344/374 (91%)	0.20	24 (6%)	16	9	64, 106, 224, 248	0
2	B	180/194 (92%)	-0.12	0	100	100	71, 103, 159, 198	0
All	All	524/568 (92%)	0.09	24 (4%)	32	20	64, 104, 215, 248	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	383	TRP	4.8
1	A	361	ILE	4.7
1	A	292	PHE	4.0
1	A	372	CYS	3.6
1	A	384	LEU	3.4
1	A	359	TRP	3.3
1	A	294	PHE	3.2
1	A	336	MET	2.8
1	A	337	GLU	2.8
1	A	340	ASN	2.8
1	A	355	VAL	2.8
1	A	345	THR	2.8
1	A	338	VAL	2.7
1	A	370	PHE	2.7
1	A	278	LEU	2.6
1	A	90	GLY	2.6
1	A	298	LEU	2.6
1	A	380	LYS	2.5
1	A	260	TRP	2.5
1	A	331	ILE	2.4
1	A	274	LEU	2.4
1	A	270	THR	2.3
1	A	293	LEU	2.1
1	A	297	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.