



wwPDB X-ray Structure Validation Summary Report ⓘ

May 18, 2020 – 12:50 am BST

PDB ID : 1FN3
Title : CRYSTAL STRUCTURE OF NICKEL RECONSTITUTED HEMOGLOBIN-
A CASE FOR PERMANENT, T-STATE HEMOGLOBIN
Authors : Venkateshrao, S.; Deepthi, S.; Pattabhi, V.; Manoharan, P.T.
Deposited on : 2000-08-20
Resolution : 2.48 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6594 (2.50-2.46)
Ramachandran outliers	138981	6469 (2.50-2.46)
Sidechain outliers	138945	6471 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	141	
1	C	141	
2	B	146	
2	D	146	

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 2 is a protein called HEMOGLOBIN BETA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	146	Total 1123	C 724	N 195	O 201	S 3	0	0	0
2	D	146	Total 1123	C 724	N 195	O 201	S 3	0	0	0

- # HNI

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 43	C 34	N 4	Ni 1	O 4	0	0
3	B	1	Total 43	C 34	N 4	Ni 1	O 4	0	0
3	C	1	Total 43	C 34	N 4	Ni 1	O 4	0	0
3	D	1	Total 43	C 34	N 4	Ni 1	O 4	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	17	Total 17	O 17	0	0
4	B	7	Total 7	O 7	0	0
4	C	9	Total 9	O 9	0	0
4	D	5	Total 5	O 5	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: HEMOGLOBIN ALPHA CHAIN



• Molecule 1: HEMOGLOBIN ALPHA CHAIN





• Molecule 2: HEMOGLOBIN BETA CHAIN



• Molecule 2: HEMOGLOBIN BETA CHAIN

Chain D:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	95.52Å 96.14Å 65.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.48 19.80 – 2.48	Depositor EDS
% Data completeness (in resolution range)	(Not available) (8.00-2.48) 19.4 (19.80-2.48)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.00 (at 2.50Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.242 , 0.324 0.321 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	35.8	Xtriage
Anisotropy	0.558	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 52.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.57$, $\langle L^2 \rangle = 0.42$	Xtriage
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	4594	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HNI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	3/1097 (0.3%)	1.01	6/1491 (0.4%)
1	C	0.84	4/1097 (0.4%)	1.23	14/1491 (0.9%)
2	B	0.66	1/1153 (0.1%)	1.24	10/1566 (0.6%)
2	D	0.73	2/1153 (0.2%)	1.33	9/1566 (0.6%)
All	All	0.73	10/4500 (0.2%)	1.21	39/6114 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
2	B	0	1
2	D	1	0
All	All	1	2

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	86	LEU	CA-C	10.55	1.80	1.52
1	A	88	ALA	CA-CB	9.38	1.72	1.52
2	D	59	LYS	CE-NZ	-8.30	1.28	1.49
2	B	59	LYS	CE-NZ	-7.93	1.29	1.49
2	D	95	LYS	CA-CB	7.18	1.69	1.53

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	40	ARG	NE-CZ-NH2	-19.20	110.70	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	104	ARG	NE-CZ-NH2	17.38	128.99	120.30
2	D	104	ARG	NE-CZ-NH1	-16.92	111.84	120.30
2	D	40	ARG	NE-CZ-NH1	16.32	128.46	120.30
2	B	40	ARG	NE-CZ-NH2	16.22	128.41	120.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	D	94	ASP	CA

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	63	HIS	Sidechain
1	C	86	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1069	0	1073	158	0
1	C	1069	0	1073	154	0
2	B	1123	0	1118	154	0
2	D	1123	0	1118	159	0
3	A	43	0	30	7	0
3	B	43	0	30	2	0
3	C	43	0	30	13	0
3	D	43	0	30	20	0
4	A	17	0	0	32	0
4	B	7	0	0	13	0
4	C	9	0	0	7	0
4	D	5	0	0	16	0
All	All	4594	0	4502	601	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 66.

The worst 5 of 601 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:86:LEU:CA	1:C:86:LEU:C	1.80	1.45
2:D:137:VAL:HG21	4:D:150:HOH:O	1.24	1.28
1:A:46:PHE:HA	4:A:153:HOH:O	1.30	1.25
1:C:16:LYS:HE2	4:C:150:HOH:O	1.30	1.25
2:B:115:ALA:HB2	4:B:154:HOH:O	1.24	1.24

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/141 (99%)	109 (78%)	20 (14%)	10 (7%)	1	1
1	C	139/141 (99%)	108 (78%)	22 (16%)	9 (6%)	1	1
2	B	144/146 (99%)	118 (82%)	20 (14%)	6 (4%)	3	3
2	D	144/146 (99%)	117 (81%)	21 (15%)	6 (4%)	3	3
All	All	566/574 (99%)	452 (80%)	83 (15%)	31 (6%)	2	1

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	46	PHE
1	A	53	ALA
1	A	86	LEU
2	B	14	LEU
2	B	49	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/113 (100%)	72 (64%)	41 (36%)	0	0
1	C	113/113 (100%)	78 (69%)	35 (31%)	0	0
2	B	118/118 (100%)	74 (63%)	44 (37%)	0	0
2	D	118/118 (100%)	73 (62%)	45 (38%)	0	0
All	All	462/462 (100%)	297 (64%)	165 (36%)	0	0

5 of 165 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	108	ASN
1	C	30	GLU
2	D	96	LEU
2	B	120	LYS
2	B	141	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	131	GLN
2	B	139	ASN
2	D	108	ASN
2	B	116	HIS
2	B	127	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HNI	A	142	-	33,50,50	4.06	17 (51%)	22,82,82	5.64	11 (50%)
3	HNI	D	147	2	33,50,50	4.29	17 (51%)	22,82,82	5.87	14 (63%)
3	HNI	C	142	-	33,50,50	4.17	18 (54%)	22,82,82	5.70	11 (50%)
3	HNI	B	147	-	33,50,50	4.33	17 (51%)	22,82,82	5.83	13 (59%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HNI	A	142	-	-	4/8/94/94	-
3	HNI	D	147	2	-	3/8/94/94	-
3	HNI	C	142	-	-	4/8/94/94	-
3	HNI	B	147	-	-	3/8/94/94	-

The worst 5 of 69 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	142	HNI	CHB-C4A	-10.51	1.35	1.53
3	A	142	HNI	CHB-C4A	-10.42	1.35	1.53
3	B	147	HNI	CHB-C4A	-9.85	1.36	1.53
3	D	147	HNI	CHB-C4A	-9.69	1.36	1.53
3	D	147	HNI	CHD-C4C	-9.34	1.37	1.53

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	147	HNI	CHD-C4C-NC	12.98	126.46	110.03
3	D	147	HNI	CHD-C4C-NC	12.92	126.38	110.03
3	C	142	HNI	CHD-C4C-NC	12.62	126.00	110.03
3	A	142	HNI	CHD-C4C-NC	12.41	125.74	110.03
3	C	142	HNI	CHA-C1A-NA	12.01	125.23	110.03

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	142	HNI	C2C-C3C-CAC-CBC
3	A	142	HNI	C4C-C3C-CAC-CBC
3	D	147	HNI	C1A-C2A-CAA-CBA
3	C	142	HNI	C2C-C3C-CAC-CBC
3	C	142	HNI	C4C-C3C-CAC-CBC


There are no ring outliers.

4 monomers are involved in 42 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	142	HNI	7	0
3	D	147	HNI	20	0
3	C	142	HNI	13	0
3	B	147	HNI	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand HNI A 142




Bond lengths



Bond angles




Torsions




Rings


Ligand HNI D 147




Bond lengths



Bond angles





Torsions



Rings

Ligand HNI C 142





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.




6.4 Ligands ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around HNI A 142:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HNI D 147:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HNI C 142:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HNI B 147:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.