



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 24, 2021 – 06:04 pm BST

PDB ID : 6FP8  
Title : mTFP1/DARPin 1238\_E11 complex in space group C2  
Authors : Jakob, R.P.; Vigano, M.A.; Bieli, D.; Matsuda, S.; Schaefer, J.V.; Pluckthun, A.; Affolter, M.; Maier, T.  
Deposited on : 2018-02-09  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	<div> <div>2%</div> <div> <div></div> <div>80%</div> <div>7%</div> <div>12%</div> </div> </div>
2	B	182	<div> <div>8%</div> <div> <div></div> <div>87%</div> <div>•</div> <div>8%</div> </div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6561 atoms, of which 3050 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GFP-like fluorescent chromoprotein cFP484.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	217	Total	C	H	N	O	S	31	2	0
			3498	1133	1728	297	334	6			

There are 61 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	MET	-	initiating methionine	UNP Q9U6Y3
A	-21	GLY	-	expression tag	UNP Q9U6Y3
A	-20	SER	-	expression tag	UNP Q9U6Y3
A	-19	SER	-	expression tag	UNP Q9U6Y3
A	-18	HIS	-	expression tag	UNP Q9U6Y3
A	-17	HIS	-	expression tag	UNP Q9U6Y3
A	-16	HIS	-	expression tag	UNP Q9U6Y3
A	-15	HIS	-	expression tag	UNP Q9U6Y3
A	-14	HIS	-	expression tag	UNP Q9U6Y3
A	-13	HIS	-	expression tag	UNP Q9U6Y3
A	-12	SER	-	expression tag	UNP Q9U6Y3
A	-11	GLN	-	expression tag	UNP Q9U6Y3
A	-10	ASP	-	expression tag	UNP Q9U6Y3
A	-9	PRO	-	expression tag	UNP Q9U6Y3
A	-8	MET	-	expression tag	UNP Q9U6Y3
A	-7	VAL	-	expression tag	UNP Q9U6Y3
A	-6	SER	-	expression tag	UNP Q9U6Y3
A	-5	LYS	-	expression tag	UNP Q9U6Y3
A	-4	GLY	-	expression tag	UNP Q9U6Y3
A	-3	GLU	-	expression tag	UNP Q9U6Y3
A	-2	GLU	-	expression tag	UNP Q9U6Y3
A	38	ASN	HIS	conflict	UNP Q9U6Y3
A	40	ILE	LEU	conflict	UNP Q9U6Y3
A	58	THR	SER	conflict	UNP Q9U6Y3
A	59	THR	ASN	conflict	UNP Q9U6Y3
A	62	PIA	GLN	conflict	UNP Q9U6Y3
A	?	-	TYR	deletion	UNP Q9U6Y3

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	GLY	deletion	UNP Q9U6Y3
A	68	PHE	LEU	conflict	UNP Q9U6Y3
A	76	PRO	ALA	conflict	UNP Q9U6Y3
A	77	ASN	ASP	conflict	UNP Q9U6Y3
A	119	HIS	ARG	conflict	UNP Q9U6Y3
A	120	LEU	PHE	conflict	UNP Q9U6Y3
A	121	LYS	ASP	conflict	UNP Q9U6Y3
A	123	GLU	MET	conflict	UNP Q9U6Y3
A	137	THR	LEU	conflict	UNP Q9U6Y3
A	138	GLY	LYS	conflict	UNP Q9U6Y3
A	140	ASP	GLU	conflict	UNP Q9U6Y3
A	141	ALA	PRO	conflict	UNP Q9U6Y3
A	145	ARG	ILE	conflict	UNP Q9U6Y3
A	154	LYS	VAL	conflict	UNP Q9U6Y3
A	157	VAL	ILE	conflict	UNP Q9U6Y3
A	158	LYS	SER	conflict	UNP Q9U6Y3
A	160	LYS	SER	conflict	UNP Q9U6Y3
A	169	HIS	TYR	conflict	UNP Q9U6Y3
A	171	VAL	CYS	conflict	UNP Q9U6Y3
A	175	THR	SER	conflict	UNP Q9U6Y3
A	178	ARG	LYS	conflict	UNP Q9U6Y3
A	182	ALA	VAL	conflict	UNP Q9U6Y3
A	209	VAL	LEU	conflict	UNP Q9U6Y3
A	212	SER	ASN	conflict	UNP Q9U6Y3
A	217	ASN	TYR	conflict	UNP Q9U6Y3
A	219	THR	-	expression tag	UNP Q9U6Y3
A	220	ASP	-	expression tag	UNP Q9U6Y3
A	221	GLY	-	expression tag	UNP Q9U6Y3
A	222	MET	-	expression tag	UNP Q9U6Y3
A	223	ASP	-	expression tag	UNP Q9U6Y3
A	224	GLU	-	expression tag	UNP Q9U6Y3
A	225	LEU	-	expression tag	UNP Q9U6Y3
A	226	TYR	-	expression tag	UNP Q9U6Y3
A	227	LYS	-	expression tag	UNP Q9U6Y3

- Molecule 2 is a protein called DARPin1238\_E11.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	167	Total	C	H	N	O	S	11	2	0
			2547	809	1268	224	242	4			

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula:

C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	1	0
			22	6	14	2		
3	A	1	Total	C	H	O	1	0
			22	6	14	2		
3	A	1	Total	C	H	O	1	0
			22	6	14	2		

- Molecule 4 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C H O 10 2 6 2	0	0
5	A	1	Total C H O 10 2 6 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	263	Total O 263 263	0	0
6	B	147	Total O 147 147	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GFP-like fluorescent chromoprotein cFP484



- Molecule 2: DARPin1238\_E11



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	135.95Å 75.01Å 48.30Å 90.00° 90.73° 90.00°	Depositor
Resolution (Å)	39.61 – 1.85 39.61 – 1.85	Depositor EDS
% Data completeness (in resolution range)	96.6 (39.61-1.85) 97.2 (39.61-1.85)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.24 (at 1.85Å)	Xtriage
Refinement program	PHENIX (1.10pre_2131: ???)	Depositor
R, $R_{free}$	0.148 , 0.176 0.151 , 0.179	Depositor DCC
$R_{free}$ test set	1996 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.8	Xtriage
Anisotropy	0.231	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 66.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.008 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.008 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.014 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.015 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.025 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	6561	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, PIA, MPD, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.66	2/1799 (0.1%)	0.76	1/2425 (0.0%)
2	B	0.42	0/1309	0.64	0/1776
All	All	0.57	2/3108 (0.1%)	0.71	1/4201 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	65	ASN	C-N	-5.70	1.21	1.34
1	A	103	VAL	CB-CG2	-5.37	1.41	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	140	ASP	CB-CG-OD2	-6.52	112.43	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1770	1728	1728	8	1
2	B	1279	1268	1270	5	0
3	A	24	42	42	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	15	0	0	0	0
4	B	5	0	0	0	0
5	A	8	12	12	0	0
6	A	263	0	0	3	1
6	B	147	0	0	2	2
All	All	3511	3050	3052	14	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:ASP:OD1	6:A:401:HOH:O	2.03	0.75
1:A:142:SER:HB3	1:A:159:HIS:CE1	2.39	0.57
1:A:3:VAL:HB	1:A:109:MET:HE1	1.88	0.56
1:A:190:PHE:CE2	1:A:218:SER:HA	2.46	0.51
2:B:36[B]:MET:HE3	6:B:344:HOH:O	2.11	0.50
1:A:110:GLU:HB3	1:A:115:ILE:HD11	1.94	0.47
2:B:119:LEU:HD11	2:B:131:VAL:HG13	1.98	0.46
2:B:36[A]:MET:CE	2:B:71:THR:HG21	2.47	0.45
3:A:303:MPD:HM2	6:A:540:HOH:O	2.16	0.44
1:A:211:GLU:HG2	1:A:212:SER:N	2.34	0.43
2:B:29:ASP:HA	2:B:64:ILE:HD11	2.02	0.42
2:B:130:ILE:HD13	6:B:318:HOH:O	2.18	0.41
1:A:120:LEU:C	1:A:120:LEU:HD23	2.41	0.41
1:A:174:LYS:HD2	6:A:620:HOH:O	2.21	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:ARG:NH2	1:A:196:GLU:OE2[2_556]	1.81	0.39
6:A:604:HOH:O	6:B:371:HOH:O[4_456]	1.93	0.27
6:B:306:HOH:O	6:B:427:HOH:O[4_445]	2.13	0.07

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/248 (86%)	212 (99%)	2 (1%)	0	100	100
2	B	167/182 (92%)	166 (99%)	1 (1%)	0	100	100
All	All	381/430 (89%)	378 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/217 (88%)	190 (100%)	1 (0%)	88	86
2	B	131/143 (92%)	130 (99%)	1 (1%)	81	76
All	All	322/360 (89%)	320 (99%)	2 (1%)	86	83

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	159	HIS
2	B	47	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	199	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	PIA	A	62	1	21,21,22	2.66	7 (33%)	27,29,31	2.01	6 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PIA	A	62	1	-	3/8/27/28	0/2/2/2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	62	PIA	CB2-CA2	9.37	1.42	1.35
1	A	62	PIA	C2-N3	-3.58	1.31	1.39
1	A	62	PIA	CA1-C1	-3.12	1.47	1.51
1	A	62	PIA	CA2-C2	-2.76	1.45	1.48
1	A	62	PIA	O2-C2	2.74	1.28	1.23
1	A	62	PIA	CA3-N3	-2.52	1.42	1.47
1	A	62	PIA	CE1-CD1	-2.11	1.34	1.38

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	62	PIA	CA2-C2-N3	5.45	105.95	103.37
1	A	62	PIA	O2-C2-CA2	-4.91	128.20	130.96
1	A	62	PIA	CA1-C1-N3	-3.31	120.77	124.75
1	A	62	PIA	O3-C3-CA3	2.35	133.49	126.39
1	A	62	PIA	CA3-N3-C1	-2.15	124.58	127.16
1	A	62	PIA	CA2-N2-C1	-2.02	104.28	105.77

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	62	PIA	C3-CA3-N3-C2
1	A	62	PIA	N2-C1-CA1-CB1
1	A	62	PIA	C3-CA3-N3-C1

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	MPD	A	301	-	7,7,7	0.29	0	9,10,10	0.47	0
3	MPD	A	302	-	7,7,7	0.55	0	9,10,10	0.93	0
4	SO4	B	201	-	4,4,4	0.82	0	6,6,6	1.01	0
5	EDO	A	307	-	3,3,3	0.97	0	2,2,2	0.78	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	EDO	A	308	-	3,3,3	0.33	0	2,2,2	1.40	0
4	SO4	A	306	-	4,4,4	0.83	0	6,6,6	0.67	0
3	MPD	A	303	-	7,7,7	0.36	0	9,10,10	0.58	0
4	SO4	A	304	-	4,4,4	0.81	0	6,6,6	0.94	0
4	SO4	A	305	-	4,4,4	0.89	0	6,6,6	1.35	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	A	301	-	-	0/5/5/5	-
3	MPD	A	302	-	-	0/5/5/5	-
5	EDO	A	308	-	-	1/1/1/1	-
3	MPD	A	303	-	-	0/5/5/5	-
5	EDO	A	307	-	-	1/1/1/1	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	305	SO4	O3-S-O2	-2.21	97.79	109.31

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	308	EDO	O1-C1-C2-O2
5	A	307	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	303	MPD	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	216/248 (87%)	-0.15	4 (1%) 66 66	19, 27, 49, 87	0
2	B	167/182 (91%)	0.25	15 (8%) 9 9	24, 37, 75, 109	0
All	All	383/430 (89%)	0.02	19 (4%) 28 27	19, 31, 68, 109	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	37	ALA	6.4
2	B	36[A]	MET	4.3
2	B	39	GLY	4.1
2	B	38	ASN	4.0
2	B	178	ASP	4.0
1	A	219	THR	3.8
2	B	35	LEU	3.5
1	A	220	ASP	3.1
2	B	13	GLY	3.1
2	B	156	ILE	3.1
2	B	12	HIS	2.9
2	B	161	GLU	2.8
2	B	34	ILE	2.8
1	A	165	GLY	2.8
2	B	16	LEU	2.4
1	A	2	GLY	2.3
2	B	162	ASP	2.2
2	B	31	GLU	2.1
2	B	42	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	PIA	A	62	20/21	0.94	0.15	2,23,27,28	1

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	MPD	A	303	8/8	0.72	0.31	51,74,89,92	1
4	SO4	A	305	5/5	0.75	0.28	73,76,79,84	0
4	SO4	B	201	5/5	0.76	0.17	69,73,77,89	0
5	EDO	A	308	4/4	0.77	0.18	45,64,72,77	0
5	EDO	A	307	4/4	0.84	0.16	28,57,69,69	0
4	SO4	A	306	5/5	0.92	0.32	70,71,79,89	0
3	MPD	A	301	8/8	0.93	0.12	20,38,50,50	1
4	SO4	A	304	5/5	0.95	0.20	69,70,79,84	0
3	MPD	A	302	8/8	0.95	0.12	25,40,62,62	1

### 6.5 Other polymers [i](#)

There are no such residues in this entry.