



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 7, 2020 – 02:49 AM BST

PDB ID : 5FSG
Title : Structure of the hantavirus nucleoprotein provides insights into the mechanism of RNA encapsidation and a template for drug design
Authors : Olal, D.; Daumke, O.
Deposited on : 2016-01-05
Resolution : 3.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	698	
2	B	4	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 10176 atoms, of which 5028 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MALTOSE-BINDING PERIPLASMIC PROTEIN, HANTAVIRUS NUCLEOPROTEIN.

Mol	Chain	Residues	Atoms								ZeroOcc	AltConf	Trace
1	A	661	Total	C	H	N	O	S	Se		0	0	0
			10089	3269	4986	854	958	5	17				

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	ALA	-	linker	UNP P05133
A	-12	ALA	-	linker	UNP P05133
A	-11	LEU	-	linker	UNP P05133
A	-10	ALA	-	linker	UNP P05133
A	-9	ALA	-	linker	UNP P05133
A	-8	GLY	-	linker	UNP P05133
A	-7	SER	-	linker	UNP P05133
A	-6	ALA	-	linker	UNP P05133
A	-5	GLN	-	linker	UNP P05133
A	-4	THR	-	linker	UNP P05133
A	-3	ASN	-	linker	UNP P05133
A	-2	ALA	-	linker	UNP P05133
A	-1	ALA	-	linker	UNP P05133
A	0	ALA	-	linker	UNP P05133
A	430	LEU	-	expression tag	UNP P05133
A	431	GLU	-	expression tag	UNP P05133
A	432	HIS	-	expression tag	UNP P05133
A	433	HIS	-	expression tag	UNP P05133
A	434	HIS	-	expression tag	UNP P05133
A	435	HIS	-	expression tag	UNP P05133
A	436	HIS	-	expression tag	UNP P05133
A	437	HIS	-	expression tag	UNP P05133
A	185	ASN	GLN	conflict	UNP P05133
A	377	ASP	GLY	conflict	UNP P05133

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-

(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	H	O			
2	B	4	87	24	42	21	0	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MALTOSE-BINDING PERIPLASMIC PROTEIN, HANTAVIRUS NUCLEOPROTEIN



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.63 Å 86.84 Å 131.50 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.80 – 3.21 47.81 – 3.21	Depositor EDS
% Data completeness (in resolution range)	99.6 (47.80-3.21) 99.8 (47.81-3.21)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.45 (at 3.19 Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.290 , 0.299 0.292 , 0.302	Depositor DCC
R_{free} test set	679 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	82.5	Xtriage
Anisotropy	0.642	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 55.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	10176	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/5197	0.43	0/7021

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5103	4986	5082	51	0
2	B	45	42	39	1	0
All	All	5148	5028	5121	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:-37:GLN:N	1:A:-37:GLN:OE1	2.22	0.72
1:A:139:TYR:O	1:A:143:THR:HG23	1.90	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:-182:ALA:O	1:A:-179:THR:OG1	2.08	0.70
1:A:-368:GLU:HG2	1:A:-368:GLU:O	1.92	0.69
1:A:327:GLU:OE2	1:A:377:ASP:N	2.27	0.68
1:A:145:GLY:O	1:A:146:ARG:HB2	1.96	0.65
1:A:261:THR:OG1	1:A:264:ASP:OD2	2.12	0.64
1:A:263:ARG:NH2	1:A:298:ASP:OD2	2.31	0.62
1:A:298:ASP:OD1	1:A:299:ILE:N	2.34	0.60
1:A:-328:GLU:OE1	2:B:3:GLC:O3	2.11	0.59
1:A:423:ASN:O	1:A:424:GLN:NE2	2.38	0.57
1:A:314:ARG:HG2	1:A:368:ARG:HB3	1.89	0.55
1:A:-3:ASN:HA	1:A:0:ALA:HB3	1.89	0.54
1:A:327:GLU:OE1	1:A:327:GLU:N	2.41	0.53
1:A:385:MSE:O	1:A:389:GLY:N	2.42	0.52
1:A:370:GLN:N	1:A:370:GLN:OE1	2.44	0.51
1:A:211:LYS:O	1:A:214:GLN:NE2	2.44	0.50
1:A:415:ASP:O	1:A:419:LYS:N	2.42	0.50
1:A:-7:SER:O	1:A:-3:ASN:ND2	2.41	0.48
1:A:-319:THR:O	1:A:-319:THR:HG22	2.14	0.47
1:A:-344:GLU:OE1	1:A:-339:ILE:N	2.48	0.47
1:A:231:ASP:OD2	1:A:235:ARG:NH2	2.48	0.47
1:A:-179:THR:O	1:A:-176:VAL:HG22	2.15	0.46
1:A:235:ARG:NH1	1:A:391:GLU:OE1	2.48	0.46
1:A:-345:PHE:O	1:A:-341:THR:HG22	2.14	0.46
1:A:-232:LYS:NZ	1:A:-170:LYS:O	2.47	0.46
1:A:-221:LEU:HD12	1:A:-164:THR:OG1	2.16	0.46
1:A:-290:ASP:N	1:A:-290:ASP:OD1	2.49	0.46
1:A:-160:ILE:HG13	1:A:-159:ALA:N	2.31	0.45
1:A:-366:LYS:HZ1	1:A:-336:THR:CB	2.30	0.45
1:A:-295:ALA:HB3	1:A:-106:ILE:HG23	1.98	0.45
1:A:-240:ILE:N	1:A:-239:PRO:HD2	2.31	0.45
1:A:258:GLY:N	1:A:259:PRO:CD	2.80	0.44
1:A:307:VAL:CG1	1:A:316:PRO:HG3	2.47	0.44
1:A:328:LEU:O	1:A:331:PHE:HB3	2.18	0.44
1:A:-369:GLU:O	1:A:-367:GLY:N	2.50	0.43
1:A:-122:PHE:O	1:A:-121:LYS:HG2	2.18	0.43
1:A:343:MSE:HE1	1:A:398:LEU:O	2.19	0.42
1:A:422:SER:O	1:A:424:GLN:N	2.52	0.42
1:A:-122:PHE:O	1:A:-121:LYS:CG	2.67	0.42
1:A:-214:TRP:N	1:A:-213:PRO:HD2	2.34	0.42
1:A:-327:GLU:O	1:A:-324:PRO:HD2	2.19	0.42
1:A:-164:THR:HA	1:A:-160:ILE:HD11	2.01	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:-307:ASP:OD1	1:A:-307:ASP:O	2.38	0.41
1:A:-307:ASP:OD1	1:A:-36:MSE:HB2	2.21	0.41
1:A:-307:ASP:OD1	1:A:-36:MSE:CB	2.68	0.41
1:A:137:ALA:O	1:A:141:LEU:HD13	2.21	0.41
1:A:-142:TRP:O	1:A:-75:LYS:NZ	2.48	0.41
1:A:-314:ASP:O	1:A:-106:ILE:HD12	2.21	0.41
1:A:295:MSE:HE3	1:A:318:THR:O	2.20	0.41
1:A:-127:THR:HG22	1:A:-126:VAL:H	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	651/698 (93%)	628 (96%)	19 (3%)	4 (1%)	25 64

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	-368	GLU
1	A	377	ASP
1	A	423	ASN
1	A	207	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	530/556 (95%)	512 (97%)	18 (3%)	37 70

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-370	ILE
1	A	-363	ILE
1	A	-328	GLU
1	A	-307	ASP
1	A	-290	ASP
1	A	-114	PHE
1	A	-42	MSE
1	A	-14	ASP
1	A	140	MSE
1	A	226	LEU
1	A	239	TRP
1	A	263	ARG
1	A	265	TYR
1	A	280	GLU
1	A	327	GLU
1	A	328	LEU
1	A	385	MSE
1	A	429	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	B	1	2	12,12,12	2.11	3 (25%)	17,17,17	0.91	0
2	GLC	B	2	2	11,11,12	2.76	6 (54%)	15,15,17	1.09	1 (6%)
2	GLC	B	3	2	11,11,12	2.67	4 (36%)	15,15,17	1.08	0
2	GLC	B	4	2	11,11,12	2.71	4 (36%)	15,15,17	1.15	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0/2/22/22	0/1/1/1
2	GLC	B	2	2	-	0/2/19/22	0/1/1/1
2	GLC	B	3	2	-	0/2/19/22	0/1/1/1
2	GLC	B	4	2	-	0/2/19/22	0/1/1/1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	4	GLC	O5-C5	6.23	1.56	1.43
2	B	3	GLC	O5-C5	6.21	1.56	1.43
2	B	2	GLC	O5-C5	6.19	1.56	1.43
2	B	1	GLC	O5-C5	4.59	1.55	1.44
2	B	2	GLC	C2-C3	-3.70	1.47	1.52
2	B	3	GLC	C6-C5	-3.47	1.40	1.51
2	B	4	GLC	C2-C3	-3.38	1.47	1.52
2	B	4	GLC	O5-C1	3.29	1.49	1.43
2	B	2	GLC	C6-C5	-3.15	1.41	1.51
2	B	3	GLC	C2-C3	-3.00	1.48	1.52
2	B	4	GLC	C6-C5	-2.96	1.41	1.51
2	B	1	GLC	C6-C5	-2.90	1.42	1.51
2	B	1	GLC	O2-C2	2.70	1.49	1.43
2	B	2	GLC	O3-C3	2.54	1.49	1.43
2	B	2	GLC	O5-C1	2.15	1.47	1.43
2	B	3	GLC	O5-C1	2.14	1.47	1.43
2	B	2	GLC	O2-C2	2.04	1.47	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	GLC	C3-C4-C5	2.16	114.09	110.24
2	B	4	GLC	C1-O5-C5	2.14	115.09	112.19
2	B	4	GLC	C1-C2-C3	2.14	112.30	109.67

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	3	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	643/698 (92%)	0.86	64 (9%) 7 4	50, 80, 107, 129	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	254	SER	5.6
1	A	-90	ASN	4.8
1	A	-371	LYS	4.5
1	A	246	LEU	4.2
1	A	-366	LYS	4.2
1	A	-199	ASN	4.1
1	A	398	LEU	4.0
1	A	248	PRO	4.0
1	A	-178	PHE	3.9
1	A	-88	LEU	3.8
1	A	-91	GLU	3.7
1	A	287	HIS	3.6
1	A	247	LEU	3.6
1	A	-272	ASN	3.4
1	A	-201	TYR	3.2
1	A	172	ARG	3.2
1	A	-106	ILE	3.2
1	A	-284	LYS	3.2
1	A	-276	ALA	3.2
1	A	177	LEU	3.1
1	A	-295	ALA	3.1
1	A	422	SER	3.0
1	A	136	LYS	2.8
1	A	293	CYS	2.8
1	A	-317	ASP	2.8
1	A	-94	GLU	2.7
1	A	-300	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	-323	GLN	2.7
1	A	185	ASN	2.6
1	A	399	GLY	2.6
1	A	369	THR	2.6
1	A	174	PRO	2.6
1	A	244	CYS	2.5
1	A	-208	ASP	2.5
1	A	159	PHE	2.5
1	A	-160	ILE	2.5
1	A	324	GLY	2.5
1	A	239	TRP	2.5
1	A	-234	GLU	2.4
1	A	-294	GLU	2.4
1	A	-165	ASP	2.4
1	A	250	THR	2.4
1	A	179	VAL	2.4
1	A	183	ASN	2.4
1	A	330	ALA	2.4
1	A	284	ILE	2.3
1	A	-305	PHE	2.3
1	A	-179	THR	2.3
1	A	-239	PRO	2.3
1	A	-343	LYS	2.2
1	A	-282	TYR	2.2
1	A	275	ASN	2.2
1	A	296	ILE	2.2
1	A	-369	GLU	2.2
1	A	-278	TRP	2.1
1	A	-184	ALA	2.1
1	A	326	ALA	2.1
1	A	362	TYR	2.1
1	A	155	THR	2.1
1	A	158	ARG	2.0
1	A	-221	LEU	2.0
1	A	-66	SER	2.0
1	A	-255	TYR	2.0
1	A	-158	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	B	4	11/12	0.86	0.41	54,61,64,65	0
2	GLC	B	3	11/12	0.91	0.28	31,38,46,48	0
2	GLC	B	1	12/12	0.92	0.27	24,26,31,32	0
2	GLC	B	2	11/12	0.93	0.23	19,21,25,26	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.