



wwPDB X-ray Structure Validation Summary Report i

May 13, 2020 – 01:20 am BST

PDB ID : 2FT1
Title : Bacteriophage HK97 Head II
Authors : Gan, L.; Speir, J.A.; Conway, J.F.; Lander, G.; Cheng, N.; Firek, B.A.; Hendrix, R.W.; Duda, R.L.; Liljas, L.; Johnson, J.E.
Deposited on : 2006-01-23
Resolution : 3.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see references ①) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1004 (4.12-3.68)
Ramachandran outliers	138981	1021 (4.14-3.66)
Sidechain outliers	138945	1014 (4.14-3.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%



2 Entry composition [\(i\)](#)

There is only 1 type of molecule in this entry. The entry contains 14777 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called major capsid protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	280	Total	C	N	O	S	0	0	0
			2151	1344	375	422	10			
1	B	280	Total	C	N	O	S	0	0	0
			2151	1344	375	422	10			
1	C	280	Total	C	N	O	S	0	0	0
			2151	1344	375	422	10			
1	D	280	Total	C	N	O	S	0	0	0
			2151	1344	375	422	10			
1	E	280	Total	C	N	O	S	0	0	0
			2151	1344	375	422	10			
1	F	258	Total	C	N	O	S	0	0	0
			2001	1251	351	390	9			
1	G	261	Total	C	N	O	S	0	0	0
			2021	1265	354	393	9			


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: major capsid protein



- Molecule 1: major capsid protein




- Molecule 1: major capsid protein




- Molecule 1: major capsid protein

Chain D:




- Molecule 1: major capsid protein

Chain E:



- Molecule 1: major capsid protein

Chain F:



- Molecule 1: major capsid protein

Chain G:



4 Data and refinement statistics i

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	1009.80Å 1009.80Å 732.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.90 30.00 – 3.90	Depositor EDS
% Data completeness (in resolution range)	52.1 (30.00-3.90) 52.1 (30.00-3.90)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	1.47 (at 3.86Å)	Xtriage
Refinement program	CNS	Depositor
R , R_{free}	0.297 , (Not available) 0.279 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	48.5	Xtriage
Anisotropy	0.201	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.24 , 36.5	EDS
L-test for twinning ²	$< L > = 0.40$, $< L^2 > = 0.22$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.09	EDS
Total number of atoms	14777	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.26	0/2188	0.52	0/2969
1	B	0.25	0/2188	0.51	0/2969
1	C	0.26	0/2188	0.52	0/2969
1	D	0.27	0/2188	0.51	0/2969
1	E	0.25	0/2188	0.53	0/2969
1	F	0.26	0/2036	0.52	0/2762
1	G	0.28	0/2056	0.51	0/2789
All	All	0.26	0/15032	0.52	0/20396

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2151	0	2119	62	0
1	B	2151	0	2119	47	0
1	C	2151	0	2119	61	0
1	D	2151	0	2119	44	0
1	E	2151	0	2119	51	0
1	F	2001	0	1967	61	0
1	G	2021	0	1992	34	0
All	All	14777	0	14554	320	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 320 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:229:THR:HG23	1:D:230:GLY:H	1.35	0.92
1:E:210:ARG:HH21	1:F:156:PHE:HD2	1.20	0.88
1:F:227:ASP:HB3	1:F:229:THR:HG22	1.54	0.88
1:E:116:ILE:HG21	1:F:146:ASN:HB2	1.59	0.82
1:D:193:SER:HB3	1:D:196:VAL:HG23	1.64	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	278/282 (99%)	242 (87%)	33 (12%)	3 (1%)	14 51
1	B	278/282 (99%)	243 (87%)	31 (11%)	4 (1%)	11 46
1	C	278/282 (99%)	246 (88%)	26 (9%)	6 (2%)	6 38
1	D	278/282 (99%)	248 (89%)	25 (9%)	5 (2%)	8 42
1	E	278/282 (99%)	252 (91%)	21 (8%)	5 (2%)	8 42
1	F	256/282 (91%)	229 (90%)	22 (9%)	5 (2%)	7 40
1	G	259/282 (92%)	230 (89%)	29 (11%)	0	100 100
All	All	1905/1974 (96%)	1690 (89%)	187 (10%)	28 (2%)	10 45

5 of 28 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	105	LEU
1	C	134	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	304	THR
1	D	229	THR
1	E	134	ILE

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	230/231 (100%)	214 (93%)	16 (7%)	15 44
1	B	230/231 (100%)	218 (95%)	12 (5%)	23 53
1	C	230/231 (100%)	220 (96%)	10 (4%)	29 57
1	D	230/231 (100%)	220 (96%)	10 (4%)	29 57
1	E	230/231 (100%)	220 (96%)	10 (4%)	29 57
1	F	213/231 (92%)	202 (95%)	11 (5%)	23 53
1	G	215/231 (93%)	208 (97%)	7 (3%)	38 63
All	All	1578/1617 (98%)	1502 (95%)	76 (5%)	25 54

5 of 76 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	277	LEU
1	D	202	MET
1	G	129	LEU
1	C	300	PRO
1	D	115	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	352	ASN
1	E	237	ASN
1	G	232	ASN
1	E	191	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	208	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.