



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 7, 2020 – 01:43 am BST

PDB ID : 6FTD
Title : Deinococcus radiodurans BphP PAS-GAF H290T mutant
Authors : Edlund, P.; Takala, H.; Westenhoff, S.; Ihalainen, J.A.
Deposited on : 2018-02-21
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	<div> <div>6%</div> <div> <div></div> <div>78%</div> <div>12%</div> <div>8%</div> </div> </div>
1	B	343	<div> <div>6%</div> <div> <div></div> <div>76%</div> <div>13%</div> <div>8%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	B	403	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5741 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bacteriophytochrome.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	314	Total	C	N	O	S	0	19	0
			2526	1611	442	463	10			
1	B	314	Total	C	N	O	S	0	17	0
			2516	1608	436	461	11			

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	initiating methionine	UNP Q9RZA4
A	-12	ALA	-	expression tag	UNP Q9RZA4
A	-11	SER	-	expression tag	UNP Q9RZA4
A	-10	MET	-	expression tag	UNP Q9RZA4
A	-9	THR	-	expression tag	UNP Q9RZA4
A	-8	GLY	-	expression tag	UNP Q9RZA4
A	-7	GLY	-	expression tag	UNP Q9RZA4
A	-6	GLN	-	expression tag	UNP Q9RZA4
A	-5	GLN	-	expression tag	UNP Q9RZA4
A	-4	MET	-	expression tag	UNP Q9RZA4
A	-3	GLY	-	expression tag	UNP Q9RZA4
A	-2	ARG	-	expression tag	UNP Q9RZA4
A	-1	GLY	-	expression tag	UNP Q9RZA4
A	0	SER	-	expression tag	UNP Q9RZA4
A	290	THR	HIS	engineered mutation	UNP Q9RZA4
A	322	LEU	-	expression tag	UNP Q9RZA4
A	323	GLU	-	expression tag	UNP Q9RZA4
A	324	HIS	-	expression tag	UNP Q9RZA4
A	325	HIS	-	expression tag	UNP Q9RZA4
A	326	HIS	-	expression tag	UNP Q9RZA4
A	327	HIS	-	expression tag	UNP Q9RZA4
A	328	HIS	-	expression tag	UNP Q9RZA4
A	329	HIS	-	expression tag	UNP Q9RZA4
B	-13	MET	-	initiating methionine	UNP Q9RZA4
B	-12	ALA	-	expression tag	UNP Q9RZA4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-11	SER	-	expression tag	UNP Q9RZA4
B	-10	MET	-	expression tag	UNP Q9RZA4
B	-9	THR	-	expression tag	UNP Q9RZA4
B	-8	GLY	-	expression tag	UNP Q9RZA4
B	-7	GLY	-	expression tag	UNP Q9RZA4
B	-6	GLN	-	expression tag	UNP Q9RZA4
B	-5	GLN	-	expression tag	UNP Q9RZA4
B	-4	MET	-	expression tag	UNP Q9RZA4
B	-3	GLY	-	expression tag	UNP Q9RZA4
B	-2	ARG	-	expression tag	UNP Q9RZA4
B	-1	GLY	-	expression tag	UNP Q9RZA4
B	0	SER	-	expression tag	UNP Q9RZA4
B	290	THR	HIS	engineered mutation	UNP Q9RZA4
B	322	LEU	-	expression tag	UNP Q9RZA4
B	323	GLU	-	expression tag	UNP Q9RZA4
B	324	HIS	-	expression tag	UNP Q9RZA4
B	325	HIS	-	expression tag	UNP Q9RZA4
B	326	HIS	-	expression tag	UNP Q9RZA4
B	327	HIS	-	expression tag	UNP Q9RZA4
B	328	HIS	-	expression tag	UNP Q9RZA4
B	329	HIS	-	expression tag	UNP Q9RZA4

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	6	2		
3	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 4 is 3-[2-[(Z)-[3-(2-carboxyethyl)-5-[(Z)-(4-ethenyl-3-methyl-5-oxidanylidene-pyrro-2-ylidene)methyl]-4-methyl-pyrrol-1-ium-2-ylidene]methyl]-5-[(Z)-[(3E)-3-ethylidene-4-methyl-5-oxidanylidene-pyrrolidin-2-ylidene]methyl]-4-methyl-1H-pyrrol-3-yl]propanoic acid (three-letter code: LBV) (formula: $C_{33}H_{37}N_4O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			43	33	4	6		
4	B	1	Total	C	N	O	0	0
			43	33	4	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	274	Total	O	0	0
			274	274		
5	B	299	Total	O	0	0
			299	299		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Bacteriophytochrome



• Molecule 1: Bacteriophytochrome



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	94.81Å 53.03Å 135.88Å 90.00° 93.30° 90.00°	Depositor
Resolution (Å)	47.33 – 1.40 47.33 – 1.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.33-1.40) 99.7 (47.33-1.40)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.148 , 0.198 0.157 , 0.205	Depositor DCC
R_{free} test set	6620 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	16.6	Xtriage
Anisotropy	0.184	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 51.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	5741	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.73 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.9659e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, ACT, LBV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.36	11/2603 (0.4%)	1.40	23/3560 (0.6%)
1	B	1.36	8/2594 (0.3%)	1.38	28/3548 (0.8%)
All	All	1.36	19/5197 (0.4%)	1.39	51/7108 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	24	CYS	CB-SG	-8.25	1.68	1.82
1	B	132	TRP	CE3-CZ3	6.43	1.49	1.38
1	A	295	TYR	CD2-CE2	-6.34	1.29	1.39
1	B	13	TYR	CG-CD1	-6.08	1.31	1.39
1	B	120[A]	GLU	CD-OE1	6.04	1.32	1.25
1	B	120[B]	GLU	CD-OE1	6.04	1.32	1.25
1	A	27	GLU	CD-OE1	-5.87	1.19	1.25
1	A	27	GLU	CD-OE2	-5.87	1.19	1.25
1	A	127	GLU	CG-CD	5.85	1.60	1.51
1	B	193	GLU	CD-OE1	5.49	1.31	1.25
1	A	276	SER	CB-OG	5.47	1.49	1.42
1	A	132	TRP	CB-CG	5.36	1.59	1.50
1	B	25	GLU	CD-OE1	5.33	1.31	1.25
1	A	133	ASP	C-O	-5.30	1.13	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	9	PHE	CB-CG	-5.27	1.42	1.51
1	B	254	ARG	CZ-NH2	-5.16	1.26	1.33
1	A	127	GLU	CB-CG	-5.13	1.42	1.52
1	A	65	GLU	CD-OE2	5.09	1.31	1.25
1	B	320	GLU	CD-OE1	5.04	1.31	1.25

All (51) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	213	ARG	NE-CZ-NH1	19.10	129.85	120.30
1	A	302	ARG	NE-CZ-NH2	-13.54	113.53	120.30
1	A	228[A]	ARG	NE-CZ-NH2	-12.43	114.09	120.30
1	A	228[B]	ARG	NE-CZ-NH2	-12.43	114.09	120.30
1	B	213	ARG	NE-CZ-NH2	-11.02	114.79	120.30
1	A	213	ARG	NE-CZ-NH1	10.63	125.62	120.30
1	A	254	ARG	NE-CZ-NH2	10.11	125.35	120.30
1	A	191	ARG	NE-CZ-NH2	-10.08	115.26	120.30
1	B	222	ARG	NE-CZ-NH2	-9.95	115.33	120.30
1	B	191	ARG	NE-CZ-NH2	-9.86	115.37	120.30
1	A	226	ASP	CB-CG-OD1	9.85	127.17	118.30
1	A	85	LEU	CA-CB-CG	9.60	137.38	115.30
1	B	302	ARG	NE-CZ-NH2	-8.63	115.99	120.30
1	A	24	CYS	CB-CA-C	-8.59	93.21	110.40
1	A	222	ARG	NE-CZ-NH2	-8.45	116.08	120.30
1	A	228[A]	ARG	NE-CZ-NH1	8.00	124.30	120.30
1	A	228[B]	ARG	NE-CZ-NH1	8.00	124.30	120.30
1	A	222	ARG	NE-CZ-NH1	7.89	124.25	120.30
1	A	302	ARG	NE-CZ-NH1	7.37	123.98	120.30
1	B	192	ARG	NE-CZ-NH2	-7.02	116.79	120.30
1	A	172	ARG	NE-CZ-NH1	7.00	123.80	120.30
1	B	192	ARG	NE-CZ-NH1	6.80	123.70	120.30
1	B	207	ASP	CB-CG-OD1	6.78	124.40	118.30
1	B	216	TYR	CG-CD2-CE2	6.73	126.69	121.30
1	B	254	ARG	NE-CZ-NH2	6.70	123.65	120.30
1	A	213	ARG	NE-CZ-NH2	-6.68	116.96	120.30
1	B	228	ARG	NE-CZ-NH1	-6.60	117.00	120.30
1	B	100	ARG	NE-CZ-NH2	-6.42	117.09	120.30
1	B	172	ARG	NE-CZ-NH2	-6.21	117.19	120.30
1	B	117	ARG	NE-CZ-NH1	-6.04	117.28	120.30
1	A	9	PHE	CB-CG-CD2	-5.99	116.61	120.80
1	B	308	LEU	CB-CG-CD2	5.92	121.06	111.00
1	A	235	ASP	CB-CG-OD1	-5.87	113.01	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	267	MET	CG-SD-CE	-5.70	91.08	100.20
1	B	228	ARG	NE-CZ-NH2	5.69	123.14	120.30
1	A	165	ARG	NE-CZ-NH1	5.67	123.14	120.30
1	A	154	ARG	NE-CZ-NH1	5.58	123.09	120.30
1	A	192	ARG	NE-CZ-NH2	-5.50	117.55	120.30
1	B	191	ARG	NE-CZ-NH1	5.47	123.04	120.30
1	B	154	ARG	NE-CZ-NH1	-5.37	117.61	120.30
1	B	62	LEU	CB-CG-CD2	5.33	120.07	111.00
1	B	13	TYR	CG-CD1-CE1	5.30	125.54	121.30
1	B	302	ARG	NE-CZ-NH1	5.18	122.89	120.30
1	B	265	ARG	NE-CZ-NH1	-5.17	117.72	120.30
1	B	319	LYS	CD-CE-NZ	5.15	123.55	111.70
1	B	216	TYR	CB-CG-CD1	5.13	124.08	121.00
1	B	300	ASP	CB-CG-OD1	5.10	122.89	118.30
1	B	127	GLU	OE1-CD-OE2	5.10	129.41	123.30
1	A	141	ARG	NE-CZ-NH1	5.08	122.84	120.30
1	B	226	ASP	CB-CG-OD2	-5.07	113.74	118.30
1	A	265	ARG	NE-CZ-NH1	-5.01	117.79	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	165	ARG	Sidechain
1	A	191	ARG	Sidechain
1	B	165	ARG	Sidechain
1	B	191	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2526	0	2545	20	0
1	B	2516	0	2531	29	0
2	A	12	0	9	1	0
2	B	12	0	9	2	0
3	A	8	0	14	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	8	0	14	0	0
4	A	43	0	33	2	0
4	B	43	0	34	0	0
5	A	274	0	0	7	0
5	B	299	0	0	8	0
All	All	5741	0	5189	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:162[A]:GLN:HE21	1:B:166[A]:GLU:HG3	1.32	0.94
1:B:54[B]:MET:HE1	1:B:69:LEU:HB2	1.67	0.76
1:A:167[B]:LEU:HD23	1:A:305[B]:LEU:HD21	1.71	0.71
1:A:167[B]:LEU:HD23	1:A:305[B]:LEU:CD2	2.21	0.71
1:B:293[A]:THR:HG23	1:B:294:PRO:HD2	1.74	0.70
1:A:162[B]:GLN:HG2	5:A:503:HOH:O	1.91	0.68
1:A:118:VAL:HG11	1:A:247[B]:PRO:CG	2.29	0.63
1:B:54[B]:MET:SD	1:B:66:PRO:HB3	2.40	0.62
1:B:162[B]:GLN:HG2	5:B:519:HOH:O	2.00	0.61
1:A:307:TYR:CE2	1:A:311[B]:LEU:HD11	2.36	0.61
1:B:54[B]:MET:HE1	1:B:69:LEU:CB	2.32	0.60
1:B:247[B]:PRO:HB3	5:B:772:HOH:O	2.02	0.59
1:A:158:GLU:O	1:A:162[B]:GLN:HG3	2.03	0.59
1:A:118:VAL:HG11	1:A:247[B]:PRO:HG3	1.84	0.58
1:B:162[B]:GLN:CG	5:B:519:HOH:O	2.54	0.56
1:A:228[B]:ARG:NH2	5:A:505:HOH:O	2.39	0.55
1:B:61:PHE:HD2	1:B:62:LEU:HD22	1.71	0.55
1:B:54[A]:MET:HE1	1:B:70:ARG:HG3	1.90	0.54
1:A:258[A]:PRO:HG2	5:A:683:HOH:O	2.08	0.54
1:B:54[B]:MET:SD	1:B:66:PRO:CB	2.96	0.53
1:B:159[A]:VAL:HG23	5:B:686:HOH:O	2.08	0.53
1:B:54[A]:MET:SD	1:B:70:ARG:HB2	2.49	0.53
1:B:258[A]:PRO:HG2	5:B:720:HOH:O	2.09	0.52
1:B:118:VAL:HG11	1:B:247[A]:PRO:CG	2.41	0.51
1:A:167[B]:LEU:HG	1:A:301:LEU:HD22	1.92	0.51
1:B:177[B]:LYS:HE2	1:B:284:TRP:CE2	2.46	0.51
1:A:259:MET:HG2	5:A:683:HOH:O	2.15	0.47
1:B:320:GLU:O	1:B:323:GLU:HB2	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:292[B]:GLN:HG3	2:A:403:ACT:H1	1.98	0.46
1:B:292[B]:GLN:CG	2:B:403:ACT:H1	2.46	0.46
1:B:162[A]:GLN:HE21	1:B:166[A]:GLU:CG	2.16	0.45
1:B:54[B]:MET:HB3	1:B:54[B]:MET:HE3	1.22	0.45
1:B:96:ALA:HA	5:B:748:HOH:O	2.16	0.45
1:B:292[A]:GLN:O	1:B:292[A]:GLN:HG2	2.16	0.45
1:B:154:ARG:HD2	5:B:736:HOH:O	2.16	0.45
1:B:259:MET:HG2	5:B:720:HOH:O	2.17	0.45
1:A:167[B]:LEU:HD23	1:A:305[B]:LEU:HD23	1.96	0.44
1:A:292[A]:GLN:HG2	1:A:292[A]:GLN:O	2.17	0.44
1:B:292[B]:GLN:HG3	2:B:403:ACT:H1	2.01	0.43
1:A:227:THR:O	1:A:261[A]:MET:HE3	2.18	0.42
1:A:90:PRO:HA	5:A:620:HOH:O	2.19	0.42
1:B:85:LEU:HD23	1:B:85:LEU:HA	1.86	0.42
1:A:306:GLU:HG3	5:A:729:HOH:O	2.18	0.42
1:B:59:ALA:HB2	1:B:66:PRO:HA	2.01	0.42
1:B:61:PHE:CD2	1:B:62:LEU:HD22	2.53	0.42
1:B:293[A]:THR:HG22	1:B:294:PRO:O	2.19	0.41
1:A:24:CYS:HB3	4:A:405:LBV:HAA1	1.94	0.41
1:A:319:LYS:HE2	5:A:525:HOH:O	2.20	0.41
4:A:405:LBV:N_D	4:A:405:LBV:HMC1	2.36	0.40
1:A:298:PRO:HA	1:A:299:PRO:HD3	1.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	329/343 (96%)	324 (98%)	5 (2%)	0	100	100
1	B	327/343 (95%)	324 (99%)	3 (1%)	0	100	100
All	All	656/686 (96%)	648 (99%)	8 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	272/275 (99%)	265 (97%)	7 (3%)	46	13
1	B	270/275 (98%)	262 (97%)	8 (3%)	41	10
All	All	542/550 (98%)	527 (97%)	15 (3%)	43	11

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	GLU
1	A	54	MET
1	A	85	LEU
1	A	140	LEU
1	A	148	GLU
1	A	222	ARG
1	A	319	LYS
1	B	54[A]	MET
1	B	54[B]	MET
1	B	64	GLN
1	B	95	ASP
1	B	110	HIS
1	B	130	GLU
1	B	222	ARG
1	B	319	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	86	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	A	402	-	1,3,3	2.27	1 (100%)	0,3,3	0.00	-
2	ACT	A	401	-	1,3,3	0.18	0	0,3,3	0.00	-
2	ACT	B	403	-	1,3,3	0.84	0	0,3,3	0.00	-
2	ACT	A	403	-	1,3,3	0.58	0	0,3,3	0.00	-
4	LBV	A	405	1	36,46,46	2.28	15 (41%)	41,67,67	1.86	12 (29%)
3	MPD	A	404	-	7,7,7	0.64	0	9,10,10	1.87	2 (22%)
4	LBV	B	405	1	36,46,46	2.64	11 (30%)	41,67,67	2.06	14 (34%)
3	MPD	B	404	-	7,7,7	0.72	0	9,10,10	1.24	1 (11%)
2	ACT	B	401	-	1,3,3	0.39	0	0,3,3	0.00	-
2	ACT	B	402	-	1,3,3	2.77	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	LBV	A	405	1	-	2/22/74/74	0/4/4/4
4	LBV	B	405	1	-	3/22/74/74	0/4/4/4
3	MPD	B	404	-	-	3/5/5/5	-
3	MPD	A	404	-	-	1/5/5/5	-

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	405	LBV	CHC-C1C	10.90	1.44	1.35
4	A	405	LBV	CHC-C1C	7.35	1.41	1.35
4	B	405	LBV	C2A-C3A	-5.44	1.44	1.51
4	B	405	LBV	C2C-C3C	4.83	1.47	1.36
4	A	405	LBV	C2A-C3A	-4.02	1.46	1.51
4	A	405	LBV	CAA-C3A	3.57	1.43	1.33
4	A	405	LBV	C4C-C3C	-3.47	1.38	1.45
4	B	405	LBV	C2A-C1A	-3.28	1.47	1.51
4	A	405	LBV	C4A-N_A	3.05	1.42	1.37
4	A	405	LBV	C1C-N_C	-2.83	1.32	1.38
2	B	402	ACT	CH3-C	-2.77	1.45	1.48
4	A	405	LBV	C1D-N_D	-2.76	1.33	1.37
4	B	405	LBV	O_D-C4D	2.74	1.28	1.23
4	A	405	LBV	C1C-C2C	-2.69	1.41	1.45
4	B	405	LBV	C1C-N_C	-2.49	1.33	1.38
4	B	405	LBV	CHD-C4C	2.48	1.46	1.40
4	B	405	LBV	CMB-C2B	2.47	1.56	1.51
4	B	405	LBV	CAA-C3A	2.28	1.39	1.33
2	A	402	ACT	CH3-C	-2.27	1.45	1.48
4	A	405	LBV	C3B-C2B	2.22	1.44	1.37
4	A	405	LBV	CAB-C3B	-2.21	1.48	1.52
4	B	405	LBV	O_A-C1A	2.19	1.27	1.23
4	A	405	LBV	C1D-C2D	-2.18	1.41	1.45
4	A	405	LBV	CHD-C4C	2.14	1.45	1.40
4	A	405	LBV	C2C-C3C	2.14	1.41	1.36
4	A	405	LBV	CHB-C4A	2.08	1.38	1.34
4	B	405	LBV	C4A-C3A	2.08	1.48	1.45
4	A	405	LBV	C2A-C1A	2.06	1.54	1.51

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	405	LBV	O_A-C1A-N_A	-4.50	119.49	124.94
4	B	405	LBV	C2D-C1D-N_D	4.12	113.02	106.99

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	405	LBV	CAA-C3A-C4A	-4.08	121.44	126.36
4	A	405	LBV	CMD-C2D-C1D	4.06	129.24	124.17
4	B	405	LBV	CMC-C3C-C4C	3.96	131.16	125.06
4	B	405	LBV	C3D-C2D-C1D	-3.89	103.33	108.03
4	A	405	LBV	C3D-C2D-C1D	-3.85	103.37	108.03
3	A	404	MPD	O2-C2-C1	3.75	120.12	108.08
4	A	405	LBV	C4C-C3C-C2C	3.70	110.75	106.51
4	B	405	LBV	C1D-N_D-C4D	-3.53	106.18	110.67
4	A	405	LBV	CAA-C3A-C4A	-3.09	122.63	126.36
4	A	405	LBV	CHD-C4C-C3C	3.02	132.66	124.90
3	A	404	MPD	CM-C2-C1	-2.98	104.36	110.57
4	A	405	LBV	CAC-C2C-C1C	2.86	130.07	125.01
4	A	405	LBV	C3C-C4C-N_C	-2.80	104.54	110.53
4	A	405	LBV	C2D-C1D-N_D	2.80	111.08	106.99
4	B	405	LBV	CHD-C1D-C2D	-2.74	121.51	126.95
4	B	405	LBV	C1C-C2C-C3C	-2.70	103.79	106.78
3	B	404	MPD	O2-C2-C1	-2.66	99.53	108.08
4	A	405	LBV	CHD-C1D-C2D	-2.59	121.82	126.95
4	A	405	LBV	CAC-CBC-CGC	2.59	117.01	112.67
4	B	405	LBV	CAC-CBC-CGC	2.56	116.96	112.67
4	A	405	LBV	C1C-N_C-C4C	2.52	111.25	106.51
4	A	405	LBV	C1C-C2C-C3C	-2.50	104.02	106.78
4	B	405	LBV	C1C-N_C-C4C	2.39	111.01	106.51
4	B	405	LBV	O_D-C4D-N_D	-2.26	119.82	125.08
4	B	405	LBV	CMB-C2B-C3B	-2.17	120.86	124.94
4	B	405	LBV	CMD-C2D-C1D	2.15	126.85	124.17
4	B	405	LBV	C3D-C4D-N_D	2.02	108.47	106.19

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	405	LBV	C3C-C4C-CHD-C1D
4	A	405	LBV	N_C-C4C-CHD-C1D
4	B	405	LBV	C3C-C4C-CHD-C1D
4	B	405	LBV	N_C-C4C-CHD-C1D
3	B	404	MPD	O2-C2-C3-C4
3	B	404	MPD	CM-C2-C3-C4
4	B	405	LBV	C2A-C3A-CAA-CBA
3	B	404	MPD	C1-C2-C3-C4
3	A	404	MPD	C2-C3-C4-C5


There are no ring outliers.


3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	403	ACT	2	0
2	A	403	ACT	1	0
4	A	405	LBV	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand LBV A 405





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	314/343 (91%)	0.10	21 (6%) 17 16	11, 20, 57, 105	0
1	B	314/343 (91%)	0.14	19 (6%) 21 19	10, 17, 55, 88	0
All	All	628/686 (91%)	0.12	40 (6%) 19 18	10, 18, 55, 105	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	132	TRP	16.2
1	B	105	TRP	10.2
1	A	132	TRP	9.7
1	B	322	LEU	7.9
1	B	318	VAL	6.6
1	B	321	ALA	6.1
1	A	323	GLU	5.0
1	A	67	THR	4.7
1	B	65	GLU	4.7
1	B	109	GLY	4.4
1	B	131	ALA	4.3
1	A	324	HIS	4.0
1	A	134	SER	3.7
1	A	64	GLN	3.7
1	A	90	PRO	3.5
1	A	65	GLU	3.5
1	B	138	HIS	3.4
1	B	137	PRO	3.3
1	A	131	ALA	3.1
1	A	140	LEU	3.1
1	A	66	PRO	3.1
1	A	138	HIS	3.0
1	B	64	GLN	2.9
1	B	139	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	61	PHE	2.7
1	B	66	PRO	2.6
1	B	60	THR	2.6
1	A	62	LEU	2.5
1	B	320	GLU	2.4
1	A	84	ALA	2.3
1	A	71	GLY	2.3
1	B	97	LEU	2.3
1	A	137	PRO	2.2
1	B	63	GLY	2.2
1	A	77	LEU	2.2
1	B	136	GLY	2.1
1	A	322	LEU	2.1
1	B	95	ASP	2.1
1	A	69	LEU	2.0
1	A	136	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ACT	A	401	4/4	0.84	0.16	41,44,45,54	0
2	ACT	B	403	4/4	0.88	0.10	25,26,26,30	0
2	ACT	A	402	4/4	0.90	0.08	28,45,49,49	0
3	MPD	A	404	8/8	0.91	0.12	22,25,31,32	8
2	ACT	B	401	4/4	0.91	0.12	61,63,64,75	0
3	MPD	B	404	8/8	0.93	0.10	17,24,28,31	8
2	ACT	A	403	4/4	0.94	0.10	31,36,43,49	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	LBV	A	405	43/43	0.96	0.08	9,11,19,31	0
2	ACT	B	402	4/4	0.96	0.06	27,36,39,60	0
4	LBV	B	405	43/43	0.97	0.08	8,10,18,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.