



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 06:07 am BST

PDB ID : 5FUF  
Title : Crystal structure of the Mep2 mutant S453D from *Candida albicans*  
Authors : van den Berg, B.; Chembath, A.; Jefferies, D.; Basle, A.; Khalid, S.; Rutherford, J.  
Deposited on : 2016-01-26  
Resolution : 2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2684 (2.08-2.04)
Clashscore	141614	2801 (2.08-2.04)
Ramachandran outliers	138981	2768 (2.08-2.04)
Sidechain outliers	138945	2768 (2.08-2.04)
RSRZ outliers	127900	2646 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	486	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	DMU	A	1457	-	-	-	X

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3675 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called AMMONIUM TRANSPORTER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	449	3459	2269	558	605	27	0	2	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	481	HIS	-	expression tag	UNP Q59UP8
A	482	HIS	-	expression tag	UNP Q59UP8
A	483	HIS	-	expression tag	UNP Q59UP8
A	484	HIS	-	expression tag	UNP Q59UP8
A	485	HIS	-	expression tag	UNP Q59UP8
A	486	HIS	-	expression tag	UNP Q59UP8
A	453	ASP	SER	engineered mutation	UNP Q59UP8

- Molecule 2 is DECYL-BETA-D-MALTOPYRANOSIDE (three-letter code: DMU) (formula:  $C_{22}H_{42}O_{11}$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			33	22	11		
2	A	1	Total	C	O	0	0
			33	22	11		
2	A	1	Total	C	O	0	0
			28	17	11		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	122	Total	O	0	0
			122	122		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: AMMONIUM TRANSPORTER



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	109.40 Å   109.40 Å   341.91 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	29.53 – 2.07 29.53 – 2.07	Depositor EDS
% Data completeness (in resolution range)	99.3 (29.53-2.07) 96.6 (29.53-2.07)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.72 (at 2.06 Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.162 , 0.174 0.162 , 0.177	Depositor DCC
$R_{free}$ test set	2000 reflections (4.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.0	Xtriage
Anisotropy	0.376	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 65.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3675	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.44	0/3570	0.54	0/4870

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3459	0	3372	17	0
2	A	94	0	113	0	0
3	A	122	0	0	0	0
All	All	3675	0	3485	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:213:LYS:O	1:A:213:LYS:HG3	1.90	0.72

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:249:MET:HA	1:A:252:TRP:CD1	2.30	0.65
1:A:420:GLU:OE1	1:A:452:ARG:NH1	2.30	0.65
1:A:438:ASP:HA	1:A:439:ASP:HB2	1.83	0.60
1:A:454:THR:O	1:A:454:THR:HG22	2.03	0.58
1:A:50[B]:SER:HB3	1:A:61:LEU:HB2	1.86	0.56
1:A:89:GLY:HA3	1:A:164:TYR:OH	2.05	0.56
1:A:50[A]:SER:HB2	1:A:61:LEU:HB2	1.89	0.54
1:A:90:SER:O	1:A:91:VAL:HB	2.07	0.53
1:A:91:VAL:HA	1:A:170:ASN:O	2.09	0.52
1:A:423:LEU:O	1:A:425:THR:N	2.43	0.51
1:A:425:THR:HG22	1:A:426:ASP:H	1.76	0.50
1:A:423:LEU:O	1:A:425:THR:OG1	2.21	0.48
1:A:253:TYR:CZ	1:A:306:VAL:HG23	2.49	0.47
1:A:257:ASN:OD1	1:A:306:VAL:HA	2.14	0.47
1:A:410:PRO:O	1:A:452:ARG:NH2	2.48	0.46
1:A:308:THR:HA	1:A:311:ILE:HD12	1.99	0.44

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	447/486 (92%)	439 (98%)	5 (1%)	3 (1%)	22 11

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	424	GLY
1	A	439	ASP
1	A	91	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	351/379 (93%)	348 (99%)	3 (1%)	78	78

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	79	TRP
1	A	236	TRP
1	A	419	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	DMU	A	1458	-	29,29,34	1.57	8 (27%)	40,40,45	1.15	3 (7%)
2	DMU	A	1457	-	34,34,34	1.45	7 (20%)	45,45,45	1.13	4 (8%)
2	DMU	A	1456	-	34,34,34	1.59	8 (23%)	45,45,45	1.10	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DMU	A	1458	-	-	7/14/54/59	0/2/2/2
2	DMU	A	1457	-	-	4/19/59/59	0/2/2/2
2	DMU	A	1456	-	-	12/19/59/59	0/2/2/2

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1456	DMU	O1-C9	4.40	1.55	1.44
2	A	1457	DMU	O1-C9	3.91	1.53	1.44
2	A	1458	DMU	O1-C9	3.91	1.53	1.44
2	A	1458	DMU	C11-C9	-2.85	1.42	1.51
2	A	1456	DMU	C7-C5	-2.72	1.45	1.52
2	A	1456	DMU	C11-C9	-2.65	1.43	1.51
2	A	1457	DMU	C11-C9	-2.65	1.43	1.51
2	A	1456	DMU	O5-C6	2.64	1.48	1.41
2	A	1456	DMU	O1-C10	2.56	1.48	1.41
2	A	1458	DMU	C7-C5	-2.53	1.45	1.52
2	A	1458	DMU	O4-C7	2.51	1.48	1.43
2	A	1456	DMU	O4-C7	2.48	1.48	1.43
2	A	1457	DMU	O4-C7	2.46	1.48	1.43
2	A	1456	DMU	O5-C4	2.42	1.50	1.44
2	A	1457	DMU	C7-C5	-2.35	1.46	1.52
2	A	1457	DMU	O5-C6	2.29	1.47	1.41
2	A	1458	DMU	O1-C10	2.23	1.47	1.41
2	A	1457	DMU	O3-C5	2.21	1.48	1.43
2	A	1458	DMU	O5-C6	2.20	1.47	1.41
2	A	1458	DMU	O3-C5	2.17	1.48	1.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1456	DMU	O3-C5	2.17	1.48	1.43
2	A	1457	DMU	O1-C10	2.14	1.47	1.41
2	A	1458	DMU	O5-C4	2.12	1.49	1.44

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1458	DMU	O5-C4-C3	3.21	116.52	109.75
2	A	1457	DMU	C10-O7-C3	-2.75	111.17	117.96
2	A	1457	DMU	C8-C7-C5	2.73	115.59	110.82
2	A	1458	DMU	O16-C6-C1	2.65	112.45	108.30
2	A	1458	DMU	C10-O7-C3	-2.46	111.87	117.96
2	A	1457	DMU	C7-C8-C9	2.27	114.29	110.24
2	A	1456	DMU	C1-C2-C3	2.27	114.86	109.68
2	A	1456	DMU	C18-O16-C6	2.14	117.39	113.84
2	A	1457	DMU	C10-C5-C7	2.11	114.39	110.00

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1456	DMU	O5-C6-O16-C18
2	A	1456	DMU	O6-C11-C9-O1
2	A	1456	DMU	O6-C11-C9-C8
2	A	1458	DMU	C1-C6-O16-C18
2	A	1456	DMU	O5-C4-C57-O61
2	A	1458	DMU	O5-C4-C57-O61
2	A	1456	DMU	O16-C18-C19-C22
2	A	1456	DMU	C5-C10-O7-C3
2	A	1456	DMU	C25-C28-C31-C34
2	A	1456	DMU	O1-C10-O7-C3
2	A	1457	DMU	C18-C19-C22-C25
2	A	1457	DMU	C25-C28-C31-C34
2	A	1458	DMU	O6-C11-C9-O1
2	A	1456	DMU	C3-C4-C57-O61
2	A	1456	DMU	C28-C31-C34-C37
2	A	1458	DMU	C19-C18-O16-C6
2	A	1458	DMU	C19-C22-C25-C28
2	A	1458	DMU	O5-C6-O16-C18
2	A	1456	DMU	C19-C22-C25-C28
2	A	1457	DMU	O16-C18-C19-C22
2	A	1456	DMU	C31-C34-C37-C40

*Continued on next page...*


*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	A	1458	DMU	C3-C4-C57-O61
2	A	1457	DMU	C19-C22-C25-C28


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	449/486 (92%)	0.13	32 (7%) 16 16	32, 45, 73, 143	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	435	TYR	7.2
1	A	427	LEU	6.9
1	A	454	THR	6.3
1	A	437	ALA	6.3
1	A	455	THR	5.6
1	A	426	ASP	4.5
1	A	436	TYR	4.2
1	A	4	GLN	4.2
1	A	92	PHE	3.8
1	A	329	LEU	3.6
1	A	411	PHE	3.1
1	A	5	PHE	2.8
1	A	133	LEU	2.8
1	A	172	TRP	2.8
1	A	424	GLY	2.6
1	A	453	ASP	2.5
1	A	137	ALA	2.4
1	A	409	ILE	2.4
1	A	408	ARG	2.4
1	A	134	MET	2.3
1	A	419	ASP	2.2
1	A	404	LEU	2.2
1	A	338	VAL	2.2
1	A	91	VAL	2.2
1	A	195	ALA	2.1
1	A	383	TYR	2.1
1	A	262	ALA	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	183	GLY	2.0
1	A	291	ILE	2.0
1	A	6	THR	2.0
1	A	129	VAL	2.0
1	A	275	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	DMU	A	1457	33/33	0.78	0.48	70,95,102,103	0
2	DMU	A	1456	33/33	0.79	0.34	69,95,111,113	0
2	DMU	A	1458	28/33	0.87	0.25	71,79,95,98	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


**Electron density around DMU A 1457:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around DMU A 1456:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers ⓘ

There are no such residues in this entry.