



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 04:14 pm BST

PDB ID : 3FWS  
Title : Crystal Structure of the CBS domains from the Bacillus subtilis CcpN repressor complexed with AppNp, phosphate and magnesium ions  
Authors : Chaix, D.; Arold, S.; Hoh, F.; Declerck, N.  
Deposited on : 2009-01-19  
Resolution : 2.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	10434 (2.04-2.00)
Clashscore	141614	11643 (2.04-2.00)
Ramachandran outliers	138981	11493 (2.04-2.00)
Sidechain outliers	138945	11492 (2.04-2.00)
RSRZ outliers	127900	10220 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	<div> <div>4%</div> <div> <div></div> <div>63%</div> <div>25%</div> <div>•• 9%</div> </div> </div>
1	B	159	<div> <div>6%</div> <div> <div></div> <div>70%</div> <div>21%</div> <div>• 8%</div> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2446 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YqzB protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	145	Total	C	N	O	S	0	0	0
			1143	729	193	214	7			
1	B	146	Total	C	N	O	S	0	0	0
			1142	726	195	214	7			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	62	MET	-	EXPRESSION TAG	UNP O34994
A	213	LEU	-	EXPRESSION TAG	UNP O34994
A	214	GLN	-	EXPRESSION TAG	UNP O34994
A	215	HIS	-	EXPRESSION TAG	UNP O34994
A	216	HIS	-	EXPRESSION TAG	UNP O34994
A	217	HIS	-	EXPRESSION TAG	UNP O34994
A	218	HIS	-	EXPRESSION TAG	UNP O34994
A	219	HIS	-	EXPRESSION TAG	UNP O34994
A	220	HIS	-	EXPRESSION TAG	UNP O34994
B	62	MET	-	EXPRESSION TAG	UNP O34994
B	213	LEU	-	EXPRESSION TAG	UNP O34994
B	214	GLN	-	EXPRESSION TAG	UNP O34994
B	215	HIS	-	EXPRESSION TAG	UNP O34994
B	216	HIS	-	EXPRESSION TAG	UNP O34994
B	217	HIS	-	EXPRESSION TAG	UNP O34994
B	218	HIS	-	EXPRESSION TAG	UNP O34994
B	219	HIS	-	EXPRESSION TAG	UNP O34994
B	220	HIS	-	EXPRESSION TAG	UNP O34994

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C<sub>10</sub>H<sub>17</sub>N<sub>6</sub>O<sub>12</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 31	C 10	N 6	O 12	P 3	0	0
2	B	1	Total 31	C 10	N 6	O 12	P 3	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula:  $O_4P$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 5	O 4	P 1	0	0
3	B	1	Total 5	O 4	P 1	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	A	2	Total	Mg	0	0
			2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	36	Total	O	0	0
			36	36		
5	B	50	Total	O	0	0
			50	50		



- Molecule 1: YqzB protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.02Å 103.97Å 98.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.17 – 2.03 29.17 – 2.03	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.17-2.03) 99.8 (29.17-2.03)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.35 (at 2.03Å)	Xtriage
Refinement program	REFMAC 5.4.0062	Depositor
R, $R_{free}$	0.189 , 0.262 0.203 , 0.270	Depositor DCC
$R_{free}$ test set	934 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.2	Xtriage
Anisotropy	0.061	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 40.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	2446	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, ANP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.96	0/1155	0.97	1/1562 (0.1%)
1	B	1.03	1/1154 (0.1%)	0.96	0/1559
All	All	1.00	1/2309 (0.0%)	0.97	1/3121 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	162	TYR	CD1-CE1	5.03	1.46	1.39

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	94	VAL	CG1-CB-CG2	5.62	119.89	110.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	211	ILE	Peptide



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1143	0	1210	35	0
1	B	1142	0	1208	21	0
2	A	31	0	13	2	0
2	B	31	0	13	1	0
3	A	5	0	0	1	0
3	B	5	0	0	0	0
4	A	2	0	0	0	0
4	B	1	0	0	0	0
5	A	36	0	0	6	0
5	B	50	0	0	4	0
All	All	2446	0	2444	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:GLN:CB	1:A:137:GLN:HB2	1.85	1.06
1:A:136:GLN:HB3	1:A:137:GLN:HB2	1.06	1.06
1:A:159:ARG:HD2	5:A:226:HOH:O	1.55	1.06
1:A:160:GLU:HG3	5:A:244:HOH:O	1.69	0.91
1:A:136:GLN:HB3	1:A:137:GLN:CB	2.01	0.88
1:A:143:PRO:HD2	1:A:146:ILE:HD13	1.57	0.85
2:A:221:ANP:O2G	2:A:221:ANP:O1A	1.96	0.83
1:A:135:GLY:H	1:A:136:GLN:HB2	1.44	0.82
2:B:221:ANP:O1A	2:B:221:ANP:PG	2.45	0.74
1:A:151:MET:SD	1:A:154:ILE:HD12	2.29	0.72
1:B:159:ARG:NH2	5:B:238:HOH:O	2.22	0.71
1:A:100:ILE:HD13	1:B:134:ILE:HG21	1.79	0.64
1:A:136:GLN:CA	1:A:137:GLN:HB2	2.31	0.61
1:A:151:MET:HG2	1:A:154:ILE:HD12	1.83	0.59
2:A:221:ANP:O1A	2:A:221:ANP:PG	2.60	0.59
1:A:81:LYS:HB3	1:A:159:ARG:HB3	1.85	0.59
1:A:126:ARG:HD3	1:B:127:LYS:HD2	1.85	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:151:MET:CG	1:A:154:ILE:HD12	2.34	0.57
1:A:197:THR:HG22	1:A:201:LYS:HE3	1.87	0.56
3:A:222:PO4:O2	5:A:258:HOH:O	2.18	0.56
1:A:200:THR:HG22	1:B:171:ILE:HD11	1.88	0.56
1:B:81:LYS:HB3	1:B:159:ARG:HB3	1.87	0.56
1:A:159:ARG:CG	5:A:226:HOH:O	2.54	0.55
1:B:80:VAL:HG13	1:B:194:VAL:HG21	1.90	0.53
1:B:135:GLY:HA3	5:B:265:HOH:O	2.07	0.53
1:A:124:LEU:HD21	1:A:147:ILE:HG13	1.93	0.51
1:A:177:ALA:HB2	1:A:195:THR:HG22	1.94	0.49
1:B:65:LYS:N	1:B:69:GLN:OE1	2.38	0.48
1:A:159:ARG:CD	5:A:226:HOH:O	2.31	0.48
1:A:94:VAL:HG13	1:A:98:ASP:HB2	1.96	0.47
1:A:99:ALA:HB2	1:A:144:VAL:HG11	1.97	0.46
1:A:135:GLY:N	1:A:136:GLN:HB2	2.22	0.46
1:A:158:ARG:NH2	1:A:183:ASP:OD2	2.48	0.46
1:B:156:VAL:HG12	1:B:179:PRO:HG2	1.97	0.45
1:A:130:LEU:HD13	1:B:130:LEU:HA	1.97	0.45
1:B:182:LYS:NZ	1:B:184:THR:HG22	2.31	0.44
1:A:159:ARG:HG2	5:A:226:HOH:O	2.15	0.43
1:A:177:ALA:CB	1:A:195:THR:HG22	2.48	0.43
1:B:135:GLY:CA	5:B:265:HOH:O	2.64	0.43
1:B:182:LYS:HZ1	1:B:184:THR:HG22	1.84	0.42
1:A:73:ASP:HA	1:A:76:LYS:HZ2	1.84	0.42
1:A:124:LEU:HD11	1:A:144:VAL:HB	2.00	0.42
1:A:184:THR:C	1:A:186:LYS:H	2.23	0.42
1:B:124:LEU:HD11	1:B:144:VAL:HB	2.02	0.42
1:A:204:VAL:HA	1:B:164:MET:CE	2.50	0.42
1:B:86:ILE:O	1:B:193:ARG:NH1	2.48	0.42
1:B:138:GLU:N	5:B:230:HOH:O	2.42	0.42
1:A:103:MET:HB2	1:A:103:MET:HE2	1.75	0.41
1:A:112:PHE:HD1	1:A:123:VAL:HG12	1.85	0.41
1:A:143:PRO:HD2	1:A:146:ILE:CD1	2.41	0.41
1:B:123:VAL:HG13	1:B:154:ILE:HD12	2.03	0.41
1:B:139:LEU:O	1:B:142:VAL:HG22	2.21	0.40
1:B:181:ILE:HA	1:B:189:GLU:O	2.21	0.40
1:A:134:ILE:HD11	1:B:100:ILE:HD13	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	143/159 (90%)	137 (96%)	5 (4%)	1 (1%)	22	15
1	B	144/159 (91%)	139 (96%)	5 (4%)	0	100	100
All	All	287/318 (90%)	276 (96%)	10 (4%)	1 (0%)	41	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	137	GLN

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/146 (92%)	120 (90%)	14 (10%)	7	3
1	B	133/146 (91%)	127 (96%)	6 (4%)	27	23
All	All	267/292 (91%)	247 (92%)	20 (8%)	13	8

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	GLN
1	A	70	LEU
1	A	94	VAL
1	A	103	MET
1	A	137	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	138	GLU
1	A	139	LEU
1	A	140	THR
1	A	141	SER
1	A	146	ILE
1	A	172	GLU
1	A	183	ASP
1	A	185	ASP
1	A	212	LEU
1	B	74	LYS
1	B	105	LEU
1	B	119	VAL
1	B	182	LYS
1	B	186	LYS
1	B	208	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	137	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ANP	B	221	-	29,33,33	3.51	11 (37%)	31,52,52	4.09	11 (35%)
2	ANP	A	221	-	29,33,33	3.16	5 (17%)	31,52,52	1.52	7 (22%)
3	PO4	A	222	-	4,4,4	0.82	0	6,6,6	0.97	0
3	PO4	B	222	-	4,4,4	1.01	0	6,6,6	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	B	221	-	-	2/14/38/38	0/3/3/3
2	ANP	A	221	-	-	6/14/38/38	0/3/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	221	ANP	PG-O1G	11.30	1.64	1.46
2	B	221	ANP	PB-O3A	11.24	1.73	1.59
2	A	221	ANP	PB-O3A	11.02	1.73	1.59
2	A	221	ANP	PG-O1G	8.91	1.60	1.46
2	A	221	ANP	PG-O3G	4.86	1.69	1.56
2	B	221	ANP	PG-O3G	4.64	1.69	1.56
2	A	221	ANP	PG-N3B	4.30	1.74	1.63
2	A	221	ANP	PB-O2B	-4.24	1.45	1.56
2	B	221	ANP	O4'-C4'	3.76	1.53	1.45
2	B	221	ANP	PG-O2G	3.65	1.66	1.56
2	B	221	ANP	PG-N3B	3.04	1.71	1.63
2	B	221	ANP	PB-O2B	-2.88	1.49	1.56
2	B	221	ANP	O3'-C3'	2.77	1.49	1.43
2	B	221	ANP	C2'-C1'	2.31	1.57	1.53
2	B	221	ANP	O4'-C1'	2.27	1.44	1.41
2	B	221	ANP	C5'-C4'	2.06	1.58	1.51

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	221	ANP	O1B-PB-N3B	17.22	137.13	111.77
2	B	221	ANP	O3G-PG-O2G	8.47	130.21	107.64
2	B	221	ANP	O2G-PG-O1G	-5.75	99.00	113.45
2	B	221	ANP	N3-C2-N1	-5.30	120.39	128.68
2	A	221	ANP	N3-C2-N1	-4.67	121.38	128.68
2	B	221	ANP	O3A-PB-N3B	-4.35	94.53	106.59
2	B	221	ANP	O2B-PB-O3A	-4.06	91.10	104.64
2	B	221	ANP	O3G-PG-O1G	-3.03	105.84	113.45
2	B	221	ANP	O1G-PG-N3B	-2.66	107.86	111.77
2	A	221	ANP	O3A-PB-N3B	2.56	113.69	106.59
2	B	221	ANP	O2B-PB-O1B	2.50	115.17	109.92
2	A	221	ANP	O4'-C1'-C2'	-2.46	103.33	106.93
2	A	221	ANP	C2-N1-C6	2.41	122.88	118.75
2	A	221	ANP	PA-O3A-PB	-2.36	124.32	132.62
2	B	221	ANP	C3'-C2'-C1'	2.32	104.47	100.98
2	A	221	ANP	O2'-C2'-C3'	2.09	118.58	111.82
2	A	221	ANP	O4'-C4'-C5'	-2.07	102.55	109.37
2	B	221	ANP	C2-N1-C6	2.06	122.27	118.75

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	221	ANP	PB-N3B-PG-O1G
2	B	221	ANP	PG-N3B-PB-O1B
2	A	221	ANP	PB-N3B-PG-O1G
2	A	221	ANP	PG-N3B-PB-O1B
2	A	221	ANP	PG-N3B-PB-O3A
2	A	221	ANP	C5'-O5'-PA-O1A
2	A	221	ANP	C5'-O5'-PA-O2A
2	A	221	ANP	C5'-O5'-PA-O3A


There are no ring outliers.


3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	221	ANP	1	0
2	A	221	ANP	2	0
3	A	222	PO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	145/159 (91%)	0.64	7 (4%) 30 30	21, 29, 40, 56	0
1	B	146/159 (91%)	0.51	9 (6%) 20 19	21, 29, 37, 53	0
All	All	291/318 (91%)	0.57	16 (5%) 25 24	21, 29, 38, 56	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	68	THR	6.2
1	B	185	ASP	5.0
1	A	135	GLY	4.7
1	A	185	ASP	4.3
1	B	209	ASN	3.2
1	A	136	GLN	3.1
1	B	208	GLU	3.0
1	B	64	GLY	3.0
1	B	186	LYS	2.9
1	B	73	ASP	2.9
1	A	91	HIS	2.7
1	A	69	GLN	2.7
1	A	138	GLU	2.7
1	B	187	GLY	2.6
1	B	116	ARG	2.4
1	B	65	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	MG	A	223	1/1	0.89	0.24	40,40,40,40	0
3	PO4	B	222	5/5	0.93	0.15	49,49,51,51	0
3	PO4	A	222	5/5	0.96	0.18	46,50,50,51	0
2	ANP	A	221	31/31	0.96	0.09	17,22,33,39	0
2	ANP	B	221	31/31	0.97	0.09	17,20,33,38	0
4	MG	B	223	1/1	0.97	0.23	31,31,31,31	0
4	MG	A	224	1/1	0.98	0.05	44,44,44,44	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around ANP A 221:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ANP B 221:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers

There are no such residues in this entry.