



wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2020 – 12:18 am BST

PDB ID : 1FZ7
Title : METHANE MONOOXYGENASE HYDROXYLASE, FORM III SOAKED
IN 0.9 M ETHANOL
Authors : Whittington, D.A.; Sazinsky, M.H.; Lippard, S.J.
Deposited on : 2000-10-03
Resolution : 1.96 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	527	
1	B	527	
2	C	389	
2	D	389	
3	E	170	
3	F	170	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 18582 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called METHANE MONOOXYGENASE COMPONENT A, ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	510	4177	2673	719	767	18	0	0	0
1	B	510	4177	2673	719	767	18	0	0	0

- Molecule 2 is a protein called METHANE MONOOXYGENASE COMPONENT A, BETA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	388	3193	2054	551	580	8	0	0	0
2	D	388	3193	2054	551	580	8	7	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	370	ARG	ALA	CONFLICT	UNP P18798
D	370	ARG	ALA	CONFLICT	UNP P18798

- Molecule 3 is a protein called METHANE MONOOXYGENASE COMPONENT A, GAMMA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	166	1368	867	246	250	5	0	0	0
3	F	167	1377	872	248	252	5	0	0	0


- Molecule 4 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Fe	0	0
			2	2		
4	A	2	Total	Fe	0	0
			2	2		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		
5	C	2	Total	Ca	0	0
			2	2		

- Molecule 6 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			3	1	2		

- Molecule 7 is ETHANOL (three-letter code: EOH) (formula: C₂H₆O).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			3	2	1		

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	246	Total	O	0	0
			246	246		
8	B	225	Total	O	0	0
			225	225		
8	C	303	Total	O	0	0
			303	303		
8	D	116	Total	O	0	0
			116	116		
8	E	152	Total	O	0	0
			152	152		
8	F	42	Total	O	0	0
			42	42		




- Molecule 2: METHANE MONOOXYGENASE COMPONENT A, BETA CHAIN



- Molecule 2: METHANE MONOOXYGENASE COMPONENT A, BETA CHAIN




- Molecule 3: METHANE MONOOXYGENASE COMPONENT A, GAMMA CHAIN



- Molecule 3: METHANE MONOOXYGENASE COMPONENT A, GAMMA CHAIN





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.40Å 172.37Å 221.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.96 29.99 – 1.96	Depositor EDS
% Data completeness (in resolution range)	90.1 (30.00-1.96) 90.2 (29.99-1.96)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.19 (at 1.96Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.214 , 0.255 0.208 , 0.249	Depositor DCC
R_{free} test set	6226 reflections (3.24%)	wwPDB-VP
Wilson B-factor (Å ²)	28.1	Xtrriage
Anisotropy	0.260	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 58.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	18582	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMT, EOH, CA, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/4302	0.55	0/5842
1	B	0.33	0/4302	0.55	0/5842
2	C	0.38	0/3289	0.57	0/4464
2	D	0.31	0/3289	0.53	0/4464
3	E	0.34	0/1396	0.59	0/1880
3	F	0.28	0/1407	0.52	0/1896
All	All	0.33	0/17985	0.55	0/24388

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4177	0	3975	110	0
1	B	4177	0	3975	113	0
2	C	3193	0	3042	66	0
2	D	3193	0	3042	106	0
3	E	1368	0	1363	19	0
3	F	1377	0	1364	59	0
4	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	2	0	0	0	0
5	A	1	0	0	0	0
5	C	2	0	0	0	0
6	A	3	0	1	0	0
7	B	3	0	5	0	0
8	A	246	0	0	8	0
8	B	225	0	0	4	0
8	C	303	0	0	5	0
8	D	116	0	0	3	0
8	E	152	0	0	1	0
8	F	42	0	0	2	0
All	All	18582	0	16767	414	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 414 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:41:THR:HG23	3:F:43:PHE:H	1.21	1.06
2:C:270:PRO:HB3	2:D:270:PRO:HB3	1.37	1.05
1:A:78:GLN:HE22	1:A:150:GLN:HE21	1.05	0.95
1:A:352:ALA:HA	1:A:404:PRO:HB2	1.49	0.95
1:B:78:GLN:HE22	1:B:150:GLN:HE21	1.17	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	508/527 (96%)	486 (96%)	20 (4%)	2 (0%)	34 22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	508/527 (96%)	483 (95%)	24 (5%)	1 (0%)	47 38
2	C	386/389 (99%)	377 (98%)	8 (2%)	1 (0%)	41 30
2	D	386/389 (99%)	357 (92%)	25 (6%)	4 (1%)	15 6
3	E	164/170 (96%)	162 (99%)	2 (1%)	0	100 100
3	F	165/170 (97%)	156 (94%)	7 (4%)	2 (1%)	13 4
All	All	2117/2172 (98%)	2021 (96%)	86 (4%)	10 (0%)	29 17

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	40	LYS
1	A	311	GLU
2	D	64	ALA
3	F	122	ILE
1	A	340	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	432/442 (98%)	419 (97%)	13 (3%)	41 30
1	B	432/442 (98%)	423 (98%)	9 (2%)	53 46
2	C	322/323 (100%)	317 (98%)	5 (2%)	62 58
2	D	322/323 (100%)	316 (98%)	6 (2%)	57 50
3	E	144/147 (98%)	141 (98%)	3 (2%)	53 46
3	F	145/147 (99%)	141 (97%)	4 (3%)	43 33
All	All	1797/1824 (98%)	1757 (98%)	40 (2%)	52 44

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	186	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	C	35	MET
3	F	11	THR
1	B	310	TYR
2	C	153	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 62 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	108	ASN
1	B	268	ASN
3	F	7	HIS
1	B	168	HIS
1	B	278	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 7 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	FMT	A	9001	4	0,2,2	0.00	-	0,1,1	0.00	-
7	EOH	B	9002	4	2,2,2	0.50	0	1,1,1	0.14	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	510/527 (96%)	0.50	48 (9%) 8 13	22, 36, 62, 78	0
1	B	510/527 (96%)	0.32	35 (6%) 16 25	22, 36, 63, 78	0
2	C	388/389 (99%)	-0.01	7 (1%) 68 76	18, 26, 46, 62	0
2	D	388/389 (99%)	0.83	54 (13%) 2 4	26, 46, 68, 82	2 (0%)
3	E	166/170 (97%)	-0.04	2 (1%) 79 84	20, 29, 46, 69	0
3	F	167/170 (98%)	1.47	44 (26%) 0 0	37, 56, 75, 86	0
All	All	2129/2172 (98%)	0.46	190 (8%) 9 15	18, 37, 66, 86	2 (0%)

The worst 5 of 190 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	244	LEU	6.1
1	A	19	ALA	5.7
2	D	205	PRO	5.5
1	B	316	ILE	5.5
1	A	316	ILE	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	EOH	B	9002	3/3	0.86	0.21	46,46,47,49	0
6	FMT	A	9001	3/3	0.93	0.12	51,51,53,57	0
5	CA	C	5006	1/1	0.94	0.04	65,65,65,65	0
4	FE	A	5002	1/1	0.98	0.03	49,49,49,49	0
5	CA	C	5007	1/1	0.99	0.06	40,40,40,40	0
4	FE	B	5004	1/1	0.99	0.03	44,44,44,44	0
5	CA	A	5005	1/1	0.99	0.06	40,40,40,40	0
4	FE	B	5003	1/1	0.99	0.03	34,34,34,34	0
4	FE	A	5001	1/1	0.99	0.02	42,42,42,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.