



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2021 – 10:26 AM EDT

PDB ID : 3FZY  
Title : Crystal Structure of Pre-cleavage Form of Cysteine Protease Domain from *Vibrio cholerae* RtxA Toxin  
Authors : Shuvalova, L.; Minasov, G.; Prochazkova, K.; Satchell, K.J.F.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)  
Deposited on : 2009-01-26  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	234	 81% 11% • 8%
1	B	234	 80% 10% 10%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3890 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RTX toxin RtxA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	216	Total	C	N	O	S	0	17	0
			1794	1101	327	363	3			
1	B	211	Total	C	N	O	S	0	9	0
			1690	1030	313	345	2			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	MET	-	expression tag	UNP Q9KS12
A	-22	HIS	-	expression tag	UNP Q9KS12
A	-21	HIS	-	expression tag	UNP Q9KS12
A	-20	HIS	-	expression tag	UNP Q9KS12
A	-19	HIS	-	expression tag	UNP Q9KS12
A	-18	HIS	-	expression tag	UNP Q9KS12
A	-17	HIS	-	expression tag	UNP Q9KS12
A	-16	SER	-	expression tag	UNP Q9KS12
A	-15	SER	-	expression tag	UNP Q9KS12
A	-14	GLY	-	expression tag	UNP Q9KS12
A	-13	VAL	-	expression tag	UNP Q9KS12
A	-12	ASP	-	expression tag	UNP Q9KS12
A	-11	LEU	-	expression tag	UNP Q9KS12
A	-10	GLY	-	expression tag	UNP Q9KS12
A	-9	THR	-	expression tag	UNP Q9KS12
A	-8	GLU	-	expression tag	UNP Q9KS12
A	-7	ASN	-	expression tag	UNP Q9KS12
A	-6	LEU	-	expression tag	UNP Q9KS12
A	-5	TYR	-	expression tag	UNP Q9KS12
A	-4	PHE	-	expression tag	UNP Q9KS12
A	-3	GLN	-	expression tag	UNP Q9KS12
A	-2	SER	-	expression tag	UNP Q9KS12
A	-1	ASN	-	expression tag	UNP Q9KS12
A	3568	SER	CYS	engineered mutation	UNP Q9KS12
B	-23	MET	-	expression tag	UNP Q9KS12

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-22	HIS	-	expression tag	UNP Q9KS12
B	-21	HIS	-	expression tag	UNP Q9KS12
B	-20	HIS	-	expression tag	UNP Q9KS12
B	-19	HIS	-	expression tag	UNP Q9KS12
B	-18	HIS	-	expression tag	UNP Q9KS12
B	-17	HIS	-	expression tag	UNP Q9KS12
B	-16	SER	-	expression tag	UNP Q9KS12
B	-15	SER	-	expression tag	UNP Q9KS12
B	-14	GLY	-	expression tag	UNP Q9KS12
B	-13	VAL	-	expression tag	UNP Q9KS12
B	-12	ASP	-	expression tag	UNP Q9KS12
B	-11	LEU	-	expression tag	UNP Q9KS12
B	-10	GLY	-	expression tag	UNP Q9KS12
B	-9	THR	-	expression tag	UNP Q9KS12
B	-8	GLU	-	expression tag	UNP Q9KS12
B	-7	ASN	-	expression tag	UNP Q9KS12
B	-6	LEU	-	expression tag	UNP Q9KS12
B	-5	TYR	-	expression tag	UNP Q9KS12
B	-4	PHE	-	expression tag	UNP Q9KS12
B	-3	GLN	-	expression tag	UNP Q9KS12
B	-2	SER	-	expression tag	UNP Q9KS12
B	-1	ASN	-	expression tag	UNP Q9KS12
B	3568	SER	CYS	engineered mutation	UNP Q9KS12

- Molecule 2 is INOSITOL HEXAKISPHOSPHATE (three-letter code: IHP) (formula:  $C_6H_{18}O_{24}P_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			36	6	24	6		
2	B	1	Total	C	O	P	0	0
			36	6	24	6		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	12	Total	X	0	0
			12	12		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	178	Total	O	0	11
			183	183		
5	B	135	Total	O	0	6
			138	138		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.05Å 66.37Å 137.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.67 – 1.95 23.67 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.6 (23.67-1.95) 99.6 (23.67-1.95)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.31 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.5.0051	Depositor
R, $R_{free}$	0.169 , 0.216 0.233 , 0.272	Depositor DCC
$R_{free}$ test set	1584 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.9	Xtriage
Anisotropy	0.637	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 46.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3890	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, UNX, IHP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.69	0/1823	0.80	0/2467
1	B	0.58	0/1717	0.74	0/2324
All	All	0.64	0/3540	0.77	0/4791

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1794	0	1711	24	0
1	B	1690	0	1609	17	0
2	A	36	0	6	1	0
2	B	36	0	6	1	0
3	A	1	0	0	0	0
4	A	12	0	0	0	0
5	A	183	0	0	2	0
5	B	138	0	0	1	0
All	All	3890	0	3332	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.



All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3547:GLN:OE1	1:A:3592[A]:LEU:HD21	1.52	1.07
1:A:3526:THR:HG22	1:A:3577[A]:LYS:HE2	1.66	0.76
1:A:3587:MET:HE3	1:A:3592[B]:LEU:HD23	1.69	0.73
1:B:3429:ALA:HB2	1:B:3568:SER:HB2	1.78	0.66
1:A:3547:GLN:OE1	1:A:3592[A]:LEU:CD2	2.39	0.65
1:A:-3[B]:GLN:HE22	1:A:3614:LYS:HZ2	1.42	0.65
1:A:3542:LYS:NZ	1:A:3542:LYS:HB2	2.12	0.64
1:A:3587:MET:CE	1:A:3592[B]:LEU:HD23	2.29	0.63
1:A:3494:SER:HA	1:A:3542:LYS:HE2	1.81	0.62
1:B:3606[A]:ASP:OD1	1:B:3612:HIS:NE2	2.31	0.62
1:A:3592[B]:LEU:HG	1:A:3594:VAL:HG13	1.83	0.60
1:A:3547:GLN:HE22	1:A:3559:LYS:HB3	1.67	0.58
1:B:3610:ARG:NH2	2:B:8001:IHP:O21	2.40	0.55
1:B:3547:GLN:OE1	1:B:3592:LEU:HD21	2.07	0.54
1:B:-3:GLN:HG3	1:B:-1:ASN:H	1.73	0.54
1:A:3557:ASN:H	1:A:3557:ASN:HD22	1.58	0.50
1:B:3534:TYR:CZ	1:B:3542:LYS:HE3	2.47	0.50
1:A:3557:ASN:HD22	1:A:3557:ASN:N	2.11	0.49
1:A:-7:ASN:O	1:A:-3[B]:GLN:HG2	2.13	0.49
1:A:3547:GLN:NE2	1:A:3559:LYS:HB3	2.28	0.47
1:A:3509[B]:ASP:OD1	1:A:3510:GLY:N	2.46	0.47
1:A:3542:LYS:HB2	1:A:3542:LYS:HZ3	1.78	0.47
1:A:3507[B]:LYS:O	1:A:3507[B]:LYS:CG	2.63	0.47
1:A:3550:PHE:CE2	1:A:3556:ILE:HD12	2.50	0.46
1:A:3556:ILE:HG22	1:A:3557:ASN:N	2.31	0.46
1:A:3582:GLN:NE2	5:A:191:HOH:O	2.49	0.46
1:B:3606[B]:ASP:OD1	1:B:3606[B]:ASP:C	2.54	0.45
1:B:3617[B]:ASN:O	1:B:3617[B]:ASN:OD1	2.35	0.45
1:B:3547:GLN:NE2	1:B:3559:LYS:HG2	2.31	0.45
1:A:3429:ALA:HB2	1:A:3568:SER:HB2	1.98	0.45
1:A:3514:TRP:HB2	1:A:3563[B]:ILE:HD13	1.99	0.44
1:B:-3:GLN:HB3	1:B:3620:TRP:CZ2	2.52	0.44
1:B:3611:LYS:HD2	5:B:50:HOH:O	2.18	0.44
1:B:3606[B]:ASP:OD1	1:B:3609:GLY:N	2.52	0.43
1:B:3603:LEU:HD11	1:B:3611:LYS:HG2	2.01	0.42
1:B:3563:ILE:HD12	1:B:3594:VAL:HG21	2.01	0.42
2:A:8000:IHP:O23	2:A:8000:IHP:O24	2.37	0.42
1:A:3551:ASN:HB3	5:A:188:HOH:O	2.19	0.41
1:B:3548[B]:GLN:NE2	1:B:3548[B]:GLN:HA	2.35	0.41
1:A:3563[A]:ILE:HD12	1:A:3594:VAL:HG21	2.01	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3504:ASP:HB3	1:B:3507:LYS:HD3	2.02	0.40
1:B:3547:GLN:HE21	1:B:3559:LYS:HA	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	231/234 (99%)	219 (95%)	10 (4%)	2 (1%)	17	8
1	B	218/234 (93%)	199 (91%)	18 (8%)	1 (0%)	29	17
All	All	449/468 (96%)	418 (93%)	28 (6%)	3 (1%)	22	11

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3454	GLY
1	B	3453	GLY
1	A	3527	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/194 (100%)	192 (98%)	3 (2%)	65	60

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	184/194 (95%)	183 (100%)	1 (0%)	88	88
All	All	379/388 (98%)	375 (99%)	4 (1%)	73	71

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3542	LYS
1	A	3552	GLN
1	A	3557	ASN
1	B	3557	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	-1	ASN
1	A	3552	GLN
1	A	3557	ASN
1	B	3552	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 1 is monoatomic and 12 are unknown - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	IHP	A	8000	-	36,36,36	0.88	1 (2%)	54,60,60	1.18	6 (11%)
2	IHP	B	8001	-	36,36,36	0.94	2 (5%)	54,60,60	1.36	6 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IHP	A	8000	-	-	3/30/54/54	0/1/1/1
2	IHP	B	8001	-	-	3/30/54/54	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	8001	IHP	P4-O14	2.50	1.64	1.59
2	A	8000	IHP	P5-O15	2.32	1.63	1.59
2	B	8001	IHP	P1-O11	2.19	1.63	1.59

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	8001	IHP	O11-P1-O21	-4.02	93.87	109.39
2	B	8001	IHP	O43-P3-O33	3.22	119.95	107.64
2	A	8000	IHP	O13-P3-O23	-2.82	98.51	109.39
2	A	8000	IHP	O11-P1-O21	-2.78	98.67	109.39
2	B	8001	IHP	C5-C4-C3	-2.64	104.62	110.41
2	A	8000	IHP	O41-P1-O31	2.57	117.45	107.64
2	B	8001	IHP	O41-P1-O31	2.41	116.85	107.64
2	B	8001	IHP	O15-C5-C4	2.27	114.04	108.69
2	A	8000	IHP	O15-C5-C6	2.27	114.03	108.69
2	B	8001	IHP	O12-C2-C3	2.24	113.97	108.69
2	A	8000	IHP	O46-P6-O36	2.21	116.10	107.64
2	A	8000	IHP	O15-P5-O25	-2.09	101.31	109.39

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	8000	IHP	C2-O12-P2-O22
2	B	8001	IHP	C2-O12-P2-O22
2	B	8001	IHP	C2-O12-P2-O42
2	B	8001	IHP	C6-O16-P6-O26
2	A	8000	IHP	C6-O16-P6-O26
2	A	8000	IHP	C6-O16-P6-O36


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	8000	IHP	1	0
2	B	8001	IHP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands



Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around IHP A 8000:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around IHP B 8001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.