



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 20, 2020 – 09:00 AM BST

PDB ID : 4FZG
Title : 20S yeast proteasome in complex with glidobactin
Authors : Stein, M.; Beck, P.; Kaiser, M.; Dudler, R.; Becker, C.F.W.; Groll, M.
Deposited on : 2012-07-06
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




























Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	<div> <div>%</div> <div>96%</div> <div>.</div> </div>
1	O	250	<div> <div>2%</div> <div>96%</div> <div>.</div> </div>
2	B	244	<div> <div>2%</div> <div>93%</div> <div>7%</div> </div>
2	P	244	<div> <div>2%</div> <div>93%</div> <div>7%</div> </div>
3	C	241	<div> <div>%</div> <div>95%</div> <div>5%</div> </div>
3	Q	241	<div> <div>2%</div> <div>95%</div> <div>5%</div> </div>
4	D	242	<div> <div>2%</div> <div>94%</div> <div>6%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	R	242	
5	E	233	
5	S	233	
6	F	244	
6	T	244	
7	G	243	
7	U	243	
8	H	222	
8	V	222	
9	I	204	
9	W	204	
10	J	198	
10	X	198	
11	K	212	
11	Y	212	
12	L	222	
12	Z	222	
13	M	233	
13	a	233	
14	N	196	
14	b	196	
15	c	4	
15	d	4	
15	e	4	
15	f	4	

2 Entry composition

There are 16 unique types of molecules in this entry. The entry contains 50986 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome component Y7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			
1	O	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			

- Molecule 2 is a protein called Proteasome component Y13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	244	Total	C	N	O	S	0	0	0
			1905	1201	321	380	3			
2	P	244	Total	C	N	O	S	0	0	0
			1905	1201	321	380	3			

- Molecule 3 is a protein called Proteasome component PRE6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	241	Total	C	N	O	S	0	0	0
			1891	1181	331	375	4			
3	Q	241	Total	C	N	O	S	0	0	0
			1891	1181	331	375	4			

- Molecule 4 is a protein called Proteasome component PUP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	242	Total	C	N	O	S	0	0	0
			1862	1162	314	379	7			
4	R	242	Total	C	N	O	S	0	0	0
			1862	1162	314	379	7			

- Molecule 5 is a protein called Proteasome component PRE5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	233	Total	C	N	O	S	0	0	0
			1795	1129	312	350	4			
5	S	233	Total	C	N	O	S	0	0	0
			1795	1129	312	350	4			

- Molecule 6 is a protein called Proteasome component C1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	244	Total	C	N	O	S	0	0	0
			1897	1205	330	358	4			
6	T	244	Total	C	N	O	S	0	0	0
			1897	1205	330	358	4			

- Molecule 7 is a protein called Proteasome component C7-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	243	Total	C	N	O	S	0	0	0
			1921	1221	322	370	8			
7	U	243	Total	C	N	O	S	0	0	0
			1921	1221	322	370	8			

- Molecule 8 is a protein called Proteasome component PUP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	222	Total	C	N	O	S	0	0	0
			1685	1061	293	324	7			
8	V	222	Total	C	N	O	S	0	0	0
			1685	1061	293	324	7			

- Molecule 9 is a protein called Proteasome component PUP3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			
9	W	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			

- Molecule 10 is a protein called Proteasome component C11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	198	Total	C	N	O	S	0	0	0
			1585	1005	269	305	6			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	X	198	Total	C	N	O	S	0	0	0
			1585	1005	269	305	6			

- Molecule 11 is a protein called Proteasome component PRE2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			
11	Y	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			

- Molecule 12 is a protein called Proteasome component C5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			
12	Z	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			

- Molecule 13 is a protein called Proteasome component PRE4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			
13	a	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			

- Molecule 14 is a protein called Proteasome component PRE3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			
14	b	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			

- Molecule 15 is a protein called Glidobactin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
15	c	4	Total	C	N	O	0	0	0
			37	27	4	6			
15	d	4	Total	C	N	O	0	0	0
			37	27	4	6			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
15	e	4	Total	C	N	O	0	0	0
			37	27	4	6			
15	f	4	Total	C	N	O	0	0	0
			37	27	4	6			

- Molecule 16 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	A	56	Total	O	0	0
			56	56		
16	B	35	Total	O	0	0
			35	35		
16	C	41	Total	O	0	0
			41	41		
16	D	39	Total	O	0	0
			39	39		
16	E	25	Total	O	0	0
			25	25		
16	F	45	Total	O	0	0
			45	45		
16	G	59	Total	O	0	0
			59	59		
16	H	52	Total	O	0	0
			52	52		
16	I	61	Total	O	0	0
			61	61		
16	J	55	Total	O	0	0
			55	55		
16	K	41	Total	O	0	0
			41	41		
16	L	57	Total	O	0	0
			57	57		
16	M	63	Total	O	0	0
			63	63		
16	N	61	Total	O	0	0
			61	61		
16	O	34	Total	O	0	0
			34	34		
16	P	27	Total	O	0	0
			27	27		
16	Q	29	Total	O	0	0
			29	29		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	R	31	Total 31	O 31	0	0
16	S	21	Total 21	O 21	0	0
16	T	36	Total 36	O 36	0	0
16	U	63	Total 63	O 63	0	0
16	V	46	Total 46	O 46	0	0
16	W	47	Total 47	O 47	0	0
16	X	47	Total 47	O 47	0	0
16	Y	44	Total 44	O 44	0	0
16	Z	49	Total 49	O 49	0	0
16	a	72	Total 72	O 72	0	0
16	b	54	Total 54	O 54	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome component Y7




- Molecule 1: Proteasome component Y7




- Molecule 2: Proteasome component Y13



- Molecule 2: Proteasome component Y13



- Molecule 3: Proteasome component PRE6



- Molecule 3: Proteasome component PRE6



- Molecule 4: Proteasome component PUP2



- Molecule 4: Proteasome component PUP2



- Molecule 5: Proteasome component PRE5



- Molecule 5: Proteasome component PRE5




- Molecule 6: Proteasome component C1




- Molecule 6: Proteasome component C1




- Molecule 7: Proteasome component C7-alpha




- Molecule 7: Proteasome component C7-alpha




- Molecule 8: Proteasome component PUP1




- Molecule 8: Proteasome component PUP1




- Molecule 9: Proteasome component PUP3




- Molecule 9: Proteasome component PUP3




- Molecule 10: Proteasome component C11




• Molecule 10: Proteasome component C11




• Molecule 11: Proteasome component PRE2




• Molecule 11: Proteasome component PRE2




• Molecule 12: Proteasome component C5




• Molecule 12: Proteasome component C5



• Molecule 13: Proteasome component PRE4




• Molecule 13: Proteasome component PRE4



- Molecule 14: Proteasome component PRE3

Chain N:  96% .



- Molecule 14: Proteasome component PRE3

Chain b:  95% 5%



- Molecule 15: Glidobactin

Chain c:  75% 25%



- Molecule 15: Glidobactin

Chain d:  100%

There are no outlier residues recorded for this chain.

- Molecule 15: Glidobactin

Chain e:  100%

There are no outlier residues recorded for this chain.

- Molecule 15: Glidobactin

Chain f:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	134.77Å 300.35Å 144.49Å 90.00° 112.67° 90.00°	Depositor
Resolution (Å)	15.00 – 3.00 24.92 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.5 (15.00-3.00) 99.6 (24.92-3.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.43 (at 2.99Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.197 , 0.226 0.195 , 0.195	Depositor DCC
R_{free} test set	10479 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	66.6	Xtriage
Anisotropy	0.875	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 61.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	50986	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 0W6, LY0, MH9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/1952	0.63	0/2642
1	O	0.38	0/1952	0.63	0/2642
2	B	0.38	0/1935	0.63	0/2618
2	P	0.39	0/1935	0.64	0/2618
3	C	0.35	0/1920	0.61	0/2598
3	Q	0.36	0/1920	0.62	0/2598
4	D	0.36	0/1887	0.63	0/2541
4	R	0.38	0/1887	0.64	0/2541
5	E	0.36	0/1823	0.59	0/2463
5	S	0.37	0/1823	0.59	0/2463
6	F	0.39	0/1937	0.61	0/2614
6	T	0.39	0/1937	0.61	0/2614
7	G	0.39	0/1959	0.62	0/2652
7	U	0.38	0/1959	0.62	0/2652
8	H	0.39	0/1716	0.66	0/2326
8	V	0.38	0/1716	0.66	0/2326
9	I	0.40	0/1611	0.65	0/2174
9	W	0.42	0/1611	0.65	0/2174
10	J	0.42	0/1613	0.64	0/2173
10	X	0.42	0/1613	0.64	0/2173
11	K	0.41	0/1681	0.65	0/2274
11	Y	0.42	0/1681	0.65	0/2274
12	L	0.41	0/1795	0.66	0/2420
12	Z	0.41	0/1795	0.66	0/2420
13	M	0.40	0/1855	0.67	2/2514 (0.1%)
13	a	0.40	0/1855	0.67	2/2514 (0.1%)
14	N	0.41	0/1541	0.64	0/2087
14	b	0.40	0/1541	0.64	0/2087
15	c	1.04	0/6	0.80	0/7
15	d	1.14	0/6	1.13	0/7
15	e	1.08	0/6	0.74	0/7
15	f	0.86	0/6	1.12	0/7

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
All	All	0.39	0/50474	0.64	4/68220 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
12	L	0	1
12	Z	0	1
All	All	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
13	a	110	LEU	N-CA-C	-5.98	94.85	111.00
13	M	110	LEU	N-CA-C	-5.75	95.47	111.00
13	a	113	ALA	N-CA-C	-5.25	96.83	111.00
13	M	113	ALA	N-CA-C	-5.02	97.44	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
12	L	174	TYR	Sidechain
12	Z	174	TYR	Sidechain

5.2 Too-close contacts ⓘ

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/250 (99%)	226 (91%)	18 (7%)	4 (2%)	9	40
1	O	248/250 (99%)	224 (90%)	20 (8%)	4 (2%)	9	40
2	B	242/244 (99%)	213 (88%)	21 (9%)	8 (3%)	4	21
2	P	242/244 (99%)	212 (88%)	23 (10%)	7 (3%)	4	24
3	C	239/241 (99%)	214 (90%)	20 (8%)	5 (2%)	7	33
3	Q	239/241 (99%)	215 (90%)	19 (8%)	5 (2%)	7	33
4	D	240/242 (99%)	212 (88%)	19 (8%)	9 (4%)	3	18
4	R	240/242 (99%)	212 (88%)	19 (8%)	9 (4%)	3	18
5	E	231/233 (99%)	202 (87%)	21 (9%)	8 (4%)	3	20
5	S	231/233 (99%)	202 (87%)	22 (10%)	7 (3%)	4	24
6	F	242/244 (99%)	224 (93%)	16 (7%)	2 (1%)	19	57
6	T	242/244 (99%)	223 (92%)	17 (7%)	2 (1%)	19	57
7	G	241/243 (99%)	219 (91%)	19 (8%)	3 (1%)	13	48
7	U	241/243 (99%)	218 (90%)	20 (8%)	3 (1%)	13	48
8	H	220/222 (99%)	197 (90%)	20 (9%)	3 (1%)	11	43
8	V	220/222 (99%)	195 (89%)	22 (10%)	3 (1%)	11	43
9	I	202/204 (99%)	185 (92%)	14 (7%)	3 (2%)	10	42
9	W	202/204 (99%)	185 (92%)	15 (7%)	2 (1%)	15	53
10	J	196/198 (99%)	183 (93%)	10 (5%)	3 (2%)	10	42
10	X	196/198 (99%)	182 (93%)	11 (6%)	3 (2%)	10	42
11	K	210/212 (99%)	199 (95%)	11 (5%)	0	100	100
11	Y	210/212 (99%)	199 (95%)	11 (5%)	0	100	100
12	L	220/222 (99%)	199 (90%)	16 (7%)	5 (2%)	6	30
12	Z	220/222 (99%)	202 (92%)	14 (6%)	4 (2%)	8	37
13	M	231/233 (99%)	206 (89%)	24 (10%)	1 (0%)	34	72
13	a	231/233 (99%)	206 (89%)	24 (10%)	1 (0%)	34	72
14	N	194/196 (99%)	183 (94%)	10 (5%)	1 (0%)	29	68
14	b	194/196 (99%)	183 (94%)	9 (5%)	2 (1%)	15	53
15	c	1/4 (25%)	1 (100%)	0	0	100	100
15	d	1/4 (25%)	1 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
15	e	1/4 (25%)	1 (100%)	0	0	100	100
15	f	1/4 (25%)	1 (100%)	0	0	100	100
All	All	6316/6384 (99%)	5724 (91%)	485 (8%)	107 (2%)	9	39

5 of 107 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	51	VAL
2	B	62	THR
2	B	221	ASP
3	C	52	LEU
4	D	122	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	203 (97%)	6 (3%)	42	76
1	O	209/209 (100%)	203 (97%)	6 (3%)	42	76
2	B	203/203 (100%)	193 (95%)	10 (5%)	25	61
2	P	203/203 (100%)	193 (95%)	10 (5%)	25	61
3	C	213/213 (100%)	205 (96%)	8 (4%)	33	69
3	Q	213/213 (100%)	205 (96%)	8 (4%)	33	69
4	D	198/198 (100%)	193 (98%)	5 (2%)	47	79
4	R	198/198 (100%)	193 (98%)	5 (2%)	47	79
5	E	192/192 (100%)	176 (92%)	16 (8%)	11	39
5	S	192/192 (100%)	176 (92%)	16 (8%)	11	39
6	F	201/201 (100%)	186 (92%)	15 (8%)	13	43
6	T	201/201 (100%)	184 (92%)	17 (8%)	10	38
7	G	207/207 (100%)	196 (95%)	11 (5%)	22	58
7	U	207/207 (100%)	196 (95%)	11 (5%)	22	58

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
8	H	181/181 (100%)	172 (95%)	9 (5%)	24	60
8	V	181/181 (100%)	173 (96%)	8 (4%)	28	65
9	I	172/172 (100%)	168 (98%)	4 (2%)	50	80
9	W	172/172 (100%)	167 (97%)	5 (3%)	42	76
10	J	175/175 (100%)	172 (98%)	3 (2%)	60	85
10	X	175/175 (100%)	172 (98%)	3 (2%)	60	85
11	K	169/169 (100%)	162 (96%)	7 (4%)	30	67
11	Y	169/169 (100%)	162 (96%)	7 (4%)	30	67
12	L	185/185 (100%)	173 (94%)	12 (6%)	17	50
12	Z	185/185 (100%)	172 (93%)	13 (7%)	15	47
13	M	199/199 (100%)	186 (94%)	13 (6%)	17	50
13	a	199/199 (100%)	188 (94%)	11 (6%)	21	57
14	N	162/162 (100%)	155 (96%)	7 (4%)	29	66
14	b	162/162 (100%)	155 (96%)	7 (4%)	29	66
15	c	1/1 (100%)	1 (100%)	0	100	100
15	d	1/1 (100%)	1 (100%)	0	100	100
15	e	1/1 (100%)	1 (100%)	0	100	100
15	f	1/1 (100%)	1 (100%)	0	100	100
All	All	5336/5336 (100%)	5083 (95%)	253 (5%)	26	63

5 of 253 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
13	M	161	ARG
2	P	220	ASN
12	Z	165	ASN
13	M	206	LEU
1	O	122	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 200 such sidechains are listed below:

Mol	Chain	Res	Type
13	M	108	ASN
3	Q	17	GLN
12	Z	165	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
13	M	194	ASN
2	P	20	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
15	LYO	d	3	15	7,9,10	0.87	0	6,10,12	0.54	0
15	OW6	f	4	11,15	6,6,7	1.12	0	5,6,8	0.65	0
15	OW6	d	4	11,15	6,6,7	1.10	0	5,6,8	0.69	0
15	OW6	e	4	8,15	6,6,7	1.08	0	5,6,8	0.91	0
15	OW6	c	4	8,15	6,6,7	1.45	1 (16%)	5,6,8	0.86	0
15	LYO	c	3	15	7,9,10	0.97	0	6,10,12	0.56	0
15	LYO	e	3	15	7,9,10	0.92	0	6,10,12	0.45	0
15	LYO	f	3	15	7,9,10	0.95	0	6,10,12	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
15	LYO	d	3	15	-	3/8/9/11	-
15	OW6	f	4	11,15	-	1/3/4/5	-
15	OW6	d	4	11,15	-	1/3/4/5	-
15	OW6	e	4	8,15	-	1/3/4/5	-
15	OW6	c	4	8,15	-	1/3/4/5	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
15	LYO	c	3	15	-	4/8/9/11	-
15	LYO	e	3	15	-	4/8/9/11	-
15	LYO	f	3	15	-	3/8/9/11	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
15	c	4	0W6	C17-C15	2.96	1.57	1.53

There are no bond angle outliers.

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
15	d	3	LYO	N-CA-CB-CG
15	d	3	LYO	C-CA-CB-CG
15	d	3	LYO	CG-CD-CE-NZ
15	e	4	0W6	C15-C17-C18-C19
15	c	3	LYO	N-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/250 (100%)	-0.43	2 (0%) 86 65	61, 83, 116, 141	0
1	O	250/250 (100%)	-0.54	4 (1%) 72 44	61, 83, 117, 141	0
2	B	244/244 (100%)	-0.36	6 (2%) 57 29	61, 83, 131, 151	0
2	P	244/244 (100%)	-0.37	6 (2%) 57 29	61, 83, 131, 151	0
3	C	241/241 (100%)	-0.38	2 (0%) 86 65	63, 89, 135, 162	0
3	Q	241/241 (100%)	-0.32	4 (1%) 70 41	65, 91, 136, 162	0
4	D	242/242 (100%)	-0.30	6 (2%) 57 29	61, 90, 125, 155	0
4	R	242/242 (100%)	-0.29	5 (2%) 63 34	63, 91, 126, 155	0
5	E	233/233 (100%)	-0.34	4 (1%) 70 41	67, 95, 125, 145	0
5	S	233/233 (100%)	-0.31	3 (1%) 77 51	68, 95, 125, 145	0
6	F	244/244 (100%)	-0.37	3 (1%) 79 54	64, 87, 124, 141	0
6	T	244/244 (100%)	-0.29	4 (1%) 72 44	65, 87, 125, 140	0
7	G	243/243 (100%)	-0.48	2 (0%) 86 65	62, 84, 116, 152	0
7	U	243/243 (100%)	-0.47	2 (0%) 86 65	63, 83, 115, 153	0
8	H	222/222 (100%)	-0.67	0 100 100	59, 75, 98, 140	0
8	V	222/222 (100%)	-0.67	1 (0%) 91 75	61, 74, 98, 142	0
9	I	204/204 (100%)	-0.71	1 (0%) 91 75	57, 75, 99, 117	0
9	W	204/204 (100%)	-0.68	0 100 100	58, 75, 99, 116	0
10	J	198/198 (100%)	-0.58	3 (1%) 73 46	56, 76, 99, 161	0
10	X	198/198 (100%)	-0.60	3 (1%) 73 46	57, 77, 98, 162	0
11	K	212/212 (100%)	-0.63	1 (0%) 91 75	55, 75, 99, 115	0
11	Y	212/212 (100%)	-0.63	1 (0%) 91 75	56, 75, 101, 117	0
12	L	222/222 (100%)	-0.65	0 100 100	59, 77, 105, 129	0
12	Z	222/222 (100%)	-0.63	0 100 100	59, 77, 104, 129	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
13	M	233/233 (100%)	-0.63	1 (0%) 92 79	59, 77, 99, 111	0
13	a	233/233 (100%)	-0.62	0 100 100	58, 77, 99, 110	0
14	N	196/196 (100%)	-0.65	0 100 100	58, 74, 100, 115	0
14	b	196/196 (100%)	-0.61	0 100 100	58, 74, 101, 114	0
15	c	1/4 (25%)	-0.79	0 100 100	75, 75, 75, 75	0
15	d	1/4 (25%)	-0.65	0 100 100	63, 63, 63, 63	0
15	e	1/4 (25%)	-0.57	0 100 100	75, 75, 75, 75	0
15	f	1/4 (25%)	-1.05	0 100 100	66, 66, 66, 66	0
All	All	6372/6384 (99%)	-0.50	64 (1%) 82 59	55, 81, 120, 162	0

The worst 5 of 64 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	118	GLY	6.2
7	U	243	ASP	6.1
4	R	121	GLY	6.0
4	D	121	GLY	5.6
2	B	220	ASN	5.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
15	LYO	e	3	10/11	0.93	0.18	73,75,78,79	0
15	0W6	c	4	7/8	0.95	0.14	70,72,76,77	0
15	0W6	e	4	7/8	0.95	0.15	71,73,77,77	0
15	0W6	d	4	7/8	0.96	0.13	67,67,72,72	0
15	LYO	c	3	10/11	0.96	0.12	72,74,78,78	0
15	LYO	d	3	10/11	0.96	0.13	64,68,71,71	0
15	0W6	f	4	7/8	0.97	0.12	67,68,72,73	0
15	LYO	f	3	10/11	0.97	0.13	64,67,71,72	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.