



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2020 – 08:47 pm BST

PDB ID : 3G05  
Title : Crystal structure of N-terminal domain (2-550) of E.coli MnmG  
Authors : Shi, R.; Matte, A.; Cygler, M.; Montreal-Kingston Bacterial Structural Genomics Initiative (BSGI)  
Deposited on : 2009-01-27  
Resolution : 3.49 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)
RSRZ outliers	127900	1559 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	576	<div> <div>5%</div> <div>25%</div> <div>48%</div> <div>16%</div> <div>•</div> <div>9%</div> </div>
1	B	576	<div> <div>8%</div> <div>21%</div> <div>48%</div> <div>19%</div> <div>•</div> <div>9%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8156 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called tRNA uridine 5-carboxymethylaminomethyl modification enzyme mnmG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	527	Total	C	N	O	S	0	0	0
			4073	2551	728	776	18			
1	B	524	Total	C	N	O	S	0	0	0
			4063	2540	730	775	18			

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-27	MET	-	EXPRESSION TAG	UNP Q8XAY0
A	-26	GLY	-	EXPRESSION TAG	UNP Q8XAY0
A	-25	SER	-	EXPRESSION TAG	UNP Q8XAY0
A	-24	SER	-	EXPRESSION TAG	UNP Q8XAY0
A	-23	HIS	-	EXPRESSION TAG	UNP Q8XAY0
A	-22	HIS	-	EXPRESSION TAG	UNP Q8XAY0
A	-21	HIS	-	EXPRESSION TAG	UNP Q8XAY0
A	-20	HIS	-	EXPRESSION TAG	UNP Q8XAY0
A	-19	HIS	-	EXPRESSION TAG	UNP Q8XAY0
A	-18	HIS	-	EXPRESSION TAG	UNP Q8XAY0
A	-17	HIS	-	EXPRESSION TAG	UNP Q8XAY0
A	-16	HIS	-	EXPRESSION TAG	UNP Q8XAY0
A	-15	ASP	-	EXPRESSION TAG	UNP Q8XAY0
A	-14	TYR	-	EXPRESSION TAG	UNP Q8XAY0
A	-13	ASP	-	EXPRESSION TAG	UNP Q8XAY0
A	-12	ILE	-	EXPRESSION TAG	UNP Q8XAY0
A	-11	PRO	-	EXPRESSION TAG	UNP Q8XAY0
A	-10	THR	-	EXPRESSION TAG	UNP Q8XAY0
A	-9	THR	-	EXPRESSION TAG	UNP Q8XAY0
A	-8	GLU	-	EXPRESSION TAG	UNP Q8XAY0
A	-7	ASN	-	EXPRESSION TAG	UNP Q8XAY0
A	-6	LEU	-	EXPRESSION TAG	UNP Q8XAY0
A	-5	TYR	-	EXPRESSION TAG	UNP Q8XAY0
A	-4	PHE	-	EXPRESSION TAG	UNP Q8XAY0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLN	-	EXPRESSION TAG	UNP Q8XAY0
A	-2	GLY	-	EXPRESSION TAG	UNP Q8XAY0
A	-1	SER	-	EXPRESSION TAG	UNP Q8XAY0
B	-27	MET	-	EXPRESSION TAG	UNP Q8XAY0
B	-26	GLY	-	EXPRESSION TAG	UNP Q8XAY0
B	-25	SER	-	EXPRESSION TAG	UNP Q8XAY0
B	-24	SER	-	EXPRESSION TAG	UNP Q8XAY0
B	-23	HIS	-	EXPRESSION TAG	UNP Q8XAY0
B	-22	HIS	-	EXPRESSION TAG	UNP Q8XAY0
B	-21	HIS	-	EXPRESSION TAG	UNP Q8XAY0
B	-20	HIS	-	EXPRESSION TAG	UNP Q8XAY0
B	-19	HIS	-	EXPRESSION TAG	UNP Q8XAY0
B	-18	HIS	-	EXPRESSION TAG	UNP Q8XAY0
B	-17	HIS	-	EXPRESSION TAG	UNP Q8XAY0
B	-16	HIS	-	EXPRESSION TAG	UNP Q8XAY0
B	-15	ASP	-	EXPRESSION TAG	UNP Q8XAY0
B	-14	TYR	-	EXPRESSION TAG	UNP Q8XAY0
B	-13	ASP	-	EXPRESSION TAG	UNP Q8XAY0
B	-12	ILE	-	EXPRESSION TAG	UNP Q8XAY0
B	-11	PRO	-	EXPRESSION TAG	UNP Q8XAY0
B	-10	THR	-	EXPRESSION TAG	UNP Q8XAY0
B	-9	THR	-	EXPRESSION TAG	UNP Q8XAY0
B	-8	GLU	-	EXPRESSION TAG	UNP Q8XAY0
B	-7	ASN	-	EXPRESSION TAG	UNP Q8XAY0
B	-6	LEU	-	EXPRESSION TAG	UNP Q8XAY0
B	-5	TYR	-	EXPRESSION TAG	UNP Q8XAY0
B	-4	PHE	-	EXPRESSION TAG	UNP Q8XAY0
B	-3	GLN	-	EXPRESSION TAG	UNP Q8XAY0
B	-2	GLY	-	EXPRESSION TAG	UNP Q8XAY0
B	-1	SER	-	EXPRESSION TAG	UNP Q8XAY0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: tRNA uridine 5-carboxymethylaminomethyl modification enzyme mnmG



- Molecule 1: tRNA uridine 5-carboxymethylaminomethyl modification enzyme mnmG



D535	L410	F347	F287	V226	L103	U38	MET
E536	G411	F348	A288	F227	Y104	I38	GLY
Q537	V412	D349	D289	S228	Q106	D40	SER
A538	L413	P350	E290	F229	LEU		SER
A539	V414	R351	Q291	M230	A107	Q44	HIS
E540	D415		D292	G231	ASP	M45	HIS
Q541	D416		E293	N232	TYR	S46	HIS
V542	L417		Q294	A233	ASN	C47	HIS
E543			T295	S234	GLY	I110	HIS
I544	L420		F296	Q235	GLY	N48	HIS
Q545	G421		L297	H236	ARG	P49	HIS
V546	T422		E298	P237	ALA	A50	HIS
K547	K423		P299	Q238	GLY	I114	HIS
Y548	E424		E300	Q239	ASP	Q115	ASP
E549	P425		G301	V240	PRO	P116	TYR
G550	Y426		L302	P241	PRO	M117	ASP
	R427		S303	C242	S130	K56	ILE
	M428		F304	V243	I181	G57	PRO
	F429		N305	T244	P182	H58	THR
	T430		E306	T245	Q121	L59	THR
	S431		T307	H246	Q122	V60	GLU
	R432		Y308	T247	S184	K61	ASN
	A433		P309	N248	R185	E62	LEU
	E434		N310	E249	R186	V63	TYR
	Y435		G311	K250	L187	D64	PHE
	R436		E312	T251	E188	A65	Q-3
	L437		S313	H252	L189	L66	G-2
	M438		T314	D253	P190	G67	S-1
	L439		S315	V254	P191	G68	F2
	R440		L316	T255	L192	L69	Y3
	E441		P317	R256	R193	M70	P4
	D442		F318		V194	K71	D5
	N443		D319	L259	G195	K72	P6
	A444		V320	D260	R196	A73	F7
	D445		Q321	R261	L197	I74	D8
	L446		N322	S262	K198	D75	V9
	R447		Q323	PRO	T199	Q76	I10
	L448		T324	MET	G200	A77	I11
	T449		V325	TYR	V139	G78	I12
	E450		R326	ALA	T140	I79	G13
	I451		S327	GLY	Q141	Q80	G14
	G452		N328	VAL	L144	F81	G15
	R453		Q329	ILE	K145	R82	T19
	E454		G330	GLU	F146	I83	
	L455		N331	GLY	R147	I84	
	G456		E332	VAL	A148	N85	
	L457		N333	GLY	K149		
	V458		A334	PRO	A150	G89	M23
	D459		K335	ARG	V151	P90	A24
	D460		I336	TVR	V152	A91	A25
	E461		V337	CYS	V153	V92	A26
	W401		R338	P278	T154	R93	M28
	A402		P339	T279	Y155	A94	G29
	P403		G340	I280	Q217	T95	Q30
	A404		Y341	E281	T157	R96	Q31
	R405		A342	D282	H219	A97	T32
	S406		T343	K283	L159	Q98	I33
	Q407		E344	V284	D160	A99	I34
	A408		Y345	W285	G161	D100	L35
	Y409		D346	R286	K162	R101	T36
					I163	V102	H37

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	144.59Å 144.59Å 271.02Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 3.49 49.74 – 3.49	Depositor EDS
% Data completeness (in resolution range)	99.7 (50.00-3.49) 99.7 (49.74-3.49)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.54 (at 3.48Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.227 , 0.265 0.224 , 0.260	Depositor DCC
$R_{free}$ test set	2137 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	91.0	Xtriage
Anisotropy	0.300	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 83.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	8156	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.84	1/4146 (0.0%)	1.15	20/5615 (0.4%)
1	B	0.75	0/4134	1.11	15/5596 (0.3%)
All	All	0.80	1/8280 (0.0%)	1.13	35/11211 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	7
1	B	0	4
All	All	1	11

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	505	ARG	CZ-NH2	24.30	1.64	1.33

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	101	ARG	NE-CZ-NH1	-11.62	114.49	120.30
1	B	290	ARG	N-CA-C	-10.74	82.01	111.00
1	A	505	ARG	NE-CZ-NH1	9.52	125.06	120.30
1	B	100	ASP	CB-CG-OD1	7.85	125.36	118.30
1	A	505	ARG	NH1-CZ-NH2	-7.68	110.95	119.40

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	291	ASN	CA

5 of 11 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	-1	SER	Mainchain,Peptide
1	A	159	LEU	Peptide
1	A	261	ARG	Peptide
1	A	287	PHE	Peptide
1	A	288	ALA	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4073	0	4026	563	0
1	B	4063	0	4029	581	0
2	A	10	0	0	1	0
2	B	10	0	0	0	0
All	All	8156	0	8055	1141	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 70.

The worst 5 of 1141 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:289:ASP:CA	1:A:291:ASN:HA	1.56	1.32
1:B:180:SER:O	1:B:181:ILE:HG22	1.24	1.31
1:B:154:THR:O	1:B:155:VAL:CG1	1.79	1.29
1:A:285:MET:CE	1:A:285:MET:HA	1.59	1.26
1:A:289:ASP:C	1:A:291:ASN:HA	1.56	1.26

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	521/576 (90%)	383 (74%)	88 (17%)	50 (10%)	0	8
1	B	518/576 (90%)	363 (70%)	98 (19%)	57 (11%)	0	6
All	All	1039/1152 (90%)	746 (72%)	186 (18%)	107 (10%)	0	7

5 of 107 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	15	GLY
1	A	142	MET
1	A	160	ASP
1	A	167	LEU
1	A	212	PHE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	426/474 (90%)	338 (79%)	88 (21%)	1	6
1	B	427/474 (90%)	320 (75%)	107 (25%)	0	3
All	All	853/948 (90%)	658 (77%)	195 (23%)	1	4

5 of 195 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	-3	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	114	ASN
1	B	471	GLU
1	B	9	VAL
1	B	46	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 37 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	541	GLN
1	B	44	GLN
1	B	383	GLN
1	B	30	GLN
1	B	31	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	551	-	4,4,4	0.17	0	6,6,6	0.52	0
2	SO4	A	552	-	4,4,4	0.17	0	6,6,6	0.59	0
2	SO4	B	1	-	4,4,4	0.21	0	6,6,6	0.54	0
2	SO4	B	551	-	4,4,4	0.19	0	6,6,6	0.33	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	551	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	527/576 (91%)	0.33	27 (5%)	28 25	53, 73, 123, 136	0
1	B	524/576 (90%)	0.53	46 (8%)	10 10	55, 79, 130, 137	0
All	All	1051/1152 (91%)	0.43	73 (6%)	16 16	53, 75, 128, 137	0

The worst 5 of 73 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	507	ALA	5.8
1	A	503	LEU	5.8
1	B	289	ASP	5.5
1	A	485	VAL	5.5
1	B	164	HIS	5.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	552	5/5	0.93	0.20	79,79,79,80	0
2	SO4	B	551	5/5	0.93	0.20	85,85,85,86	0
2	SO4	B	1	5/5	0.96	0.25	77,78,78,78	0
2	SO4	A	551	5/5	0.98	0.17	71,72,72,72	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.