



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 09:27 pm BST

PDB ID : 3G2C
Title : Mth0212 in complex with a short ssDNA (CGTA)
Authors : Lakomek, K.; Dickmanns, A.; Ficner, R.
Deposited on : 2009-01-31
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	265	<div> <div>4%</div> <div> <div></div> <div>81%</div> <div>12%</div> <div>• •</div> </div> </div>
1	B	265	<div> <div>8%</div> <div> <div></div> <div>77%</div> <div>16%</div> <div>• •</div> </div> </div>
2	I	4	<div> <div>25%</div> <div>75%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	270	-	-	-	X
4	MG	A	271	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4586 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Exodeoxyribonuclease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	1	0
			2146	1379	374	385	8			
1	B	255	Total	C	N	O	S	0	1	0
			2130	1368	372	382	8			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	THR	ENGINEERED MUTATION	UNP O26314
A	258	LEU	-	EXPRESSION TAG	UNP O26314
A	259	GLU	-	EXPRESSION TAG	UNP O26314
A	260	HIS	-	EXPRESSION TAG	UNP O26314
A	261	HIS	-	EXPRESSION TAG	UNP O26314
A	262	HIS	-	EXPRESSION TAG	UNP O26314
A	263	HIS	-	EXPRESSION TAG	UNP O26314
A	264	HIS	-	EXPRESSION TAG	UNP O26314
A	265	HIS	-	EXPRESSION TAG	UNP O26314
B	2	ALA	THR	ENGINEERED MUTATION	UNP O26314
B	258	LEU	-	EXPRESSION TAG	UNP O26314
B	259	GLU	-	EXPRESSION TAG	UNP O26314
B	260	HIS	-	EXPRESSION TAG	UNP O26314
B	261	HIS	-	EXPRESSION TAG	UNP O26314
B	262	HIS	-	EXPRESSION TAG	UNP O26314
B	263	HIS	-	EXPRESSION TAG	UNP O26314
B	264	HIS	-	EXPRESSION TAG	UNP O26314
B	265	HIS	-	EXPRESSION TAG	UNP O26314

- Molecule 2 is a DNA chain called 5'-D(P*CP*GP*TP*A)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	4	Total	C	N	O	P	0	0	0
			83	39	15	25	4			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	I	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		
5	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	107	Total	O	0	0
			107	107		
6	B	59	Total	O	0	0
			59	59		
6	I	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Exodeoxyribonuclease



• Molecule 1: Exodeoxyribonuclease



• Molecule 2: 5'-D(P*CP*GP*TP*A)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	80.33Å 80.33Å 79.61Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.27 – 2.30 28.27 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (28.27-2.30) 99.7 (28.27-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.77 (at 2.31Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.204 , 0.263 0.203 , 0.262	Depositor DCC
R_{free} test set	1295 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	36.9	Xtriage
Anisotropy	0.213	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.009 for -h,-k,l 0.039 for h,-h-k,-l 0.019 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4586	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/2207	0.79	2/2976 (0.1%)
1	B	0.60	0/2188	0.68	0/2950
2	I	1.69	1/92 (1.1%)	3.69	14/138 (10.1%)
All	All	0.68	1/4487 (0.0%)	0.92	16/6064 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	1	DC	OP3-P	-9.94	1.49	1.61

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	2	DG	O4'-C1'-N9	18.83	121.18	108.00
2	I	2	DG	C1'-O4'-C4'	-14.39	95.71	110.10
2	I	3	DT	P-O3'-C3'	11.76	133.81	119.70
2	I	3	DT	C1'-O4'-C4'	-10.88	99.22	110.10
2	I	1	DC	P-O3'-C3'	10.81	132.67	119.70
2	I	4	DA	C1'-O4'-C4'	-9.10	101.00	110.10
2	I	2	DG	C3'-C2'-C1'	-9.04	91.66	102.50
2	I	1	DC	O4'-C1'-N1	8.55	113.99	108.00
2	I	1	DC	C1'-O4'-C4'	-8.27	101.83	110.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	3	DT	O4'-C1'-N1	7.95	113.56	108.00
2	I	4	DA	O4'-C1'-N9	7.76	113.44	108.00
2	I	2	DG	P-O3'-C3'	7.18	128.32	119.70
2	I	2	DG	N9-C1'-C2'	5.41	122.88	112.60
1	A	57	ARG	NE-CZ-NH1	-5.40	117.60	120.30
2	I	4	DA	O4'-C1'-C2'	-5.24	101.71	105.90
1	A	34	LEU	CB-CG-CD2	5.04	119.57	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	210	THR	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2146	0	2090	29	0
1	B	2130	0	2069	22	0
2	I	83	0	46	3	0
3	A	36	0	48	2	0
3	I	6	0	8	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	10	0	0	0	0
5	B	5	0	0	0	0
6	A	107	0	0	4	0
6	B	59	0	0	2	0
6	I	2	0	0	0	0
All	All	4586	0	4261	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:90:ARG:HD3	6:B:303:HOH:O	1.60	0.98
1:B:89:GLU:O	1:B:93:THR:HG23	1.90	0.72
1:A:17:VAL:HG13	1:A:22:PHE:HB2	1.72	0.71
1:B:207:SER:O	1:B:212:ALA:HB3	1.91	0.70
1:B:29:GLU:O	1:B:31:PRO:HD3	1.94	0.68
1:A:165:LYS:H	1:A:165:LYS:HD3	1.65	0.62
1:A:159:ILE:HA	3:A:270:GOL:H2	1.83	0.60
1:A:7:ILE:HD12	1:A:31:PRO:HB3	1.83	0.60
1:A:149:CYS:HB3	1:A:224:PHE:CE2	2.37	0.60
1:B:192:THR:O	1:B:195:MET:HG3	2.01	0.60
1:A:43:PRO:HA	1:A:46:LEU:HD12	1.84	0.59
1:A:11:VAL:HG22	1:A:17:VAL:HG11	1.86	0.58
1:A:236:ARG:HB2	1:A:254:GLU:HB2	1.85	0.57
1:A:118:SER:HB2	1:A:121:ARG:H	1.69	0.57
1:A:235:LYS:HE3	1:A:256:GLU:HB3	1.85	0.57
1:A:164:PRO:HG3	3:A:270:GOL:O1	2.05	0.56
1:A:149:CYS:HB3	1:A:224:PHE:CD2	2.41	0.55
1:A:44:GLU:OE1	1:A:44:GLU:N	2.26	0.54
1:A:176:VAL:HG22	6:A:328:HOH:O	2.06	0.54
1:B:213:ARG:NH2	1:B:243:VAL:O	2.36	0.54
1:B:166:GLU:H	1:B:166:GLU:CD	2.09	0.54
1:A:43:PRO:HA	1:A:46:LEU:CD1	2.40	0.52
1:B:46:LEU:HD22	1:B:50:LEU:HD23	1.91	0.51
1:A:140:ARG:NH2	6:A:290:HOH:O	2.37	0.50
1:A:48:ARG:HB3	6:A:325:HOH:O	2.13	0.48
1:B:11:VAL:HG22	1:B:17:VAL:HG11	1.95	0.48
1:A:210:THR:HG1	1:A:215:ARG:HE	1.62	0.47
1:A:164:PRO:HG2	1:A:165:LYS:HD3	1.96	0.47
1:A:65:ARG:HG2	1:A:68:TYR:CB	2.45	0.47
1:A:159:ILE:HG23	1:A:194:ARG:NH2	2.30	0.46
1:A:15:ARG:HH11	1:A:15:ARG:HG3	1.80	0.46
2:I:4:DA:O4'	2:I:4:DA:OP2	2.33	0.46
1:A:11:VAL:CG2	1:A:17:VAL:HG11	2.45	0.45
1:B:179:ALA:O	1:B:182:ASP:HB2	2.17	0.45
1:A:159:ILE:HG23	1:A:194:ARG:HH21	1.81	0.44
1:B:136:VAL:HG21	1:B:148:ILE:HD11	1.99	0.44
1:A:209:ARG:C	1:A:211:ARG:N	2.71	0.43
1:B:65:ARG:HD2	1:B:68:TYR:HB2	1.99	0.43
1:B:65:ARG:HD2	1:B:68:TYR:CB	2.49	0.43
1:A:165:LYS:NZ	1:A:166:GLU:OE2	2.52	0.43
1:B:9:TRP:CD2	1:B:250:PRO:HG3	2.54	0.42
1:A:192:THR:HG22	1:A:195:MET:HE1	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:140:ARG:HD3	6:B:286:HOH:O	2.19	0.42
1:B:98:GLN:HB2	1:B:109:ASN:HB3	2.02	0.42
1:B:149:CYS:HB3	1:B:224:PHE:CE2	2.54	0.41
1:A:176:VAL:CG2	6:A:328:HOH:O	2.66	0.41
1:B:65:ARG:HD2	1:B:68:TYR:CG	2.55	0.41
1:A:149:CYS:HA	1:A:223:TYR:O	2.21	0.41
1:B:7:ILE:HG21	1:B:25:TRP:HH2	1.85	0.41
2:I:2:DG:C2	2:I:3:DT:H2'	2.55	0.41
1:B:163:ARG:HB3	1:B:166:GLU:HG2	2.03	0.41
1:B:205:TRP:HD1	1:B:248:HIS:CD2	2.39	0.41
1:B:211:ARG:HB2	1:B:215:ARG:HE	1.86	0.40
2:I:3:DT:O2	2:I:4:DA:H4'	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/265 (97%)	243 (95%)	11 (4%)	2 (1%)	19	23
1	B	253/265 (96%)	238 (94%)	12 (5%)	3 (1%)	13	14
All	All	509/530 (96%)	481 (94%)	23 (4%)	5 (1%)	15	17

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	211	ARG
1	B	20	LYS
1	B	210	THR
1	A	210	THR
1	B	103[A]	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/235 (97%)	212 (93%)	16 (7%)	15	19
1	B	226/235 (96%)	207 (92%)	19 (8%)	11	13
All	All	454/470 (97%)	419 (92%)	35 (8%)	13	16

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	VAL
1	A	15	ARG
1	A	17	VAL
1	A	27	MET
1	A	34	LEU
1	A	48	ARG
1	A	49	LYS
1	A	80	SER
1	A	89	GLU
1	A	117	MET
1	A	133	LEU
1	A	165	LYS
1	A	176	VAL
1	A	186	GLU
1	A	210	THR
1	A	256	GLU
1	B	3	VAL
1	B	11	VAL
1	B	17	VAL
1	B	20	LYS
1	B	22	PHE
1	B	29	GLU
1	B	30	LYS
1	B	34	LEU
1	B	48	ARG
1	B	49	LYS
1	B	54	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	89	GLU
1	B	90	ARG
1	B	129	TYR
1	B	133	LEU
1	B	140	ARG
1	B	169	ASN
1	B	194	ARG
1	B	195	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	167	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	270	-	5,5,5	0.38	0	5,5,5	0.65	0
3	GOL	A	267	-	5,5,5	0.40	0	5,5,5	0.73	0
3	GOL	A	274	-	5,5,5	0.37	0	5,5,5	0.26	0
5	PO4	A	272	4	4,4,4	0.63	0	6,6,6	0.73	0
3	GOL	A	269	-	5,5,5	0.36	0	5,5,5	0.26	0
3	GOL	I	5	-	5,5,5	0.42	0	5,5,5	0.33	0
3	GOL	A	266	-	5,5,5	0.34	0	5,5,5	0.62	0
3	GOL	A	268	-	5,5,5	0.34	0	5,5,5	0.38	0
5	PO4	B	267	4	4,4,4	0.82	0	6,6,6	0.50	0
5	PO4	A	273	-	4,4,4	0.75	0	6,6,6	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	270	-	-	0/4/4/4	-
3	GOL	A	267	-	-	1/4/4/4	-
3	GOL	A	274	-	-	2/4/4/4	-
3	GOL	A	269	-	-	4/4/4/4	-
3	GOL	I	5	-	-	2/4/4/4	-
3	GOL	A	266	-	-	2/4/4/4	-
3	GOL	A	268	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	274	GOL	C1-C2-C3-O3
3	A	274	GOL	O2-C2-C3-O3
3	A	269	GOL	O1-C1-C2-C3
3	I	5	GOL	C1-C2-C3-O3
3	A	266	GOL	O1-C1-C2-C3
3	A	268	GOL	O1-C1-C2-C3
3	A	268	GOL	C1-C2-C3-O3
3	A	266	GOL	O1-C1-C2-O2
3	A	268	GOL	O2-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	269	GOL	O1-C1-C2-O2
3	A	268	GOL	O1-C1-C2-O2
3	I	5	GOL	O2-C2-C3-O3
3	A	269	GOL	C1-C2-C3-O3
3	A	267	GOL	O1-C1-C2-C3
3	A	269	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	270	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/265 (96%)	-0.04	10 (3%) 39 46	18, 30, 49, 57	2 (0%)
1	B	255/265 (96%)	0.30	20 (7%) 13 17	25, 41, 65, 71	2 (0%)
2	I	4/4 (100%)	0.24	0 100 100	55, 58, 62, 70	0
All	All	516/534 (96%)	0.13	30 (5%) 23 29	18, 36, 61, 71	4 (0%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	19	ARG	4.0
1	B	27	MET	3.4
1	B	3	VAL	3.3
1	B	152	PHE	3.0
1	B	21	GLY	2.9
1	B	24	LYS	2.7
1	B	211	ARG	2.6
1	A	146	VAL	2.5
1	B	149	CYS	2.5
1	B	112	PHE	2.5
1	A	2	ALA	2.5
1	A	221	LEU	2.5
1	A	210	THR	2.4
1	B	148	ILE	2.4
1	A	117	MET	2.4
1	B	20	LYS	2.4
1	B	210	THR	2.4
1	B	175	PRO	2.3
1	B	195	MET	2.3
1	A	119	GLU	2.3
1	B	242	ASP	2.3
1	B	8	SER	2.3
1	B	221	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	165	LYS	2.2
1	B	223	TYR	2.2
1	B	196	PHE	2.2
1	A	110	ILE	2.2
1	B	207	SER	2.1
1	A	90	ARG	2.1
1	A	258	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	269	6/6	0.59	0.27	68,70,70,71	0
3	GOL	A	274	6/6	0.63	0.19	62,63,64,64	0
3	GOL	A	270	6/6	0.64	0.42	61,62,62,62	0
4	MG	A	271	1/1	0.77	0.41	49,49,49,49	0
3	GOL	A	267	6/6	0.80	0.16	47,48,50,50	0
5	PO4	A	273	5/5	0.83	0.20	82,82,83,83	0
4	MG	B	266	1/1	0.86	0.29	48,48,48,48	0
3	GOL	A	266	6/6	0.87	0.15	40,43,45,46	0
3	GOL	A	268	6/6	0.89	0.12	43,46,48,50	0
3	GOL	I	5	6/6	0.89	0.15	62,63,64,64	0
5	PO4	A	272	5/5	0.94	0.14	46,46,49,50	0
5	PO4	B	267	5/5	0.96	0.14	47,49,51,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.