



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 07:11 am BST

PDB ID : 3G56  
Title : Structure of the macrolide biosensor protein, MphR(A)  
Authors : Zheng, J.; Sagar, V.; Smolinsky, A.; Bourke, C.; LaRonde-LeBlanc, N.; Cropp, T.A.  
Deposited on : 2009-02-04  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	195	<div> <div>7%</div> <div> <div></div> <div>59%</div> <div>25%</div> <div>7%</div> <div>8%</div> </div> </div>
1	B	195	<div> <div>4%</div> <div> <div></div> <div>53%</div> <div>33%</div> <div>6%</div> <div>8%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	196	-	-	-	X
3	GOL	B	196	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2810 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Regulator of macrolide 2'-phosphotransferase I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	180	Total	C	N	O	S	0	0	0
			1371	878	240	246	7			
1	B	179	Total	C	N	O	S	0	0	0
			1354	867	242	241	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	EXPRESSION TAG	UNP Q9EVJ6
B	0	GLY	-	EXPRESSION TAG	UNP Q9EVJ6

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	27	Total	O	0	0
			27	27		
4	B	36	Total	O	0	0
			36	36		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Regulator of macrolide 2'-phosphotransferase I



- Molecule 1: Regulator of macrolide 2'-phosphotransferase I



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	37.42Å 113.76Å 43.39Å 90.00° 93.87° 90.00°	Depositor
Resolution (Å)	29.27 – 2.10 29.27 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.5 (29.27-2.10) 98.5 (29.27-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.58 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.220 , 0.284 0.226 , 0.294	Depositor DCC
$R_{free}$ test set	1070 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	43.7	Xtriage
Anisotropy	0.261	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 57.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2810	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.37	7/1397 (0.5%)	1.22	6/1903 (0.3%)
1	B	1.46	7/1378 (0.5%)	1.24	8/1877 (0.4%)
All	All	1.41	14/2775 (0.5%)	1.23	14/3780 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	64	GLU	CG-CD	8.79	1.65	1.51
1	B	64	GLU	CB-CG	7.15	1.65	1.52
1	A	169	VAL	CB-CG2	-6.94	1.38	1.52
1	A	69	TYR	CE2-CZ	6.75	1.47	1.38
1	A	69	TYR	CD1-CE1	6.42	1.49	1.39
1	A	69	TYR	CD2-CE2	5.96	1.48	1.39
1	B	77	ALA	CA-CB	5.83	1.64	1.52
1	B	169	VAL	CB-CG1	5.80	1.65	1.52
1	B	148	SER	CA-CB	5.65	1.61	1.52
1	B	174	ALA	CA-CB	5.40	1.63	1.52
1	B	126	VAL	CB-CG1	5.21	1.63	1.52
1	A	183	GLU	CG-CD	5.19	1.59	1.51
1	A	175	ALA	CA-CB	5.04	1.63	1.52
1	A	149	VAL	CA-CB	5.03	1.65	1.54

All (14) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	160	ASP	CB-CG-OD2	-9.56	109.69	118.30
1	A	41	ARG	NE-CZ-NH2	-9.10	115.75	120.30
1	A	144	LEU	CB-CG-CD1	-7.01	99.08	111.00
1	B	51	ARG	NE-CZ-NH1	-6.70	116.95	120.30
1	B	57	ARG	NE-CZ-NH1	6.36	123.48	120.30
1	B	173	ILE	CG1-CB-CG2	-6.25	97.64	111.40
1	A	41	ARG	NE-CZ-NH1	6.11	123.36	120.30
1	A	9	ASP	CB-CG-OD2	5.81	123.53	118.30
1	B	133	ARG	NE-CZ-NH1	-5.77	117.42	120.30
1	B	160	ASP	CB-CG-OD1	5.77	123.49	118.30
1	A	138	ALA	N-CA-C	-5.50	96.16	111.00
1	B	124	ARG	NE-CZ-NH1	-5.46	117.57	120.30
1	B	180	MET	CG-SD-CE	-5.27	91.77	100.20
1	A	51	ARG	NE-CZ-NH1	-5.15	117.72	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	188	GLN	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1371	0	1362	58	0
1	B	1354	0	1357	60	0
2	A	4	0	6	0	0
3	A	6	0	8	0	0
3	B	12	0	16	6	0
4	A	27	0	0	3	0
4	B	36	0	0	2	0
All	All	2810	0	2749	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:6:LEU:CB	1:B:37:VAL:HG22	1.60	1.31
1:B:6:LEU:CB	1:B:37:VAL:CG2	2.32	1.06
1:A:114:GLU:O	1:A:118:LEU:HD12	1.59	1.02
1:A:100:SER:O	1:B:107:TRP:HD1	1.55	0.89
1:B:141:ALA:H	3:B:196:GOL:H2	1.36	0.87
1:A:31:SER:HB2	1:A:41:ARG:HD2	1.56	0.87
1:A:156:GLN:HE21	1:B:147:HIS:HD2	1.28	0.80
1:A:100:SER:O	1:B:107:TRP:CD1	2.36	0.78
1:A:129:GLY:O	1:A:133:ARG:HD3	1.85	0.75
1:A:147:HIS:HD2	1:B:156:GLN:HE21	1.33	0.74
1:A:68:HIS:HB3	4:A:262:HOH:O	1.89	0.71
1:A:151:ALA:O	1:A:155:MET:HG3	1.90	0.69
1:A:25:PRO:HD2	1:A:104:LEU:HD23	1.75	0.69
1:A:156:GLN:HE21	1:B:147:HIS:CD2	2.09	0.68
1:A:9:ASP:HB2	1:A:47:ARG:HH21	1.57	0.68
1:B:52:ASP:O	1:B:56:VAL:HG23	1.93	0.68
1:A:99:PHE:O	1:A:99:PHE:CD2	2.47	0.67
1:B:143:GLU:OE1	3:B:196:GOL:H31	1.93	0.67
1:A:147:HIS:CD2	1:B:156:GLN:HE21	2.14	0.65
1:B:60:GLU:HG3	1:B:118:LEU:HD11	1.81	0.63
1:A:101:VAL:O	1:A:105:ILE:HG13	2.00	0.62
1:A:79:PRO:HG2	1:A:184:HIS:ND1	2.15	0.62
1:A:57:ARG:NH1	1:A:60:GLU:OE1	2.32	0.61
1:A:99:PHE:C	1:A:99:PHE:CD2	2.73	0.61
1:B:6:LEU:CB	1:B:37:VAL:O	2.49	0.61
1:B:161:PRO:O	1:B:162:ASP:HB2	2.01	0.60
1:B:28:PHE:HD2	1:B:51:ARG:HG3	1.68	0.59
1:B:108:TYR:O	1:B:111:GLN:HB2	2.02	0.59
1:B:141:ALA:N	3:B:196:GOL:H2	2.13	0.58
1:A:25:PRO:HD2	1:A:104:LEU:CD2	2.32	0.58
1:B:6:LEU:CB	1:B:37:VAL:HG23	2.31	0.58
1:B:155:MET:HB3	4:B:205:HOH:O	2.04	0.58
1:B:34:ALA:HB1	1:B:39:LEU:O	2.04	0.58
1:B:105:ILE:O	1:B:109:GLU:HG3	2.04	0.57
1:A:73:ILE:CB	1:A:133:ARG:HH21	2.19	0.56
1:B:126:VAL:O	1:B:130:ILE:HG13	2.06	0.55
1:A:159:VAL:O	1:B:116:ARG:NH1	2.39	0.55
1:A:131:ARG:O	1:A:133:ARG:N	2.39	0.55
1:B:39:LEU:C	1:B:39:LEU:HD12	2.27	0.55
1:B:109:GLU:OE1	1:B:122:ARG:NH2	2.41	0.54
1:B:83:TRP:O	1:B:87:GLN:HG3	2.07	0.53
1:B:28:PHE:CD2	1:B:51:ARG:HG3	2.43	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:99:PHE:HD2	1:A:99:PHE:C	2.11	0.53
1:A:131:ARG:O	1:A:132:LYS:C	2.45	0.53
1:A:99:PHE:HD2	1:A:99:PHE:O	1.88	0.53
1:A:66:VAL:O	1:A:68:HIS:N	2.42	0.52
1:B:88:VAL:O	1:B:92:SER:HB3	2.10	0.52
1:B:51:ARG:NH2	1:B:108:TYR:OH	2.43	0.52
1:B:9:ASP:OD1	1:B:47:ARG:NH2	2.38	0.52
1:B:7:LYS:O	1:B:47:ARG:NH1	2.38	0.52
1:B:39:LEU:HD11	1:B:44:LEU:HG	1.93	0.51
1:B:150:ILE:CG1	1:B:173:ILE:HD11	2.40	0.51
1:B:59:MET:HB3	1:B:118:LEU:HD13	1.94	0.50
1:A:155:MET:SD	1:B:155:MET:CG	2.99	0.50
1:A:9:ASP:H	1:A:47:ARG:NH2	2.09	0.49
1:A:68:HIS:CB	4:A:262:HOH:O	2.55	0.49
1:B:150:ILE:HG12	1:B:173:ILE:HD11	1.95	0.49
1:A:107:TRP:CD1	1:B:100:SER:O	2.66	0.48
1:B:108:TYR:HA	1:B:111:GLN:HG2	1.95	0.48
1:A:179:LEU:HB3	1:B:179:LEU:HB3	1.95	0.48
1:A:131:ARG:HA	1:A:134:LEU:HG	1.96	0.48
1:A:28:PHE:HD2	1:A:51:ARG:HG3	1.78	0.48
1:B:22:ARG:HG3	4:B:202:HOH:O	2.14	0.48
1:A:9:ASP:CB	1:A:47:ARG:HH21	2.27	0.48
1:B:128:GLU:OE2	1:B:132:LYS:HE3	2.14	0.47
1:A:134:LEU:HA	1:A:135:PRO:HD2	1.45	0.47
1:A:190:LEU:HD13	1:A:190:LEU:HA	1.66	0.47
1:B:83:TRP:HB2	1:B:177:LEU:HD13	1.95	0.46
1:A:52:ASP:O	1:A:56:VAL:HG23	2.15	0.46
1:A:57:ARG:NH1	1:A:60:GLU:CD	2.69	0.46
1:A:100:SER:HB3	1:B:110:LEU:HD12	1.97	0.46
1:A:131:ARG:O	1:A:134:LEU:N	2.27	0.46
1:A:73:ILE:CB	1:A:133:ARG:NH2	2.79	0.46
1:B:141:ALA:H	3:B:196:GOL:C2	2.19	0.45
1:A:46:GLN:NE2	4:A:208:HOH:O	2.50	0.45
1:A:107:TRP:O	1:A:111:GLN:HG2	2.16	0.45
3:B:195:GOL:H2	3:B:196:GOL:H12	1.99	0.45
1:A:66:VAL:C	1:A:68:HIS:N	2.69	0.44
1:B:20:LEU:O	1:B:20:LEU:HG	2.18	0.44
1:A:185:ASP:O	1:A:186:ASP:O	2.36	0.44
1:A:66:VAL:O	1:A:67:ARG:C	2.55	0.44
1:A:67:ARG:O	1:A:71:ASN:ND2	2.50	0.44
1:B:92:SER:O	1:B:94:ASN:HA	2.18	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:75:ILE:H	1:A:75:ILE:HD12	1.83	0.44
1:B:61:ARG:HD2	1:B:65:GLN:HE21	1.81	0.44
1:A:70:LEU:HD23	1:A:70:LEU:HA	1.87	0.43
1:A:150:ILE:HG13	1:A:173:ILE:HD11	1.99	0.43
1:A:28:PHE:CD2	1:A:51:ARG:HG3	2.53	0.43
1:A:60:GLU:HA	1:A:118:LEU:HD22	2.01	0.43
1:B:12:VAL:HG11	1:B:48:PHE:CE2	2.54	0.43
1:A:82:LEU:O	1:A:83:TRP:C	2.57	0.42
1:A:143:GLU:HG2	1:A:144:LEU:N	2.34	0.42
1:B:37:VAL:HG13	1:B:39:LEU:HG	2.00	0.42
1:B:61:ARG:CD	1:B:65:GLN:HE21	2.33	0.42
1:A:111:GLN:HE22	1:B:100:SER:HB2	1.85	0.42
1:B:143:GLU:CD	3:B:196:GOL:H31	2.40	0.42
1:B:173:ILE:HG21	1:B:173:ILE:HD13	1.61	0.42
1:A:122:ARG:O	1:A:126:VAL:HG23	2.20	0.42
1:B:28:PHE:O	1:B:51:ARG:CD	2.68	0.41
1:B:101:VAL:O	1:B:105:ILE:HG12	2.20	0.41
1:B:143:GLU:HG2	1:B:144:LEU:N	2.36	0.41
1:A:69:TYR:O	1:A:70:LEU:C	2.59	0.41
1:B:28:PHE:O	1:B:51:ARG:HD2	2.20	0.41
1:B:73:ILE:HA	1:B:74:PRO:HD3	1.90	0.41
1:A:185:ASP:O	1:A:186:ASP:C	2.59	0.41
1:B:173:ILE:O	1:B:176:ILE:HG22	2.20	0.40
1:B:63:VAL:O	1:B:67:ARG:HG3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	176/195 (90%)	159 (90%)	8 (4%)	9 (5%)	<b>2</b> <b>0</b>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	175/195 (90%)	164 (94%)	6 (3%)	5 (3%)	4	1
All	All	351/390 (90%)	323 (92%)	14 (4%)	14 (4%)	3	1

All (14) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	9	ASP
1	A	71	ASN
1	B	49	THR
1	B	162	ASP
1	B	185	ASP
1	A	67	ARG
1	A	132	LYS
1	A	137	GLY
1	B	184	HIS
1	A	70	LEU
1	A	99	PHE
1	A	131	ARG
1	A	135	PRO
1	B	186	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	139/161 (86%)	131 (94%)	8 (6%)	20	17
1	B	135/161 (84%)	121 (90%)	14 (10%)	7	4
All	All	274/322 (85%)	252 (92%)	22 (8%)	12	8

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	LEU
1	A	22	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	57	ARG
1	A	98	ASP
1	A	99	PHE
1	A	101	VAL
1	A	114	GLU
1	A	190	LEU
1	B	7	LYS
1	B	8	SER
1	B	11	GLU
1	B	20	LEU
1	B	46	GLN
1	B	51	ARG
1	B	59	MET
1	B	92	SER
1	B	93	MET
1	B	94	ASN
1	B	101	VAL
1	B	105	ILE
1	B	176	ILE
1	B	187	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	147	HIS
1	B	65	GLN
1	B	147	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	B	196	-	5,5,5	0.62	0	5,5,5	0.70	0
3	GOL	A	196	-	5,5,5	0.35	0	5,5,5	0.48	0
2	EDO	A	195	-	3,3,3	0.82	0	2,2,2	0.55	0
3	GOL	B	195	-	5,5,5	0.44	0	5,5,5	1.01	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	196	-	-	2/4/4/4	-
3	GOL	A	196	-	-	4/4/4/4	-
2	EDO	A	195	-	-	1/1/1/1	-
3	GOL	B	195	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	196	GOL	O1-C1-C2-C3
3	A	196	GOL	C1-C2-C3-O3
3	B	196	GOL	C1-C2-C3-O3
3	B	195	GOL	C1-C2-C3-O3
3	B	196	GOL	O2-C2-C3-O3
3	B	195	GOL	O1-C1-C2-O2
3	A	196	GOL	O1-C1-C2-O2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	A	195	EDO	O1-C1-C2-O2
3	A	196	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	196	GOL	6	0
3	B	195	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	180/195 (92%)	0.42	13 (7%) 15 19	27, 52, 71, 75	5 (2%)
1	B	179/195 (91%)	0.22	8 (4%) 33 38	22, 47, 64, 83	4 (2%)
All	All	359/390 (92%)	0.32	21 (5%) 23 28	22, 49, 69, 83	9 (2%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	6	LEU	4.3
1	B	185	ASP	3.6
1	B	186	ASP	3.5
1	A	136	PRO	3.4
1	A	189	LEU	3.4
1	A	99	PHE	3.3
1	A	69	TYR	3.2
1	B	75	ILE	3.2
1	A	73	ILE	3.1
1	A	190	LEU	2.5
1	A	186	ASP	2.5
1	A	37	VAL	2.3
1	A	106	SER	2.3
1	A	98	ASP	2.3
1	A	75	ILE	2.3
1	B	95	THR	2.2
1	B	145	LEU	2.1
1	B	58	MET	2.1
1	B	68	HIS	2.1
1	A	91	ARG	2.0
1	A	71	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	196	6/6	0.63	0.51	106,107,108,108	0
3	GOL	B	195	6/6	0.83	0.25	67,72,73,73	0
3	GOL	B	196	6/6	0.84	0.25	55,62,63,64	0
2	EDO	A	195	4/4	0.86	0.18	78,80,83,83	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.