



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 27, 2020 – 02:12 am BST

PDB ID : 3G78  
Title : Insight into group II intron catalysis from revised crystal structure  
Authors : Wang, J.  
Deposited on : 2009-02-09  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)
RNA backbone	3102	1227 (3.10-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	412	<div> <div>9%</div> <div>42%</div> <div>33%</div> <div>16%</div> <div>6%</div> </div>
2	Z	9	<div> <div>67%</div> <div>44%</div> <div>44%</div> <div>11%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MG	A	413	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MG	A	414	-	-	-	X
3	MG	A	418	-	-	-	X
3	MG	A	424	-	-	-	X
3	MG	A	426	-	-	-	X
3	MG	A	427	-	-	-	X
3	MG	A	448	-	-	-	X
3	MG	A	452	-	-	-	X
3	MG	A	456	-	-	-	X
3	MG	A	457	-	-	-	X
3	MG	A	460	-	-	-	X
4	K	A	472	-	-	-	X
4	K	A	473	-	-	-	X
4	K	A	482	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9057 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called Group II intron.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	389	Total	C	N	O	P	0	0	0
			8363	3734	1554	2687	388			

- Molecule 2 is a RNA chain called Ligated EXON product.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	Z	9	Total	C	N	O	P	0	0	0
			182	83	24	67	8			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	Z	3	Total	Mg	0	0
			3	3		
3	A	48	Total	Mg	0	0
			48	48		

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	26	Total	K	0	0
			26	26		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	432	Total	O	0	0
			432	432		
5	Z	3	Total	O	0	0
			3	3		



- Molecule 1: Group II intron



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.11Å 94.97Å 226.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.29 – 2.80 39.29 – 2.80	Depositor EDS
% Data completeness (in resolution range)	97.2 (39.29-2.80) 97.3 (39.29-2.80)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.03 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.4.0077	Depositor
R, $R_{free}$	0.194 , 0.227 0.234 , 0.271	Depositor DCC
$R_{free}$ test set	2346 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	55.9	Xtriage
Anisotropy	0.251	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 36.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	9057	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: K, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.53	0/9373	1.13	56/14623 (0.4%)
2	Z	0.58	0/201	1.04	0/310
All	All	0.53	0/9574	1.13	56/14933 (0.4%)

There are no bond length outliers.

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	207	A	C3'-C2'-C1'	-11.18	92.56	101.50
1	A	224	A	C3'-C2'-C1'	-10.38	93.20	101.50
1	A	7	C	C4'-C3'-C2'	-9.52	93.08	102.60
1	A	297	A	C3'-C2'-C1'	-8.95	94.34	101.50
1	A	13	U	P-O5'-C5'	-8.86	106.72	120.90

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8363	0	4201	300	0
2	Z	182	0	94	4	0
3	A	48	0	0	0	0
3	Z	3	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	26	0	0	0	0
5	A	432	0	0	1	0
5	Z	3	0	0	0	0
All	All	9057	0	4295	304	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 304 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181:A:C8	1:A:181:A:H5''	1.44	1.50
1:A:207:A:H3'	1:A:208:A:C8	1.49	1.44
1:A:206:A:C2'	1:A:207:A:H5''	1.54	1.37
1:A:2:U:H6	1:A:2:U:C5'	1.38	1.35
1:A:2:U:C6	1:A:2:U:H5'	1.73	1.22

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	389/412 (94%)	117 (30%)	27 (6%)
2	Z	8/9 (88%)	3 (37%)	0
All	All	397/421 (94%)	120 (30%)	27 (6%)

5 of 120 RNA backbone outliers are listed below:



Mol	Chain	Res	Type
1	A	3	G
1	A	7	C
1	A	9	G
1	A	12	A
1	A	13	U

5 of 27 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	208	A
1	A	246	A
1	A	341	C
1	A	223	A
1	A	58	G

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 77 ligands modelled in this entry, 77 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	389/412 (94%)	1.02	38 (9%) <b>7</b> <b>4</b>	2, 12, 68, 84	0
2	Z	9/9 (100%)	3.26	6 (66%) <b>0</b> <b>0</b>	31, 46, 64, 72	0
All	All	398/421 (94%)	1.07	44 (11%) <b>5</b> <b>3</b>	2, 12, 69, 84	0

The worst 5 of 44 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	Z	1	U	8.7
1	A	311	G	7.3
1	A	390	G	5.8
1	A	310	C	5.8
1	A	216	C	5.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	A	427	1/1	-0.19	2.32	108,108,108,108	0
3	MG	A	413	1/1	0.14	1.35	30,30,30,30	0
3	MG	A	424	1/1	0.49	0.62	110,110,110,110	0
4	K	A	484	1/1	0.51	0.25	52,52,52,52	0
4	K	A	470	1/1	0.51	0.19	56,56,56,56	0
3	MG	A	452	1/1	0.51	1.01	32,32,32,32	0
3	MG	A	418	1/1	0.52	0.60	36,36,36,36	0
3	MG	A	451	1/1	0.54	0.35	52,52,52,52	0
3	MG	A	425	1/1	0.58	0.37	34,34,34,34	0
3	MG	A	431	1/1	0.60	0.31	51,51,51,51	0
4	K	A	482	1/1	0.61	0.43	44,44,44,44	0
3	MG	A	455	1/1	0.61	0.29	12,12,12,12	0
3	MG	A	426	1/1	0.65	0.43	44,44,44,44	0
3	MG	A	430	1/1	0.66	0.34	61,61,61,61	0
3	MG	A	445	1/1	0.66	0.30	17,17,17,17	0
3	MG	A	460	1/1	0.68	0.42	95,95,95,95	0
3	MG	Z	11	1/1	0.69	0.24	2,2,2,2	0
3	MG	A	423	1/1	0.70	0.20	30,30,30,30	0
3	MG	A	457	1/1	0.70	0.45	77,77,77,77	0
3	MG	A	456	1/1	0.71	0.60	40,40,40,40	0
3	MG	A	448	1/1	0.72	0.47	16,16,16,16	0
4	K	A	483	1/1	0.74	0.24	58,58,58,58	0
4	K	A	472	1/1	0.77	0.68	73,73,73,73	0
3	MG	A	414	1/1	0.77	0.59	48,48,48,48	0
3	MG	A	421	1/1	0.78	0.35	59,59,59,59	0
4	K	A	473	1/1	0.78	0.41	27,27,27,27	0
3	MG	A	449	1/1	0.79	0.37	53,53,53,53	0
4	K	A	485	1/1	0.80	0.18	60,60,60,60	0
3	MG	A	428	1/1	0.81	0.27	73,73,73,73	0
4	K	A	471	1/1	0.82	0.12	49,49,49,49	0
3	MG	A	422	1/1	0.82	0.30	70,70,70,70	0
3	MG	Z	12	1/1	0.83	0.43	43,43,43,43	0
4	K	A	486	1/1	0.83	0.26	42,42,42,42	0
3	MG	A	416	1/1	0.83	0.38	18,18,18,18	0
3	MG	A	415	1/1	0.84	0.18	42,42,42,42	0
4	K	A	477	1/1	0.85	0.32	17,17,17,17	0
3	MG	A	437	1/1	0.85	0.18	26,26,26,26	0
3	MG	A	454	1/1	0.85	0.10	23,23,23,23	0
3	MG	A	420	1/1	0.86	0.28	62,62,62,62	0
3	MG	A	447	1/1	0.86	0.17	17,17,17,17	0
3	MG	A	429	1/1	0.86	0.07	16,16,16,16	0
4	K	A	474	1/1	0.87	0.16	29,29,29,29	0
3	MG	A	440	1/1	0.87	0.13	22,22,22,22	0

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	A	441	1/1	0.87	0.29	38,38,38,38	0
4	K	A	479	1/1	0.87	0.21	53,53,53,53	0
3	MG	A	439	1/1	0.88	0.17	2,2,2,2	0
3	MG	A	434	1/1	0.89	0.14	12,12,12,12	0
4	K	A	469	1/1	0.89	0.13	52,52,52,52	0
3	MG	A	417	1/1	0.89	0.45	64,64,64,64	0
4	K	A	480	1/1	0.89	0.16	4,4,4,4	0
3	MG	Z	10	1/1	0.90	0.24	10,10,10,10	0
4	K	A	481	1/1	0.90	0.23	54,54,54,54	0
4	K	A	463	1/1	0.91	0.10	36,36,36,36	0
3	MG	A	438	1/1	0.91	0.07	13,13,13,13	0
3	MG	A	450	1/1	0.91	0.31	8,8,8,8	0
4	K	A	467	1/1	0.91	0.14	46,46,46,46	0
3	MG	A	443	1/1	0.91	0.12	2,2,2,2	0
3	MG	A	436	1/1	0.91	0.15	35,35,35,35	0
3	MG	A	458	1/1	0.91	0.16	42,42,42,42	0
3	MG	A	432	1/1	0.92	0.16	2,2,2,2	0
4	K	A	468	1/1	0.92	0.30	36,36,36,36	0
4	K	A	466	1/1	0.92	0.28	43,43,43,43	0
3	MG	A	446	1/1	0.92	0.17	10,10,10,10	0
4	K	A	476	1/1	0.92	0.22	44,44,44,44	0
4	K	A	478	1/1	0.93	0.14	35,35,35,35	0
4	K	A	465	1/1	0.93	0.12	38,38,38,38	0
4	K	A	464	1/1	0.93	0.30	48,48,48,48	0
3	MG	A	419	1/1	0.93	0.18	30,30,30,30	0
3	MG	A	453	1/1	0.93	0.23	61,61,61,61	0
4	K	A	475	1/1	0.94	0.11	15,15,15,15	0
4	K	A	461	1/1	0.94	0.16	4,4,4,4	0
4	K	A	462	1/1	0.94	0.15	7,7,7,7	0
3	MG	A	442	1/1	0.94	0.08	2,2,2,2	0
3	MG	A	435	1/1	0.95	0.12	2,2,2,2	0
3	MG	A	444	1/1	0.96	0.24	18,18,18,18	0
3	MG	A	459	1/1	0.97	0.26	66,66,66,66	0
3	MG	A	433	1/1	0.98	0.07	2,2,2,2	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.