



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 07:29 am BST

PDB ID : 1GA6  
Title : CRYSTAL STRUCTURE ANALYSIS OF PSCP (PSEUDOMONAS SERINE-CARBOXYL PROTEINASE) COMPLEXED WITH A FRAGMENT OF TY-ROSTATIN (THIS ENZYME RENAMED "SEDOLISIN" IN 2003)  
Authors : Wlodawer, A.; Li, M.; Dauter, Z.; Gustchina, A.; Uchida, K.  
Deposited on : 2000-11-29  
Resolution : 1.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1117 (1.06-0.94)
Ramachandran outliers	138981	1043 (1.06-0.94)
Sidechain outliers	138945	1045 (1.06-0.94)
RSRZ outliers	127900	1023 (1.06-0.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	372	<div> <div></div> <div> <div></div> <div>91%</div> <div>7%</div> <div>..</div> </div> </div>
2	I	3	<div> <div>33%</div> <div>33%</div> <div>33%</div> </div>

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3345 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SERINE-CARBOXYL PROTEINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	369	Total	C	N	O	S	4	34	0
			2821	1756	471	591	3			


- Molecule 2 is a protein called FRAGMENT OF TYROSTATIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	I	2	Total	C	N	O	0	0	0
			16	11	2	3			

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 7 3 4	0	1
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	463	Total O 463 463	0	0
6	I	3	Total O 3 3	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SERINE-CARBOXYL PROTEINASE



- Molecule 2: FRAGMENT OF TYROSTATIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.40 Å 97.40 Å 83.35 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 1.00 10.00 – 1.00	Depositor EDS
% Data completeness (in resolution range)	83.9 (10.00-1.00) 100.0 (10.00-1.00)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.79 (at 1.00 Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.123 , 0.131 0.123 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	7.4	Xtriage
Anisotropy	0.125	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 58.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.54$ , $\langle L^2 \rangle = 0.38$	Xtriage
Estimated twinning fraction	0.015 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	3345	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.74	2/3024 (0.1%)	1.12	23/4129 (0.6%)
2	I	0.59	0/12	3.04	2/15 (13.3%)
All	All	0.74	2/3036 (0.1%)	1.14	25/4144 (0.6%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	150	GLU	CD-OE2	5.78	1.32	1.25
1	A	241	GLU	CD-OE1	5.05	1.31	1.25

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	256[A]	ARG	NE-CZ-NH2	-11.79	114.41	120.30
1	A	256[B]	ARG	NE-CZ-NH2	-11.79	114.41	120.30
1	A	82[A]	ASP	CB-CG-OD1	-9.89	109.40	118.30
1	A	82[B]	ASP	CB-CG-OD1	-9.89	109.40	118.30
1	A	152	ARG	NE-CZ-NH2	-8.18	116.21	120.30
1	A	181	TYR	CB-CG-CD2	-8.02	116.19	121.00
2	I	2	TYR	CB-CG-CD1	7.90	125.74	121.00
1	A	179	ARG	NE-CZ-NH2	-7.54	116.53	120.30
1	A	82[A]	ASP	CB-CG-OD2	7.32	124.89	118.30
1	A	82[B]	ASP	CB-CG-OD2	7.32	124.89	118.30
1	A	256[A]	ARG	NE-CZ-NH1	6.99	123.79	120.30
1	A	256[B]	ARG	NE-CZ-NH1	6.99	123.79	120.30
1	A	366[A]	ARG	NE-CZ-NH1	-6.76	116.92	120.30
1	A	366[B]	ARG	NE-CZ-NH1	-6.76	116.92	120.30
1	A	336	TYR	CG-CD1-CE1	-5.73	116.71	121.30
1	A	160[A]	GLN	CB-CA-C	-5.69	99.02	110.40
1	A	160[B]	GLN	CB-CA-C	-5.69	99.02	110.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	60	GLN	CA-CB-CG	5.57	125.64	113.40
1	A	152	ARG	NH1-CZ-NH2	5.26	125.18	119.40
1	A	152	ARG	CD-NE-CZ	-5.16	116.38	123.60
1	A	366[A]	ARG	NH1-CZ-NH2	5.14	125.06	119.40
1	A	366[B]	ARG	NH1-CZ-NH2	5.14	125.06	119.40
2	I	2	TYR	CG-CD2-CE2	5.14	125.41	121.30
1	A	280[A]	LEU	CA-CB-CG	-5.04	103.70	115.30
1	A	280[B]	LEU	CA-CB-CG	-5.04	103.70	115.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2821	0	2613	19	1
2	I	16	0	8	4	0
3	A	1	0	0	0	0
4	A	16	0	12	2	0
5	A	25	0	29	0	1
6	A	463	0	0	7	5
6	I	3	0	0	1	0
All	All	3345	0	2662	22	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:287[B]:SER:OG	6:A:865:HOH:O	1.70	1.09
1:A:43[A]:GLN:OE1	6:A:855:HOH:O	1.76	1.04
4:A:397:ACT:H1	6:A:635:HOH:O	1.66	0.95
1:A:273[B]:LEU:HD12	1:A:280[B]:LEU:HG	1.54	0.90
1:A:111:ASN:HD21	2:I:2:TYR:HD2	1.27	0.83

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:111:ASN:ND2	2:I:2:TYR:HD2	1.90	0.68
1:A:160[A]:GLN:NE2	6:A:604:HOH:O	2.23	0.66
1:A:15[A]:THR:HG23	6:A:575:HOH:O	2.01	0.60
1:A:116[A]:GLN:HE22	1:A:119:ASN:HD22	1.51	0.58
2:I:2:TYR:N	2:I:3:UNK:CA	2.67	0.57
1:A:366[A]:ARG:NH2	6:A:677:HOH:O	2.11	0.55
1:A:326:VAL:O	4:A:398:ACT:H2	2.07	0.54
1:A:273[B]:LEU:HD12	1:A:280[B]:LEU:CG	2.34	0.51
1:A:191:TRP:CD2	1:A:192:PRO:HA	2.46	0.50
1:A:281[A]:GLN:NE2	6:A:735:HOH:O	2.47	0.47
2:I:2:TYR:HB3	6:I:817:HOH:O	2.16	0.46
1:A:238:SER:O	1:A:256[B]:ARG:HD3	2.16	0.45
1:A:41:THR:HA	1:A:82[A]:ASP:OD2	2.17	0.45
1:A:9[A]:ASN:OD1	1:A:10:PRO:HD2	2.20	0.41
1:A:273[B]:LEU:CD1	1:A:280[B]:LEU:HG	2.38	0.41
1:A:60:GLN:HG3	1:A:98:GLN:NE2	2.36	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:640:HOH:O	6:A:640:HOH:O[4_665]	1.52	0.68
1:A:46[B]:GLN:OE1	6:A:744:HOH:O[6_555]	1.65	0.55
6:A:473:HOH:O	6:A:764:HOH:O[4_655]	1.78	0.42
6:A:628:HOH:O	6:A:861:HOH:O[2_654]	1.94	0.26
5:A:391[B]:GOL:O3	6:A:515:HOH:O[6_555]	2.18	0.02

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	401/372 (108%)	394 (98%)	7 (2%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	314/281 (112%)	312 (99%)	2 (1%)	86	61
2	I	1/1 (100%)	0	1 (100%)	0	0
All	All	315/282 (112%)	312 (99%)	3 (1%)	73	47

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	257	ARG
1	A	370	PHE
2	I	2	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 1 is monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	GOL	A	391[A]	-	5,5,5	0.62	0	5,5,5	0.66	0
5	GOL	A	391[B]	-	5,5,5	1.35	1 (20%)	5,5,5	0.98	0
4	ACT	A	395	-	1,3,3	2.98	1 (100%)	0,3,3	0.00	-
5	GOL	A	392	-	5,5,5	1.04	1 (20%)	5,5,5	0.65	0
4	ACT	A	398	-	1,3,3	3.19	1 (100%)	0,3,3	0.00	-
5	GOL	A	393	-	5,5,5	0.59	0	5,5,5	1.33	1 (20%)
5	GOL	A	394	-	5,5,5	0.48	0	5,5,5	0.82	0
4	ACT	A	397	-	1,3,3	1.31	0	0,3,3	0.00	-
4	ACT	A	396	-	1,3,3	2.77	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	391[A]	-	-	0/4/4/4	-
5	GOL	A	392	-	-	0/4/4/4	-
5	GOL	A	391[B]	-	-	0/4/4/4	-
5	GOL	A	393	-	-	1/4/4/4	-
5	GOL	A	394	-	-	0/4/4/4	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	398	ACT	CH3-C	3.19	1.52	1.48
4	A	395	ACT	CH3-C	2.98	1.52	1.48
4	A	396	ACT	CH3-C	2.77	1.52	1.48
5	A	391[B]	GOL	O3-C3	2.69	1.53	1.42
5	A	392	GOL	O1-C1	-2.10	1.33	1.42

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
5	A	393	GOL	O2-C2-C1	2.28	119.15	109.12

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	393	GOL	C1-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	391[B]	GOL	0	1
4	A	398	ACT	1	0
4	A	397	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	369/372 (99%)	-0.77	2 (0%) 91 86	5, 8, 17, 42	1 (0%)
2	I	1/3 (33%)	2.33	1 (100%) 0 0	25, 25, 25, 25	1 (100%)
All	All	370/375 (98%)	-0.76	3 (0%) 86 81	5, 8, 18, 42	2 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ALA	5.1
2	I	2	TYR	2.3
1	A	68	ASN	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	ACT	A	398	4/4	0.80	0.14	21,21,34,38	0
5	GOL	A	393	6/6	0.86	0.14	19,21,24,27	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	ACT	A	395	4/4	0.87	0.15	13,14,14,18	0
5	GOL	A	394	6/6	0.91	0.14	18,23,25,38	0
4	ACT	A	397	4/4	0.93	0.13	15,16,23,23	0
5	GOL	A	391[B]	6/6	0.98	0.08	8,12,13,14	1
5	GOL	A	391[A]	6/6	0.98	0.08	8,11,13,14	1
4	ACT	A	396	4/4	0.98	0.06	10,11,12,13	0
5	GOL	A	392	6/6	0.99	0.10	9,10,13,13	0
3	CA	A	374	1/1	1.00	0.02	7,7,7,7	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.