



# Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 11:51 pm BST

PDB ID : 2GAA  
Title : Crystal structure of YFH7 from *Saccharomyces cerevisiae*: a putative P-loop containing kinase with a circular permutation.  
Authors : Chaptal, V.; Morera, S.  
Deposited on : 2006-03-08  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	359	<div> <div>5%</div> <div> <div></div> <div>71%</div> <div>19%</div> <div>• 8%</div> </div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2810 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Hypothetical 39.9 kDa protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	329	2625	1655	456	501	13	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	354	HIS	-	EXPRESSION TAG	UNP P43591
A	355	HIS	-	EXPRESSION TAG	UNP P43591
A	356	HIS	-	EXPRESSION TAG	UNP P43591
A	357	HIS	-	EXPRESSION TAG	UNP P43591
A	358	HIS	-	EXPRESSION TAG	UNP P43591
A	359	HIS	-	EXPRESSION TAG	UNP P43591

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	180	Total	O	0	0
			180	180		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hypothetical 39.9 kDa protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	134.76 Å   134.76 Å   48.68 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	20.00 – 1.95 18.69 – 1.95	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-1.95) 99.1 (18.69-1.95)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.03 (at 1.94 Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.193 , 0.232 0.188 , 0.223	Depositor DCC
$R_{free}$ test set	1156 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.6	Xtriage
Anisotropy	0.137	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 56.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.029 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2810	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CME, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	2/2635 (0.1%)	0.60	2/3558 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	209	LYS	C-N	8.41	1.53	1.34
1	A	46	GLN	C-N	-6.90	1.18	1.34

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	46	GLN	O-C-N	-7.37	110.92	122.70
1	A	209	LYS	CA-C-N	-5.59	104.90	117.20

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	120	GLU	Mainchain
1	A	121	CME	Mainchain
1	A	208	CME	Mainchain

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	A	209	LYS	Mainchain
1	A	45	CME	Mainchain
1	A	46	GLN	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2625	0	2632	49	0
2	A	5	0	0	1	0
3	A	180	0	0	4	0
All	All	2810	0	2632	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:68:ARG:HE	1:A:68:ARG:H	1.09	0.95
1:A:68:ARG:NE	1:A:68:ARG:H	1.85	0.74
1:A:176:ASP:OD1	1:A:178:GLN:HG2	1.87	0.73
1:A:68:ARG:N	1:A:68:ARG:HE	1.88	0.71
1:A:308:LYS:O	1:A:312:GLN:HG3	1.91	0.69
1:A:315:LEU:HD23	1:A:325:LYS:NZ	2.09	0.68
1:A:169:ARG:NH1	1:A:169:ARG:HB2	2.11	0.66
1:A:315:LEU:HD23	1:A:325:LYS:HZ3	1.60	0.63
1:A:58:GLU:C	1:A:60:PRO:HD3	2.20	0.61
1:A:82:LEU:HD12	1:A:86:LYS:HE3	1.85	0.58
1:A:313:SER:HB2	1:A:315:LEU:HD13	1.87	0.57
1:A:59:HIS:N	1:A:60:PRO:HD3	2.19	0.57
1:A:38:SER:O	1:A:42:GLU:HG3	2.05	0.56
1:A:96:LEU:HD11	1:A:169:ARG:HG2	1.87	0.56
1:A:250:THR:HG23	1:A:253:GLN:CG	2.36	0.55
1:A:183:ARG:HD3	3:A:2997:HOH:O	2.07	0.54
1:A:76:VAL:HB	1:A:110:LYS:HD3	1.89	0.54

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:183:ARG:HG2	1:A:186:SER:HB3	1.90	0.54
1:A:344:VAL:HG23	1:A:347:ILE:HG13	1.90	0.54
1:A:178:GLN:HG3	1:A:179:THR:N	2.24	0.52
1:A:349:HIS:HE1	3:A:2956:HOH:O	1.95	0.48
1:A:169:ARG:HH11	1:A:169:ARG:HB2	1.77	0.48
1:A:182:LYS:HD3	1:A:182:LYS:O	2.13	0.48
1:A:245:ALA:O	1:A:247:LYS:HE3	2.15	0.46
1:A:52:TYR:CE1	1:A:226:LEU:HD11	2.50	0.46
1:A:40:ILE:HD12	1:A:295:ILE:HD11	1.98	0.46
1:A:239:VAL:CG2	1:A:254:TYR:HB2	2.46	0.46
1:A:178:GLN:CG	1:A:179:THR:N	2.79	0.45
1:A:75:LEU:HD13	1:A:201:ILE:HG23	1.99	0.45
1:A:310:HIS:HD2	1:A:325:LYS:HD3	1.82	0.45
1:A:250:THR:HG23	1:A:253:GLN:HG3	1.98	0.44
1:A:187:PRO:HG3	1:A:269:TYR:CE1	2.53	0.44
1:A:7:LEU:HD22	1:A:293:TYR:CZ	2.52	0.44
1:A:74:ASN:O	1:A:204:LYS:NZ	2.43	0.44
1:A:101:VAL:HG23	1:A:102:GLU:HG3	2.00	0.43
1:A:87:VAL:O	1:A:91:ILE:HG12	2.18	0.43
1:A:281:TYR:CE2	1:A:344:VAL:HG11	2.54	0.43
1:A:80:LYS:NZ	1:A:86:LYS:NZ	2.66	0.43
1:A:181:HIS:HD2	3:A:3027:HOH:O	2.01	0.43
1:A:226:LEU:HD12	1:A:227:SER:N	2.34	0.42
1:A:226:LEU:C	1:A:226:LEU:HD12	2.40	0.41
1:A:66:ASN:ND2	1:A:229:THR:HA	2.35	0.41
1:A:330:ASP:HA	1:A:333:ASN:OD1	2.20	0.41
1:A:209:LYS:HE2	1:A:209:LYS:HB3	1.74	0.41
1:A:354:HIS:HB3	3:A:2959:HOH:O	2.20	0.41
1:A:175:LYS:HD2	1:A:175:LYS:HA	1.89	0.40
1:A:37:LYS:HG2	2:A:2847:SO4:O2	2.21	0.40
1:A:241:GLY:HA3	1:A:253:GLN:HG3	2.04	0.40
1:A:274:GLN:O	1:A:275:GLU:HG3	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	318/359 (89%)	307 (96%)	10 (3%)	1 (0%)	41 30

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	47	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	293/322 (91%)	286 (98%)	7 (2%)	49 40

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	68	ARG
1	A	105	ASN
1	A	119	GLU
1	A	193	ASN
1	A	194	ASN
1	A	209	LYS
1	A	275	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	ASN
1	A	46	GLN
1	A	105	ASN
1	A	181	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	194	ASN
1	A	276	ASN
1	A	310	HIS
1	A	346	ASN
1	A	349	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CME	A	121	1	8,9,10	1.05	1 (12%)	5,9,11	0.44	0
1	CME	A	208	1	8,9,10	1.08	1 (12%)	5,9,11	0.51	0
1	CME	A	45	1	8,9,10	1.04	1 (12%)	5,9,11	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	121	1	-	0/5/8/10	-
1	CME	A	208	1	-	0/5/8/10	-
1	CME	A	45	1	-	1/5/8/10	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	208	CME	OH-CZ	-2.76	1.27	1.42
1	A	121	CME	OH-CZ	-2.75	1.27	1.42
1	A	45	CME	OH-CZ	-2.70	1.28	1.42

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	45	CME	CE-SD-SG-CB

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	2847	-	4,4,4	0.28	0	6,6,6	0.13	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2847	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	46:GLN	C	47:ILE	N	1.18
1	A	207:LEU	C	208:CME	N	1.14

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	326/359 (90%)	0.31	19 (5%)	23 31	14, 25, 43, 65	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	354	HIS	7.5
1	A	218	SER	5.3
1	A	119	GLU	4.6
1	A	138	ALA	4.1
1	A	120	GLU	4.1
1	A	345	ASP	3.9
1	A	178	GLN	3.6
1	A	209	LYS	3.6
1	A	211	SER	3.5
1	A	246	LEU	3.4
1	A	210	VAL	3.0
1	A	353	ASP	2.9
1	A	175	LYS	2.9
1	A	113	ALA	2.8
1	A	179	THR	2.5
1	A	152	SER	2.5
1	A	219	THR	2.3
1	A	85	ASN	2.1
1	A	346	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	CME	A	208	10/11	0.79	0.26	21,24,30,32	10
1	CME	A	121	10/11	0.91	0.14	45,48,50,53	0
1	CME	A	45	10/11	0.92	0.14	21,26,45,49	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	2847	5/5	0.97	0.07	32,36,37,38	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.