



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2021 – 07:47 AM EDT

PDB ID : 3GA0
Title : CtBP1/BARS Gly172->Glu mutant structure: impairing NAD(H) binding and dimerization
Authors : Nardini, M.; Valente, C.; Ricagno, S.; Luini, A.; Corda, D.; Bolognesi, M.
Deposited on : 2009-02-16
Resolution : 3.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1026 (3.48-3.32)
Clashscore	141614	1055 (3.48-3.32)
Ramachandran outliers	138981	1038 (3.48-3.32)
Sidechain outliers	138945	1038 (3.48-3.32)
RSRZ outliers	127900	2173 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	358	<div> <div>6%</div> <div> <div></div> <div>73%</div> <div>18%</div> <div>6%</div> </div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2616 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called C-terminal-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	337	2609	1629	478	489	13	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	expression tag	UNP Q9Z2F5
A	-6	GLY	-	expression tag	UNP Q9Z2F5
A	-5	HIS	-	expression tag	UNP Q9Z2F5
A	-4	HIS	-	expression tag	UNP Q9Z2F5
A	-3	HIS	-	expression tag	UNP Q9Z2F5
A	-2	HIS	-	expression tag	UNP Q9Z2F5
A	-1	HIS	-	expression tag	UNP Q9Z2F5
A	0	HIS	-	expression tag	UNP Q9Z2F5
A	172	GLU	GLY	engineered mutation	UNP Q9Z2F5

- Molecule 2 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			3	1	2		
2	A	1	Total	C	O	0	0
			3	1	2		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O	0	0
			1	1		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: C-terminal-binding protein 1

Chain A: 6% 73% 18% 6%



4 Data and refinement statistics

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants a, b, c, α , β , γ	89.20Å 89.20Å 160.26Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.75 – 3.40 29.80 – 3.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.75-3.40) 99.8 (29.80-3.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.76 (at 3.39Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.262 , 0.334 0.227 , 0.308	Depositor DCC
R_{free} test set	560 reflections (9.97%)	wwPDB-VP
Wilson B-factor (Å ²)	83.3	Xtriage
Anisotropy	0.230	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 32.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2616	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/2652	0.54	0/3590

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	11	GLY	Peptide
1	A	13	MET	Peptide
1	A	14	HIS	Peptide
1	A	16	ARG	Peptide
1	A	283	SER	Peptide
1	A	286	PHE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2609	0	2615	41	0
2	A	6	0	2	0	0
3	A	1	0	0	0	0
All	All	2616	0	2617	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:284:GLU:HB3	1:A:285:PRO:HD2	1.55	0.89
1:A:254:ALA:O	1:A:280:VAL:HG21	1.72	0.88
1:A:256:GLY:C	1:A:282:GLU:HB2	1.97	0.85
1:A:252:ASN:HB3	1:A:278:LEU:HD23	1.65	0.78
1:A:218:HIS:HA	1:A:245:ARG:HH21	1.52	0.74
1:A:201:GLU:HG2	1:A:208:ARG:HD2	1.72	0.71
1:A:325:ARG:O	1:A:329:THR:HG22	1.94	0.66
1:A:37:VAL:HG21	1:A:325:ARG:HG3	1.84	0.58
1:A:284:GLU:HB3	1:A:285:PRO:CD	2.32	0.58
1:A:43:CYS:SG	1:A:55:VAL:HG21	2.44	0.58
1:A:16:ARG:O	1:A:16:ARG:HG3	2.07	0.53
1:A:287:SER:O	1:A:289:SER:N	2.41	0.53
1:A:14:HIS:O	1:A:15:PRO:C	2.47	0.53
1:A:173:ARG:O	1:A:175:GLY:N	2.43	0.52
1:A:287:SER:C	1:A:289:SER:H	2.14	0.51
1:A:256:GLY:CA	1:A:282:GLU:HB2	2.40	0.51
1:A:161:ILE:O	1:A:164:GLU:HB2	2.11	0.51
1:A:72:ARG:HD3	1:A:98:SER:OG	2.13	0.48
1:A:11:GLY:HA3	1:A:12:PRO:HD3	1.45	0.48
1:A:128:LEU:HB3	1:A:249:PHE:CD1	2.50	0.47
1:A:23:ASP:HB2	1:A:67:THR:HB	1.96	0.46
1:A:252:ASN:HB3	1:A:278:LEU:CD2	2.40	0.46
1:A:30:GLU:OE1	1:A:30:GLU:N	2.50	0.45
1:A:97:LYS:HG2	1:A:344:HIS:CE1	2.51	0.45
1:A:284:GLU:O	1:A:286:PHE:N	2.50	0.45
1:A:283:SER:O	1:A:284:GLU:O	2.35	0.44
1:A:31:MET:O	1:A:35:LYS:HB2	2.17	0.44
1:A:211:THR:O	1:A:214:ASP:HB2	2.18	0.44
1:A:105:ALA:HB2	1:A:345:LEU:HD11	2.01	0.43
1:A:173:ARG:O	1:A:174:VAL:C	2.57	0.43
1:A:25:ARG:HA	1:A:42:PHE:CD2	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:78:PHE:HB3	1:A:81:LEU:HB2	2.00	0.43
1:A:256:GLY:HA3	1:A:282:GLU:HB2	2.00	0.43
1:A:15:PRO:HB2	1:A:16:ARG:H	1.67	0.42
1:A:33:ILE:O	1:A:325:ARG:NH1	2.49	0.42
1:A:19:VAL:HA	1:A:61:GLY:O	2.19	0.41
1:A:296:ALA:HA	1:A:297:PRO:HD3	1.91	0.41
1:A:242:LYS:HA	1:A:242:LYS:HD2	1.90	0.41
1:A:169:ILE:HD12	1:A:224:LEU:CD2	2.50	0.41
1:A:51:ILE:HG22	1:A:55:VAL:HB	2.03	0.40
1:A:201:GLU:H	1:A:201:GLU:HG3	1.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	335/358 (94%)	302 (90%)	27 (8%)	6 (2%)	8	32

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	15	PRO
1	A	254	ALA
1	A	285	PRO
1	A	14	HIS
1	A	283	SER
1	A	12	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/295 (94%)	257 (92%)	21 (8%)	13	41

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	ASN
1	A	16	ARG
1	A	48	THR
1	A	82	ARG
1	A	98	SER
1	A	131	ARG
1	A	139	LEU
1	A	144	ARG
1	A	188	ASN
1	A	201	GLU
1	A	216	LEU
1	A	226	CYS
1	A	229	ASN
1	A	255	ARG
1	A	281	HIS
1	A	282	GLU
1	A	283	SER
1	A	285	PRO
1	A	288	PHE
1	A	298	ASN
1	A	313	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	ASN
1	A	127	ASN
1	A	146	GLN
1	A	234	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	252	ASN
1	A	311	GLN
1	A	344	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FMT	A	801	-	0,2,2	-	-	0,1,1	-	-
2	FMT	A	800	-	0,2,2	-	-	0,1,1	-	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	337/358 (94%)	0.20	20 (5%) 22 23	26, 59, 76, 94	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	14	HIS	8.2
1	A	13	MET	6.9
1	A	12	PRO	5.5
1	A	10	ASN	5.2
1	A	346	THR	4.4
1	A	11	GLY	4.3
1	A	289	SER	3.2
1	A	288	PHE	2.7
1	A	144	ARG	2.6
1	A	286	PHE	2.6
1	A	15	PRO	2.6
1	A	230	GLU	2.3
1	A	330	GLY	2.3
1	A	101	ASP	2.2
1	A	232	ASN	2.2
1	A	287	SER	2.1
1	A	231	HIS	2.1
1	A	253	THR	2.0
1	A	123	CYS	2.0
1	A	120	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FMT	A	800	3/3	0.83	0.53	55,55,55,55	0
2	FMT	A	801	3/3	0.92	0.41	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.