



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2021 – 06:47 PM EDT

PDB ID : 3GC6
Title : Structural insights into the catalytic mechanism of CD38: Evidence for a conformationally flexible covalent enzyme-substrate complex.
Authors : Egea, P.F.; Muller-Steffner, H.; Stroud, R.M.; Oppenheimer, N.; Kellenberger, E.; Schuber, F.
Deposited on : 2009-02-21
Resolution : 1.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4009 (1.54-1.50)
Clashscore	141614	4249 (1.54-1.50)
Ramachandran outliers	138981	4148 (1.54-1.50)
Sidechain outliers	138945	4146 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	247	85% 13% .
1	B	247	81% 14% . .
2	C	2	50% 50%
2	D	2	50% 50%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4591 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ecto-NAD⁺ glycohydrolase (CD38 molecule).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	242	Total	C	N	O	S	0	18	0
			2056	1279	377	384	16			
1	B	237	Total	C	N	O	S	0	10	0
			1966	1228	358	364	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	218	GLN	GLU	engineered mutation	UNP Q9TTF5
B	218	GLN	GLU	engineered mutation	UNP Q9TTF5

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	297	Total	O	0	0
			297	297		
4	B	191	Total	O	0	0
			191	191		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ecto-NAD⁺ glycohydrolase (CD38 molecule)



- Molecule 1: Ecto-NAD⁺ glycohydrolase (CD38 molecule)



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	47.03Å 80.16Å 152.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.95 – 1.51 44.94 – 1.51	Depositor EDS
% Data completeness (in resolution range)	81.6 (44.95-1.51) 81.6 (44.94-1.51)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.25 (at 1.51Å)	Xtriage
Refinement program	REFMAC 5.4.0067	Depositor
R, R_{free}	0.187 , 0.210 0.197 , 0.219	Depositor DCC
R_{free} test set	5756 reflections (7.72%)	wwPDB-VP
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 35.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4591	wwPDB-VP
Average B, all atoms (Å ²)	7.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/2110	0.66	0/2847
1	B	0.46	0/2014	0.57	0/2721
All	All	0.53	0/4124	0.62	0/5568

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2056	0	2007	33	0
1	B	1966	0	1925	44	0
2	C	28	0	25	0	0
2	D	28	0	25	1	0
3	A	15	0	0	0	0
3	B	10	0	0	0	0
4	A	297	0	0	3	1
4	B	191	0	0	4	0
All	All	4591	0	3982	76	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:264[B]:ARG:HG2	1:A:266[B]:MET:HE1	1.22	1.16
1:B:61:THR:O	1:B:65[B]:THR:HG22	1.55	1.06
1:A:264[B]:ARG:CG	1:A:266[B]:MET:HE1	1.93	0.99
1:A:264[B]:ARG:HG2	1:A:266[B]:MET:CE	1.98	0.93
1:B:66:ILE:HG22	1:B:67[B]:ARG:HG2	1.57	0.87
1:A:123:GLU:OE1	4:A:410:HOH:O	1.93	0.85
1:B:213:ILE:HG22	4:B:363:HOH:O	1.78	0.82
1:B:110:VAL:CG2	1:B:115:GLN:HG3	2.09	0.82
1:B:122[B]:LYS:O	1:B:126:HIS:HD2	1.64	0.80
1:B:110:VAL:HG21	1:B:115:GLN:HG3	1.68	0.76
1:B:213:ILE:C	1:B:213:ILE:HD13	2.05	0.74
1:A:122:LYS:O	1:A:126:HIS:HD2	1.71	0.74
1:B:236:THR:OG1	4:B:469:HOH:O	2.06	0.73
1:B:122[A]:LYS:O	1:B:126:HIS:HD2	1.74	0.70
1:A:264[B]:ARG:NE	1:A:266[B]:MET:HE3	2.08	0.68
1:A:264[B]:ARG:CD	1:A:266[B]:MET:HE1	2.23	0.68
1:B:61:THR:O	1:B:65[B]:THR:CG2	2.38	0.66
1:A:175:ASN:ND2	1:A:178:SER:H	1.96	0.64
1:A:264[B]:ARG:NE	1:A:266[B]:MET:CE	2.62	0.63
1:B:61:THR:HG23	1:B:65[B]:THR:HG21	1.79	0.63
1:A:175:ASN:HD21	1:A:178:SER:H	1.46	0.62
1:A:66:ILE:HG22	1:A:67:ARG:HD3	1.80	0.62
1:B:110:VAL:HG22	1:B:115:GLN:HG3	1.79	0.62
1:B:124:LEU:HD13	1:B:271:ARG:HG3	1.82	0.62
1:A:211[B]:MET:O	1:A:216:ARG:NH1	2.32	0.61
1:A:211[A]:MET:O	1:A:216:ARG:NH1	2.32	0.61
1:A:99:TYR:O	1:A:103[B]:ILE:HG12	2.01	0.61
1:B:65[B]:THR:HG23	1:B:66:ILE:HG13	1.82	0.61
1:B:175:ASN:ND2	1:B:178:SER:H	1.99	0.60
1:A:117:PHE:HB2	1:A:198[B]:VAL:HG22	1.85	0.59
1:B:175:ASN:HD21	1:B:178:SER:H	1.49	0.58
1:B:191:SER:HB2	4:B:477:HOH:O	2.03	0.57
1:B:196:VAL:HG23	1:B:228:LEU:CD1	2.34	0.57
1:B:122[B]:LYS:O	1:B:126:HIS:CD2	2.53	0.57
1:A:264[B]:ARG:CG	1:A:266[B]:MET:CE	2.71	0.56
1:B:51:PHE:HB3	1:B:160[B]:LEU:HD11	1.88	0.55
1:A:264[B]:ARG:HG2	1:A:266[B]:MET:SD	2.48	0.53
1:A:83:MET:HE1	1:A:160:LEU:HD11	1.90	0.53
1:A:122:LYS:O	1:A:126:HIS:CD2	2.59	0.53
1:B:192:ALA:HB1	1:B:196:VAL:HG11	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:73:ARG:NH1	1:B:105:LEU:O	2.44	0.51
1:B:126:HIS:HE1	1:B:138:GLU:OE1	1.94	0.51
1:B:196:VAL:HG23	1:B:228:LEU:HD13	1.94	0.50
1:B:213:ILE:CD1	1:B:218:GLN:NE2	2.75	0.49
1:A:236:THR:HB	4:A:464:HOH:O	2.13	0.48
1:A:264[B]:ARG:CZ	1:A:266[B]:MET:HE3	2.43	0.48
1:A:266[B]:MET:SD	1:A:266[B]:MET:N	2.87	0.48
1:B:61:THR:C	1:B:65[B]:THR:HG22	2.32	0.48
1:B:67[A]:ARG:O	1:B:67[A]:ARG:HG2	2.14	0.48
1:B:122[A]:LYS:O	1:B:126:HIS:CD2	2.63	0.47
1:B:196:VAL:HG23	1:B:228:LEU:HD12	1.97	0.47
1:B:213:ILE:HD13	1:B:213:ILE:O	2.15	0.47
1:B:115:GLN:NE2	1:B:192:ALA:HA	2.30	0.46
1:B:61:THR:HG23	1:B:65[B]:THR:CG2	2.44	0.46
1:B:87:ILE:HD11	1:B:160[A]:LEU:HD22	1.97	0.45
1:A:264[B]:ARG:HH11	1:A:264[B]:ARG:HG3	1.80	0.45
1:B:185:SER:CB	1:B:213:ILE:HD11	2.45	0.45
1:B:67[A]:ARG:NH2	4:B:388:HOH:O	2.48	0.45
1:B:55[A]:ILE:HD11	1:B:149:LEU:HD12	1.99	0.45
1:B:126:HIS:CE1	1:B:138:GLU:OE1	2.70	0.44
1:A:269:LEU:HD21	1:A:277[A]:ARG:HH11	1.81	0.44
1:A:126:HIS:HE1	1:A:138:GLU:OE1	2.01	0.44
1:A:211[A]:MET:HA	1:A:211[A]:MET:CE	2.49	0.43
1:A:262:LYS:HE3	1:A:264[A]:ARG:NH1	2.34	0.41
1:B:55[A]:ILE:HD11	1:B:149:LEU:CD1	2.50	0.41
1:B:96:LYS:HG2	1:B:183:VAL:CG2	2.50	0.41
1:B:186:GLU:HG3	1:B:217:VAL:HG13	2.03	0.41
1:A:100:ASN:HD21	1:B:104:ASN:HD21	1.69	0.41
1:A:264[B]:ARG:NE	1:A:266[B]:MET:HE1	2.33	0.41
1:A:79[B]:ARG:O	1:A:83:MET:HG2	2.19	0.41
1:B:83:MET:CE	1:B:160[A]:LEU:HD21	2.50	0.41
1:B:83:MET:HE1	1:B:160[A]:LEU:HD21	2.02	0.41
1:A:208:PHE:O	1:A:248:SER:HB2	2.21	0.41
1:B:124:LEU:HD22	2:D:1:NAG:H81	2.03	0.41
1:A:235:ASP:HB2	1:A:238:LYS:HE2	2.03	0.40
1:A:109[B]:THR:CG2	4:A:416:HOH:O	2.70	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:344:HOH:O	4:A:454:HOH:O[3_655]	1.95	0.25

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/247 (104%)	253 (98%)	4 (2%)	0	100	100
1	B	245/247 (99%)	240 (98%)	5 (2%)	0	100	100
All	All	502/494 (102%)	493 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/218 (106%)	226 (98%)	4 (2%)	60	32
1	B	218/218 (100%)	213 (98%)	5 (2%)	50	20
All	All	448/436 (103%)	439 (98%)	9 (2%)	53	24

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	GLN
1	A	67	ARG
1	A	82	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	193	CYS
1	B	39	LEU
1	B	82	PHE
1	B	110	VAL
1	B	197	ARG
1	B	213	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	GLN
1	A	100	ASN
1	A	126	HIS
1	A	175	ASN
1	A	221	ASN
1	B	115	GLN
1	B	126	HIS
1	B	175	ASN
1	B	221	ASN
1	B	273	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	2,1	14,14,15	0.59	0	17,19,21	0.89	1 (5%)
2	NAG	C	2	2	14,14,15	0.56	0	17,19,21	0.74	0
2	NAG	D	1	2,1	14,14,15	0.47	0	17,19,21	1.00	1 (5%)
2	NAG	D	2	2	14,14,15	0.49	0	17,19,21	0.88	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	NAG	D	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	NAG	O5-C1-C2	-2.40	107.50	111.29
2	D	2	NAG	O5-C5-C6	2.21	110.67	107.20
2	C	1	NAG	O5-C1-C2	-2.00	108.12	111.29

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	NAG	O5-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
2	D	2	NAG	C4-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	280	-	4,4,4	0.15	0	6,6,6	0.17	0
3	SO4	A	279	-	4,4,4	0.16	0	6,6,6	0.08	0
3	SO4	A	281	-	4,4,4	0.12	0	6,6,6	0.39	0
3	SO4	A	280	-	4,4,4	0.14	0	6,6,6	0.17	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	279	-	4,4,4	0.13	0	6,6,6	0.09	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.