



Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 10:09 pm BST

PDB ID : 4GDO
Title : Structure of a fragment of the rod domain of plectin
Authors : De Pereda, J.M.; Buey, R.M.; Uson, I.; Sammito, M.D.; De Marino, I.
Deposited on : 2012-08-01
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	43	
1	B	43	
1	C	43	
1	D	43	
1	E	43	
1	F	43	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 3410 atoms, of which 1594 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Plectin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	42	Total	C	H	N	O	S	0	0	0
			666	200	329	68	68	1			
1	B	41	Total	C	H	N	O	S	0	0	0
			655	197	324	67	66	1			
1	C	38	Total	C	H	N	O		0	0	0
			571	173	283	56	59				
1	D	41	Total	C	H	N	O	S	0	0	0
			619	186	304	64	64	1			
1	E	31	Total	C	H	N	O		0	0	0
			388	121	188	40	39				
1	F	30	Total	C	H	N	O		0	0	0
			348	110	166	36	36				

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1378	GLY	-	EXPRESSION TAG	UNP Q15149
A	1379	SER	-	EXPRESSION TAG	UNP Q15149
A	1380	HIS	-	EXPRESSION TAG	UNP Q15149
A	1381	MET	-	EXPRESSION TAG	UNP Q15149
B	1378	GLY	-	EXPRESSION TAG	UNP Q15149
B	1379	SER	-	EXPRESSION TAG	UNP Q15149
B	1380	HIS	-	EXPRESSION TAG	UNP Q15149
B	1381	MET	-	EXPRESSION TAG	UNP Q15149
C	1378	GLY	-	EXPRESSION TAG	UNP Q15149
C	1379	SER	-	EXPRESSION TAG	UNP Q15149
C	1380	HIS	-	EXPRESSION TAG	UNP Q15149
C	1381	MET	-	EXPRESSION TAG	UNP Q15149
D	1378	GLY	-	EXPRESSION TAG	UNP Q15149
D	1379	SER	-	EXPRESSION TAG	UNP Q15149
D	1380	HIS	-	EXPRESSION TAG	UNP Q15149
D	1381	MET	-	EXPRESSION TAG	UNP Q15149
E	1378	GLY	-	EXPRESSION TAG	UNP Q15149

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	1379	SER	-	EXPRESSION TAG	UNP Q15149
E	1380	HIS	-	EXPRESSION TAG	UNP Q15149
E	1381	MET	-	EXPRESSION TAG	UNP Q15149
F	1378	GLY	-	EXPRESSION TAG	UNP Q15149
F	1379	SER	-	EXPRESSION TAG	UNP Q15149
F	1380	HIS	-	EXPRESSION TAG	UNP Q15149
F	1381	MET	-	EXPRESSION TAG	UNP Q15149

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	48	Total O 48 48	0	0
2	B	41	Total O 41 41	0	0
2	C	26	Total O 26 26	0	0
2	D	39	Total O 39 39	0	0
2	E	2	Total O 2 2	0	0
2	F	7	Total O 7 7	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Plectin

Chain A: 





• Molecule 1: Plectin

Chain B: 




• Molecule 1: Plectin

Chain C: 




• Molecule 1: Plectin

Chain D: 




• Molecule 1: Plectin

Chain E: 



• Molecule 1: Plectin



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	90.62Å 35.87Å 75.69Å 90.00° 94.11° 90.00°	Depositor
Resolution (Å)	45.19 – 1.70 45.19 – 1.70	Depositor EDS
% Data completeness (in resolution range)	85.3 (45.19-1.70) 85.3 (45.19-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 1.70Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1117)	Depositor
R, R_{free}	0.213 , 0.248 0.215 , 0.254	Depositor DCC
R_{free} test set	1189 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	19.8	Xtriage
Anisotropy	0.500	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 62.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3410	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/338	0.54	0/447
1	B	0.46	0/332	0.54	0/439
1	C	0.40	0/288	0.47	0/383
1	D	0.43	0/315	0.57	0/418
1	E	0.28	0/200	0.34	0/270
1	F	0.26	0/182	0.28	0/247
All	All	0.42	0/1655	0.49	0/2204

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	337	329	331	5	0
1	B	331	324	326	5	0
1	C	288	283	278	6	0
1	D	315	304	297	3	0
1	E	200	188	171	3	0
1	F	182	166	145	2	0
2	A	48	0	0	0	0
2	B	41	0	0	1	0
2	C	26	0	0	0	0
2	D	39	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	2	0	0	0	0
2	F	7	0	0	0	0
All	All	1816	1594	1548	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1380:HIS:N	2:B:1528:HOH:O	2.22	0.72
1:D:1382:ARG:NH2	2:D:1520:HOH:O	2.26	0.68
1:A:1381:MET:HE2	1:C:1397:GLU:HG3	1.82	0.62
1:C:1419:GLU:OE2	1:D:1388:ARG:NH1	2.35	0.60
1:E:1406:HIS:HA	1:F:1402:LEU:HD13	1.86	0.56
1:A:1380:HIS:NE2	1:C:1393:GLU:OE1	2.40	0.55
1:B:1397:GLU:OE1	1:B:1397:GLU:HA	2.09	0.51
1:A:1419:GLU:HB3	1:B:1388:ARG:HE	1.75	0.51
1:A:1381:MET:CE	1:C:1397:GLU:HG3	2.43	0.48
1:E:1406:HIS:HA	1:F:1402:LEU:CD1	2.46	0.45
1:B:1414:GLU:O	1:B:1418:LYS:HG3	2.16	0.45
1:A:1381:MET:HE2	1:C:1397:GLU:CG	2.48	0.42
1:C:1393:GLU:OE2	1:E:1396:LEU:HB2	2.20	0.41
1:B:1382:ARG:NH1	1:B:1384:GLU:OE1	2.53	0.41
1:D:1404:GLU:OE2	2:D:1516:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	40/43 (93%)	40 (100%)	0	0	100	100
1	B	39/43 (91%)	39 (100%)	0	0	100	100
1	C	36/43 (84%)	36 (100%)	0	0	100	100
1	D	39/43 (91%)	39 (100%)	0	0	100	100
1	E	29/43 (67%)	29 (100%)	0	0	100	100
1	F	28/43 (65%)	28 (100%)	0	0	100	100
All	All	211/258 (82%)	211 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	31/31 (100%)	30 (97%)	1 (3%)	39	20
1	B	30/31 (97%)	30 (100%)	0	100	100
1	C	24/31 (77%)	24 (100%)	0	100	100
1	D	26/31 (84%)	26 (100%)	0	100	100
1	E	11/31 (36%)	10 (91%)	1 (9%)	9	2
1	F	8/31 (26%)	6 (75%)	2 (25%)	0	0
All	All	130/186 (70%)	126 (97%)	4 (3%)	40	21

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1382	ARG
1	E	1412	GLN
1	F	1397	GLU
1	F	1416	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	42/43 (97%)	0.51	0 100 100	12, 20, 49, 75	0
1	B	41/43 (95%)	0.44	1 (2%) 59 63	13, 22, 57, 70	0
1	C	38/43 (88%)	0.62	4 (10%) 6 7	15, 30, 101, 138	0
1	D	41/43 (95%)	0.72	6 (14%) 2 2	14, 27, 90, 104	0
1	E	31/43 (72%)	2.44	16 (51%) 0 0	59, 68, 100, 125	0
1	F	30/43 (69%)	3.98	26 (86%) 0 0	64, 94, 124, 150	0
All	All	223/258 (86%)	1.29	53 (23%) 0 0	12, 35, 108, 150	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	1404	GLU	8.9
1	F	1407	ALA	8.1
1	F	1408	GLN	7.7
1	E	1402	LEU	7.0
1	F	1411	ALA	6.9
1	E	1396	LEU	5.9
1	E	1405	ALA	5.7
1	F	1396	LEU	5.6
1	F	1394	ALA	5.6
1	C	1389	LEU	5.5
1	F	1392	VAL	5.4
1	F	1401	GLN	5.2
1	D	1420	LEU	4.9
1	F	1409	ALA	4.9
1	F	1400	ARG	4.7
1	E	1390	ALA	4.5
1	E	1420	LEU	4.5
1	F	1402	LEU	4.3
1	E	1417	ALA	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	F	1403	ALA	4.0
1	F	1410	LYS	4.0
1	F	1414	GLU	3.9
1	F	1389	LEU	3.6
1	F	1405	ALA	3.6
1	F	1413	ALA	3.6
1	E	1419	GLU	3.5
1	E	1394	ALA	3.5
1	F	1399	GLN	3.5
1	E	1401	GLN	3.4
1	D	1381	MET	3.3
1	F	1397	GLU	3.3
1	E	1403	ALA	3.2
1	E	1397	GLU	3.1
1	F	1398	LYS	3.0
1	D	1414	GLU	2.9
1	B	1380	HIS	2.9
1	F	1406	HIS	2.9
1	C	1387	GLU	2.8
1	D	1380	HIS	2.7
1	F	1390	ALA	2.5
1	D	1418	LYS	2.5
1	E	1398	LYS	2.5
1	F	1412	GLN	2.4
1	C	1384	GLU	2.3
1	E	1391	GLU	2.3
1	F	1393	GLU	2.3
1	F	1395	ALA	2.2
1	C	1383	ALA	2.2
1	E	1409	ALA	2.2
1	E	1399	GLN	2.2
1	F	1391	GLU	2.1
1	D	1415	ARG	2.1
1	E	1406	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.