



wwPDB X-ray Structure Validation Summary Report ⓘ

May 23, 2020 – 04:25 am BST

PDB ID : 1GFF
Title : THE ATOMIC STRUCTURE OF THE DEGRADED PROCAPSID PARTICLE OF THE BACTERIOPHAGE G4: INDUCED STRUCTURAL CHANGES IN THE PRESENCE OF CALCIUM IONS AND FUNCTIONAL IMPLICATIONS
Authors : Rossmann, M.G.
Deposited on : 1995-11-06
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	1	426	
2	2	177	
3	3	25	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4779 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BACTERIOPHAGE G4 CAPSID PROTEINS GPF, GPG, GPJ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	1	417	Total	C	N	O	S	0	0	0
			3357	2144	571	623	19			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
1	378	ASP	GLU	CONFLICT	UNP P03642

- Molecule 2 is a protein called BACTERIOPHAGE G4 CAPSID PROTEINS GPF, GPG, GPJ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	2	177	Total	C	N	O	S	0	0	0
			1325	840	229	252	4			

- Molecule 3 is a protein called BACTERIOPHAGE G4 CAPSID PROTEINS GPF, GPG, GPJ.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	3	12	Total	C	N	O	0	0	0
			97	64	17	16			

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: BACTERIOPHAGE G4 CAPSID PROTEINS GPF, GPG, GPJ



- Molecule 2: BACTERIOPHAGE G4 CAPSID PROTEINS GPF, GPG, GPJ



● Molecule 3: BACTERIOPHAGE G4 CAPSID PROTEINS GPF, GPG, GPJ



MET	LYS	LYS	SER	ILE	ARG	ARG	SER	GLY	GLY	LYS	SER	LYS	G14	A15	R16	L17	W18	Y19	V20	G21	Q24	Y25
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	414.20 Å 414.20 Å 263.00 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	6.00 – 3.00	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-3.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	unknown	Depositor
R, R_{free}	0.352 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4779	wwPDB-VP
Average B, all atoms (Å ²)	0.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	1	0.93	1/3461 (0.0%)	1.30	40/4713 (0.8%)
2	2	0.90	0/1354	1.40	10/1859 (0.5%)
3	3	1.23	0/100	1.47	1/135 (0.7%)
All	All	0.93	1/4915 (0.0%)	1.33	51/6707 (0.8%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	1	43	CYS	CB-SG	5.03	1.90	1.82

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	1	214	ARG	NE-CZ-NH2	7.91	124.26	120.30
1	1	357	ARG	NE-CZ-NH2	7.81	124.20	120.30
1	1	55	ARG	NE-CZ-NH2	7.78	124.19	120.30
2	2	61	ARG	NE-CZ-NH2	7.76	124.18	120.30
1	1	208	ARG	NE-CZ-NH2	7.75	124.17	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	3357	0	3220	549	627

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	2	1325	0	1334	349	817
3	3	97	0	89	41	70
All	All	4779	0	4643	895	1437

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 95.

The worst 5 of 895 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:2:18:LEU:CB	2:2:43:ILE:HG22	1.19	1.64
2:2:18:LEU:HD23	2:2:41:ILE:CD1	1.26	1.63
1:1:138:PRO:HG2	3:3:20:VAL:CG1	1.20	1.60
1:1:138:PRO:CG	3:3:20:VAL:HG12	1.13	1.58
2:2:18:LEU:CD2	2:2:41:ILE:HD11	1.32	1.51

The worst 5 of 1437 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:1:112:ILE:CA	1:1:201:LYS:CD[2_555]	0.12	2.08
1:1:91:ALA:CA	1:1:141:ASP:CA[2_555]	0.25	1.95
1:1:349:GLN:CA	1:1:423:ILE:CG1[3_555]	0.27	1.93
2:2:8:LYS:O	2:2:89:PHE:CZ[8_556]	0.30	1.90
1:1:74:ARG:CZ	1:1:231:ASP:N[2_555]	0.30	1.90

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	1	415/426 (97%)	362 (87%)	40 (10%)	13 (3%)	4 23

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	2	175/177 (99%)	137 (78%)	30 (17%)	8 (5%)	2	14
3	3	10/25 (40%)	7 (70%)	3 (30%)	0	100	100
All	All	600/628 (96%)	506 (84%)	73 (12%)	21 (4%)	3	20

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	1	114	SER
1	1	377	THR
1	1	417	PRO
2	2	44	ASN
2	2	166	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	369/377 (98%)	308 (84%)	61 (16%)	2	11
2	2	148/148 (100%)	106 (72%)	42 (28%)	0	2
3	3	8/19 (42%)	6 (75%)	2 (25%)	0	3
All	All	525/544 (96%)	420 (80%)	105 (20%)	1	7

5 of 105 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	1	393	ILE
1	1	423	ILE
2	2	156	THR
1	1	396	SER
1	1	411	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	1	300	HIS
1	1	386	ASN
2	2	118	ASN
1	1	349	GLN
1	1	129	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.