



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 14, 2020 – 11:35 pm BST

PDB ID : 3GF5
Title : Crystal structure of the P21 R1-R7 N-terminal domain of murine MVP
Authors : Querol-Audi, J.; Casanas, A.; Luque, D.; Caston, J.R.; Fita, I.; Verdaguer, N.
Deposited on : 2009-02-26
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	387	<div> <div>8%</div> <div>82%</div> <div>14%</div> <div>• •</div> </div>
1	B	387	<div> <div>7%</div> <div>84%</div> <div>11%</div> <div>• •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	B	384	-	-	-	X
2	GOL	B	387	-	-	-	X
2	GOL	B	388	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6095 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major vault protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	378	Total	C	N	O	S	0	0	0
			2998	1905	531	555	7			
1	B	380	Total	C	N	O	S	0	0	0
			3010	1911	533	559	7			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	EXPRESSION TAG	UNP Q9EQK5
A	-2	PRO	-	EXPRESSION TAG	UNP Q9EQK5
A	-1	LEU	-	EXPRESSION TAG	UNP Q9EQK5
A	0	GLY	-	EXPRESSION TAG	UNP Q9EQK5
B	-3	GLY	-	EXPRESSION TAG	UNP Q9EQK5
B	-2	PRO	-	EXPRESSION TAG	UNP Q9EQK5
B	-1	LEU	-	EXPRESSION TAG	UNP Q9EQK5
B	0	GLY	-	EXPRESSION TAG	UNP Q9EQK5

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	13	Total	O	0	0
			13	13		
3	B	32	Total	O	0	0
			32	32		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Major vault protein



• Molecule 1: Major vault protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	36.52Å 98.89Å 140.58Å 90.00° 97.18° 90.00°	Depositor
Resolution (Å)	30.00 – 2.50 29.80 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.9 (30.00-2.50) 96.9 (29.80-2.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.38 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.232 , 0.267 0.238 , 0.271	Depositor DCC
R_{free} test set	1677 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	57.3	Xtriage
Anisotropy	0.149	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.044 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6095	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	3/3059 (0.1%)	0.49	1/4158 (0.0%)
1	B	0.38	3/3071 (0.1%)	0.51	1/4174 (0.0%)
All	All	0.38	6/6130 (0.1%)	0.50	2/8332 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	50	MET	CG-SD	6.16	1.97	1.81
1	B	138	MET	CG-SD	5.88	1.96	1.81
1	A	50	MET	CG-SD	5.59	1.95	1.81
1	A	138	MET	CG-SD	5.49	1.95	1.81
1	A	1	MET	CG-SD	5.30	1.95	1.81
1	B	1	MET	CG-SD	5.29	1.94	1.81

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	-2	PRO	N-CA-CB	6.77	111.43	103.30
1	B	-2	PRO	N-CA-CB	5.57	109.98	103.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	303	LYS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2998	0	3013	45	0
1	B	3010	0	3012	50	0
2	A	12	0	16	0	0
2	B	30	0	40	3	0
3	A	13	0	0	0	0
3	B	32	0	0	0	0
All	All	6095	0	6081	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:76:ASP:HA	1:B:82:ARG:NE	1.33	1.40
1:A:76:ASP:OD1	1:A:82:ARG:HG2	1.30	1.26
1:A:76:ASP:HA	1:A:82:ARG:NE	1.58	1.19
1:B:76:ASP:OD2	1:B:82:ARG:HG2	1.44	1.16
1:B:76:ASP:CA	1:B:82:ARG:HE	1.62	1.12
1:A:76:ASP:OD1	1:A:82:ARG:CG	1.99	1.09
1:A:76:ASP:HA	1:A:82:ARG:HE	1.01	1.08
1:B:76:ASP:HA	1:B:82:ARG:CZ	1.91	1.00
1:B:76:ASP:CA	1:B:82:ARG:NE	2.20	0.96
1:B:77:VAL:N	1:B:82:ARG:HH21	1.65	0.95
1:A:76:ASP:CA	1:A:82:ARG:HE	1.82	0.93
1:B:77:VAL:H	1:B:82:ARG:HH21	0.93	0.92
1:B:77:VAL:H	1:B:82:ARG:NH2	1.70	0.90
1:B:76:ASP:CG	1:B:82:ARG:HE	1.74	0.89
1:B:76:ASP:CB	1:B:82:ARG:HE	1.87	0.88
1:B:76:ASP:HA	1:B:82:ARG:HE	1.04	0.87
1:B:76:ASP:CG	1:B:82:ARG:HG2	1.97	0.84

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:76:ASP:OD2	1:B:82:ARG:CG	2.28	0.77
1:A:76:ASP:HA	1:A:82:ARG:CZ	2.18	0.73
1:B:76:ASP:OD1	1:B:82:ARG:NE	2.22	0.72
1:B:77:VAL:N	1:B:82:ARG:NH2	2.32	0.72
1:B:334:LEU:HD11	1:B:359:ILE:HD13	1.72	0.72
1:B:76:ASP:CG	1:B:82:ARG:NE	2.42	0.71
1:B:76:ASP:CG	1:B:82:ARG:CG	2.59	0.69
1:A:219:VAL:HG11	1:A:227:LEU:HD13	1.77	0.66
1:A:77:VAL:H	1:A:82:ARG:HH21	1.43	0.66
1:A:8:ILE:CD1	1:A:43:VAL:HG21	2.29	0.62
1:B:78:THR:HG23	1:B:80:GLN:H	1.63	0.62
1:B:120:ALA:HB2	1:B:205:LEU:HD21	1.82	0.60
1:B:77:VAL:HG13	1:B:82:ARG:HH22	1.68	0.59
1:A:76:ASP:OD1	1:A:82:ARG:CD	2.54	0.56
1:A:76:ASP:OD1	1:A:82:ARG:NE	2.39	0.55
1:A:285:LEU:HD13	1:A:317:GLU:HB2	1.89	0.55
1:B:288:MET:O	1:B:289:GLY:O	2.25	0.55
1:B:35:TYR:OH	1:B:41:GLU:OE1	2.22	0.55
1:B:208:VAL:HG13	1:B:209:PHE:CD2	2.43	0.54
1:B:310:LEU:HD21	1:B:316:LEU:CD1	2.38	0.54
1:A:323:VAL:HG22	1:A:366:VAL:CG1	2.37	0.53
1:B:325:VAL:HG13	1:B:325:VAL:O	2.09	0.53
1:A:144:LEU:HD13	1:A:208:VAL:HB	1.91	0.52
1:B:76:ASP:N	1:B:82:ARG:HG3	2.24	0.52
1:A:77:VAL:N	1:A:82:ARG:HH21	2.08	0.52
1:A:8:ILE:HD11	1:A:43:VAL:HG21	1.91	0.52
1:B:241:VAL:HG11	2:B:388:GOL:H2	1.92	0.51
1:A:8:ILE:HD12	1:A:43:VAL:HG21	1.92	0.51
1:A:310:LEU:HD21	1:A:316:LEU:HD13	1.92	0.50
1:B:26:SER:HB2	1:B:80:GLN:HE22	1.77	0.49
1:A:288:MET:HE2	1:A:311:GLN:HG3	1.93	0.49
1:A:6:ALA:HB3	1:A:7:ILE:HD12	1.95	0.49
1:A:61:VAL:HG21	1:A:91:ARG:NH1	2.27	0.49
1:A:85:HIS:O	1:A:86:ALA:HB3	2.12	0.48
1:A:222:THR:HG22	1:A:223:GLU:H	1.78	0.48
1:B:76:ASP:HA	1:B:82:ARG:NH2	2.27	0.48
1:A:30:VAL:HG13	1:A:50:MET:CE	2.43	0.48
1:A:334:LEU:HD11	1:A:359:ILE:HD12	1.95	0.47
1:B:323:VAL:HG22	1:B:366:VAL:HG22	1.95	0.47
1:B:241:VAL:HG11	2:B:388:GOL:C1	2.44	0.47
1:B:68:ASP:HB3	1:B:74:LEU:HD21	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:77:VAL:HG13	1:B:82:ARG:NH2	2.30	0.47
1:B:241:VAL:HG11	2:B:388:GOL:C2	2.44	0.47
1:B:310:LEU:HD21	1:B:316:LEU:HD13	1.97	0.46
1:A:175:ARG:NH2	1:A:263:VAL:HG23	2.30	0.46
1:A:323:VAL:HG13	1:A:366:VAL:HG13	1.97	0.46
1:B:348:ARG:O	1:B:349:VAL:CB	2.64	0.46
1:A:173:ALA:HB3	1:A:260:VAL:HG11	1.98	0.45
1:A:65:VAL:HG12	1:A:83:LEU:HD23	1.98	0.45
1:B:220:ILE:N	1:B:220:ILE:HD12	2.31	0.45
1:A:237:ASP:OD2	1:A:241:VAL:HG12	2.17	0.45
1:B:326:LEU:HD21	1:B:332:LEU:HG	1.99	0.44
1:A:76:ASP:CA	1:A:82:ARG:NE	2.50	0.43
1:A:1:MET:HB2	1:A:46:ALA:HB3	2.01	0.43
1:B:326:LEU:HD11	1:B:332:LEU:HG	1.99	0.43
1:B:227:LEU:HD21	1:B:229:LEU:HD21	2.00	0.43
1:B:290:PRO:O	1:B:291:ASP:C	2.56	0.43
1:B:59:CYS:HB3	1:B:109:ILE:HD13	2.01	0.42
1:B:276:LEU:O	1:B:305:GLU:HA	2.19	0.42
1:A:197:LEU:HD22	1:A:262:ASP:HA	2.00	0.42
1:A:76:ASP:N	1:A:82:ARG:HG3	2.33	0.42
1:A:220:ILE:N	1:A:220:ILE:HD12	2.35	0.42
1:A:76:ASP:HA	1:A:82:ARG:NH2	2.34	0.42
1:B:76:ASP:HB3	1:B:78:THR:HG22	2.02	0.42
1:B:164:GLN:HE21	1:B:164:GLN:HA	1.84	0.42
1:A:62:ALA:HB3	1:A:104:LEU:CD1	2.50	0.42
1:A:120:ALA:HB2	1:A:205:LEU:HD21	2.02	0.41
1:A:83:LEU:HD11	1:A:102:GLY:HA3	2.02	0.41
1:A:81:VAL:HG21	1:A:100:TYR:HB3	2.02	0.41
1:A:221:LEU:HD21	1:A:227:LEU:HB2	2.03	0.41
1:B:127:LEU:HD13	1:B:128:ASP:O	2.21	0.41
1:A:73:VAL:HG21	1:A:100:TYR:CZ	2.55	0.41
1:A:122:HIS:CD2	1:A:208:VAL:HG23	2.55	0.41
1:A:326:LEU:HD11	1:A:332:LEU:HD13	2.02	0.41
1:B:83:LEU:HD11	1:B:101:PRO:O	2.21	0.41
1:B:85:HIS:O	1:B:86:ALA:HB3	2.20	0.41
1:A:62:ALA:HB2	1:A:106:GLU:HB2	2.03	0.41
1:B:303:LYS:HG2	1:B:304:GLY:N	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/387 (97%)	358 (96%)	14 (4%)	2 (0%)	29	48
1	B	376/387 (97%)	355 (94%)	14 (4%)	7 (2%)	8	13
All	All	750/774 (97%)	713 (95%)	28 (4%)	9 (1%)	13	24

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	-2	PRO
1	B	289	GLY
1	B	290	PRO
1	A	353	ALA
1	B	328	GLU
1	B	348	ARG
1	B	349	VAL
1	B	291	ASP
1	B	304	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	324/335 (97%)	318 (98%)	6 (2%)	57	80
1	B	324/335 (97%)	317 (98%)	7 (2%)	52	77
All	All	648/670 (97%)	635 (98%)	13 (2%)	55	79

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	ASP
1	A	87	ASP
1	A	127	LEU
1	A	222	THR
1	A	285	LEU
1	A	316	LEU
1	B	127	LEU
1	B	164	GLN
1	B	223	GLU
1	B	241	VAL
1	B	291	ASP
1	B	316	LEU
1	B	363	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	GLN
1	A	295	GLN
1	A	338	GLN
1	B	22	ASN
1	B	40	ASN
1	B	164	GLN
1	B	228	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	387	-	5,5,5	0.39	0	5,5,5	0.21	0
2	GOL	B	388	-	5,5,5	0.35	0	5,5,5	0.28	0
2	GOL	A	385	-	5,5,5	0.36	0	5,5,5	0.26	0
2	GOL	B	386	-	5,5,5	0.38	0	5,5,5	0.24	0
2	GOL	B	385	-	5,5,5	0.42	0	5,5,5	0.14	0
2	GOL	A	384	-	5,5,5	0.37	0	5,5,5	0.14	0
2	GOL	B	384	-	5,5,5	0.36	0	5,5,5	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	387	-	-	1/4/4/4	-
2	GOL	B	388	-	-	2/4/4/4	-
2	GOL	A	385	-	-	0/4/4/4	-
2	GOL	B	386	-	-	1/4/4/4	-
2	GOL	B	385	-	-	3/4/4/4	-
2	GOL	A	384	-	-	2/4/4/4	-
2	GOL	B	384	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	388	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	385	GOL	C1-C2-C3-O3
2	A	384	GOL	C1-C2-C3-O3
2	B	388	GOL	O1-C1-C2-O2
2	B	385	GOL	O1-C1-C2-C3
2	B	386	GOL	O2-C2-C3-O3
2	B	385	GOL	O2-C2-C3-O3
2	B	384	GOL	O1-C1-C2-C3
2	B	387	GOL	O1-C1-C2-C3
2	A	384	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	388	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	378/387 (97%)	0.86	32 (8%) 10 10	23, 53, 55, 59	0
1	B	380/387 (98%)	0.84	28 (7%) 14 15	23, 52, 56, 59	0
All	All	758/774 (97%)	0.85	60 (7%) 12 12	23, 53, 56, 59	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	77	VAL	8.1
1	B	347	GLU	7.4
1	B	343	GLY	6.8
1	B	329	GLN	6.5
1	A	379	ALA	6.2
1	A	82	ARG	6.1
1	A	77	VAL	5.7
1	A	292	GLY	5.6
1	B	82	ARG	5.3
1	B	349	VAL	5.1
1	A	347	GLU	4.2
1	B	353	ALA	4.2
1	A	348	ARG	4.1
1	A	293	LYS	3.9
1	B	22	ASN	3.8
1	B	201	VAL	3.7
1	B	350	ALA	3.5
1	A	289	GLY	3.5
1	B	325	VAL	3.5
1	A	25	VAL	3.4
1	B	76	ASP	3.3
1	B	348	ARG	3.2
1	A	338	GLN	3.0
1	B	68	ASP	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	350	ALA	2.9
1	B	379	ALA	2.8
1	B	328	GLU	2.8
1	A	66	SER	2.8
1	A	349	VAL	2.7
1	A	-2	PRO	2.7
1	A	290	PRO	2.6
1	A	78	THR	2.6
1	A	352	GLN	2.6
1	A	-3	GLY	2.5
1	B	-2	PRO	2.4
1	A	9	ARG	2.4
1	A	65	VAL	2.4
1	A	304	GLY	2.4
1	B	245	THR	2.3
1	A	360	ARG	2.3
1	A	134	GLY	2.3
1	A	370	LYS	2.2
1	A	72	SER	2.2
1	B	257	GLU	2.2
1	A	291	ASP	2.2
1	A	296	LEU	2.1
1	B	276	LEU	2.1
1	A	323	VAL	2.1
1	B	362	PRO	2.1
1	B	289	GLY	2.1
1	A	333	LEU	2.1
1	B	290	PRO	2.1
1	A	327	SER	2.1
1	A	6	ALA	2.0
1	B	70	GLN	2.0
1	B	171	ASN	2.0
1	B	54	PRO	2.0
1	B	341	GLU	2.0
1	A	353	ALA	2.0
1	B	69	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	B	387	6/6	0.44	0.45	84,85,85,85	0
2	GOL	B	386	6/6	0.51	0.26	86,86,86,86	0
2	GOL	A	384	6/6	0.61	0.30	84,84,84,84	0
2	GOL	B	385	6/6	0.65	0.25	79,79,79,79	0
2	GOL	B	384	6/6	0.68	0.46	97,97,97,98	0
2	GOL	B	388	6/6	0.71	0.41	89,90,90,90	0
2	GOL	A	385	6/6	0.81	0.34	77,78,78,78	0

6.5 Other polymers [i](#)

There are no such residues in this entry.