



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 13, 2020 – 07:04 pm BST

PDB ID : 3GG1
Title : Klebsiella pneumoniae BlrP1 pH 8.0 calcium/cy-diGMP complex
Authors : Barends, T.; Hartmann, E.; Griesse, J.; Beitlich, T.; Kirienko, N.; Ryjenkov, D.; Reinstein, J.; Shoeman, R.; Gomelsky, M.; Schlichting, I.
Deposited on : 2009-02-27
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	413	
1	B	413	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	C2E	A	501	X	-	-	-
2	C2E	B	501	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6500 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called *Klebsiella pneumoniae* Blrp1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	392	Total	C	N	O	S	0	0	0
			3069	1967	529	555	18			
1	B	394	Total	C	N	O	S	0	0	0
			3094	1983	534	559	18			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	ILE	-	EXPRESSION TAG	UNP A6T8V8
A	-6	SER	-	EXPRESSION TAG	UNP A6T8V8
A	-5	GLU	-	EXPRESSION TAG	UNP A6T8V8
A	-4	PHE	-	EXPRESSION TAG	UNP A6T8V8
A	-3	GLY	-	EXPRESSION TAG	UNP A6T8V8
A	-2	SER	-	EXPRESSION TAG	UNP A6T8V8
A	-1	SER	-	EXPRESSION TAG	UNP A6T8V8
A	0	ARG	-	EXPRESSION TAG	UNP A6T8V8
B	-7	ILE	-	EXPRESSION TAG	UNP A6T8V8
B	-6	SER	-	EXPRESSION TAG	UNP A6T8V8
B	-5	GLU	-	EXPRESSION TAG	UNP A6T8V8
B	-4	PHE	-	EXPRESSION TAG	UNP A6T8V8
B	-3	GLY	-	EXPRESSION TAG	UNP A6T8V8
B	-2	SER	-	EXPRESSION TAG	UNP A6T8V8
B	-1	SER	-	EXPRESSION TAG	UNP A6T8V8
B	0	ARG	-	EXPRESSION TAG	UNP A6T8V8

- Molecule 2 is 9,9'-[(2R,3R,3aS,5S,7aR,9R,10R,10aS,12S,14aR)-3,5,10,12-tetrahydroxy-5,12-dioxidoctahydro-2H,7H-difuro[3,2-d:3',2'-j][1,3,7,9,2,8]tetraoxadiphosphacyclododecine-2,9-diyl]bis(2-amino-1,9-dihydro-6H-purin-6-one) (three-letter code: C2E) (formula: C₂₀H₂₄N₁₀O₁₄P₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total 46	C 20	N 10	O 14	P 2	0	0
2	B	1	Total 46	C 20	N 10	O 14	P 2	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Ca 2 2	0	0
3	A	2	Total Ca 2 2	0	0


- Molecule 4 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: $\text{C}_{17}\text{H}_{21}\text{N}_4\text{O}_9\text{P}$).




3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: *Klebsiella pneumoniae* Blrp1



• Molecule 1: *Klebsiella pneumoniae* Blrp1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.85Å 96.69Å 127.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 19.81 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-2.30) 100.0 (19.81-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.90 (at 2.30Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.244 , 0.296 0.248 , 0.297	Depositor DCC
R_{free} test set	1947 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	27.1	Xtriage
Anisotropy	0.554	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6500	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, CA, C2E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/3137	0.55	2/4252 (0.0%)
1	B	0.31	0/3163	0.49	0/4287
All	All	0.31	0/6300	0.52	2/8539 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	238	ILE	N-CA-C	7.62	131.57	111.00
1	A	317	PHE	N-CA-C	5.45	125.71	111.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	237	ALA	Peptide
1	A	309	SER	Peptide
1	A	316	ARG	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3069	0	3087	15	0
1	B	3094	0	3110	14	0
2	A	46	0	18	1	0
2	B	46	0	18	1	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	31	0	19	0	0
4	B	31	0	19	0	0
5	A	93	0	0	0	0
5	B	86	0	0	0	0
All	All	6500	0	6271	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:24:VAL:HG21	1:A:94:ILE:HB	1.83	0.60
1:B:161:PRO:HG3	1:B:387:CYS:SG	2.43	0.59
1:A:308:TYR:N	1:A:309:SER:HB3	2.18	0.58
1:B:246:TYR:CE2	1:B:284:GLN:HB2	2.39	0.56
1:B:68:ASP:OD1	1:B:70:ARG:HD3	2.09	0.53
1:A:355:THR:HG23	1:A:377:LEU:HD12	1.92	0.51
1:B:314:LEU:O	1:B:318:GLN:NE2	2.41	0.51
1:B:256:MET:HE2	1:B:266:PRO:HG3	1.92	0.50
1:B:174:GLN:OE1	2:B:501:C2E:H8	2.12	0.50
1:A:308:TYR:CA	1:A:309:SER:HB3	2.43	0.49
1:A:270:LEU:HD21	1:A:300:ALA:HB2	1.95	0.48
1:A:270:LEU:HD21	1:A:300:ALA:CB	2.44	0.48
1:A:186:SER:HB3	1:A:235:GLN:HB2	1.96	0.47
1:A:308:TYR:HA	1:A:309:SER:CB	2.46	0.46
1:B:32:LEU:HD13	1:B:90:THR:HB	1.99	0.45
1:B:32:LEU:HB3	1:B:33:PRO:HD3	1.99	0.45
1:A:308:TYR:HA	1:A:309:SER:HB3	1.98	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:308:TYR:CA	1:A:309:SER:CB	2.95	0.44
1:B:217:GLU:HA	1:B:254:TRP:CH2	2.52	0.43
1:B:256:MET:SD	1:B:295:ALA:HB1	2.59	0.43
1:A:292:LEU:CD2	1:A:297:MET:HE3	2.49	0.43
1:B:160:ALA:N	1:B:161:PRO:CD	2.81	0.43
1:A:355:THR:CG2	1:A:377:LEU:HD12	2.49	0.42
1:B:186:SER:HB3	1:B:235:GLN:HB3	2.03	0.41
1:B:256:MET:HE3	1:B:256:MET:HA	2.01	0.41
1:A:174:GLN:OE1	2:A:501:C2E:H8	2.20	0.41
1:A:195:THR:HG22	1:A:195:THR:O	2.21	0.40
1:A:143:GLU:N	1:A:144:PRO:CD	2.84	0.40
1:B:31:ASN:HB3	1:B:36:ILE:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/413 (94%)	378 (98%)	7 (2%)	1 (0%)	41	50
1	B	390/413 (94%)	378 (97%)	11 (3%)	1 (0%)	41	50
All	All	776/826 (94%)	756 (97%)	18 (2%)	2 (0%)	41	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	309	SER
1	B	102	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	329/346 (95%)	322 (98%)	7 (2%)	53	70
1	B	332/346 (96%)	321 (97%)	11 (3%)	38	53
All	All	661/692 (96%)	643 (97%)	18 (3%)	44	61

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	LEU
1	A	70	ARG
1	A	71	HIS
1	A	153	MET
1	A	173	LEU
1	A	276	THR
1	A	287	LYS
1	B	10	GLN
1	B	25	HIS
1	B	56	GLU
1	B	71	HIS
1	B	80	ASP
1	B	97	LEU
1	B	103	ASP
1	B	112	PHE
1	B	231	LEU
1	B	279	ILE
1	B	317	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	10	GLN
1	B	12	HIS
1	B	66	GLN
1	B	88	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	268	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	C2E	A	501	3	44,52,52	3.20	8 (18%)	54,82,82	2.09	18 (33%)
4	FMN	A	504	-	31,33,33	1.41	4 (12%)	40,50,50	1.65	7 (17%)
2	C2E	B	501	3	44,52,52	3.19	8 (18%)	54,82,82	2.10	19 (35%)
4	FMN	B	504	-	31,33,33	1.47	4 (12%)	40,50,50	1.59	6 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	C2E	A	501	3	1/1/10/10	9/22/62/62	0/6/7/7
4	FMN	A	504	-	-	1/18/18/18	0/3/3/3
2	C2E	B	501	3	1/1/10/10	6/22/62/62	0/6/7/7
4	FMN	B	504	-	-	1/18/18/18	0/3/3/3

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	C2E	C2A-C1A	-13.57	1.33	1.53
2	A	501	C2E	C2'-C1'	-13.52	1.33	1.53
2	A	501	C2E	C2A-C1A	-13.44	1.33	1.53
2	B	501	C2E	C2'-C1'	-13.09	1.33	1.53
4	B	504	FMN	C10-N1	4.21	1.38	1.33
2	A	501	C2E	O2'-C2'	-4.16	1.33	1.43
2	B	501	C2E	O2'-C2'	-4.16	1.33	1.43
2	B	501	C2E	O2A-C2A	-3.93	1.33	1.43
2	A	501	C2E	O2A-C2A	-3.81	1.34	1.43
4	A	504	FMN	C4A-N5	3.74	1.38	1.33
4	A	504	FMN	C10-N1	3.60	1.37	1.33
2	A	501	C2E	O4'-C1'	3.59	1.46	1.41
2	B	501	C2E	O4'-C1'	3.58	1.46	1.41
4	B	504	FMN	C4A-N5	3.52	1.38	1.33
4	B	504	FMN	C4-N3	3.39	1.38	1.33
2	B	501	C2E	O4A-C1A	3.19	1.45	1.41
4	A	504	FMN	C4-N3	3.17	1.38	1.33
2	A	501	C2E	C8-N7	-3.12	1.29	1.34
2	B	501	C2E	C8-N7	-3.05	1.29	1.34
4	A	504	FMN	C1'-N10	2.97	1.51	1.48
4	B	504	FMN	C1'-N10	2.94	1.51	1.48
2	A	501	C2E	O4A-C1A	2.76	1.44	1.41
2	B	501	C2E	P11-O11	2.70	1.60	1.50
2	A	501	C2E	P11-O11	2.64	1.60	1.50

All (50) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	504	FMN	C4-N3-C2	5.34	119.65	115.14
4	B	504	FMN	C4-N3-C2	5.32	119.64	115.14
4	A	504	FMN	C1'-N10-C9A	5.25	122.42	118.29
2	B	501	C2E	O4A-C1A-C2A	4.90	114.08	106.93
2	A	501	C2E	N31-C21-N11	-4.63	121.04	127.22
2	B	501	C2E	N31-C21-N11	-4.59	121.10	127.22
2	B	501	C2E	N3-C2-N1	-4.54	121.17	127.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	C2E	N3-C2-N1	-4.53	121.17	127.22
2	B	501	C2E	C4-C5-N7	-4.44	104.77	109.40
2	A	501	C2E	O4A-C1A-C2A	4.19	113.05	106.93
2	A	501	C2E	C4-C5-N7	-4.10	105.12	109.40
4	B	504	FMN	C1'-N10-C9A	4.02	121.46	118.29
2	B	501	C2E	C3'-C2'-C1'	3.77	108.24	99.89
2	A	501	C2E	C3'-C2'-C1'	3.75	108.19	99.89
4	B	504	FMN	C4A-N5-C5A	3.65	120.42	116.77
2	B	501	C2E	C5-C6-N1	-3.62	118.47	123.43
2	A	501	C2E	C5-C6-N1	-3.56	118.57	123.43
2	A	501	C2E	C21-N31-C41	3.55	119.42	115.36
2	A	501	C2E	C6-N1-C2	3.51	121.51	115.93
2	B	501	C2E	C2-N3-C4	3.50	119.35	115.36
4	A	504	FMN	C4A-N5-C5A	3.49	120.26	116.77
2	B	501	C2E	C6-N1-C2	3.42	121.36	115.93
2	A	501	C2E	C51-C61-N11	-3.32	118.90	123.43
2	A	501	C2E	C61-N11-C21	3.32	121.20	115.93
2	B	501	C2E	C21-N31-C41	3.30	119.12	115.36
2	B	501	C2E	C61-N11-C21	3.24	121.08	115.93
2	A	501	C2E	C2-N3-C4	3.19	119.00	115.36
2	B	501	C2E	C51-C61-N11	-3.17	119.09	123.43
2	B	501	C2E	O2'-C2'-C1'	3.10	122.32	110.85
2	B	501	C2E	O5A-C5A-C4A	-3.08	98.38	108.99
2	A	501	C2E	O2'-C2'-C1'	3.08	122.22	110.85
2	A	501	C2E	O21-P11-O5A	2.99	121.65	107.75
2	A	501	C2E	O2'-C2'-C3'	2.81	119.14	111.17
4	B	504	FMN	C5A-C9A-N10	2.76	119.72	117.72
4	A	504	FMN	C10-C4A-N5	-2.66	119.42	121.26
2	B	501	C2E	O2'-C2'-C3'	2.65	118.70	111.17
2	A	501	C2E	O5A-C5A-C4A	-2.62	99.99	108.99
2	A	501	C2E	O4'-C1'-C2'	2.61	110.75	106.93
4	B	504	FMN	C4A-C4-N3	-2.57	119.92	123.43
4	B	504	FMN	C10-C4A-N5	-2.45	119.56	121.26
4	A	504	FMN	C4A-C4-N3	-2.31	120.27	123.43
4	A	504	FMN	C4-C4A-N5	2.30	121.23	118.60
4	A	504	FMN	C5A-C9A-N10	2.29	119.38	117.72
2	B	501	C2E	O21-P11-O5A	2.27	118.29	107.75
2	B	501	C2E	O5A-P11-O11	-2.15	100.68	109.07
2	B	501	C2E	O4'-C1'-C2'	2.11	110.01	106.93
2	A	501	C2E	C61-C51-C41	-2.11	118.79	120.80
2	B	501	C2E	C61-C51-C41	-2.08	118.81	120.80
2	A	501	C2E	O5A-P11-O11	-2.02	101.16	109.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	C2E	C1'-N9-C4	-2.01	123.11	126.64

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	501	C2E	C1A
2	B	501	C2E	C1A

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	C2E	C5'-O5'-P1-O1P
2	B	501	C2E	C5'-O5'-P1-O1P
4	B	504	FMN	C5'-O5'-P-O2P
2	A	501	C2E	C2A-C3A-O3A-P1
2	B	501	C2E	C5'-O5'-P1-O3A
2	A	501	C2E	C4A-C3A-O3A-P1
2	A	501	C2E	C4'-C5'-O5'-P1
2	B	501	C2E	C4'-C5'-O5'-P1
2	B	501	C2E	C5'-O5'-P1-O2P
2	A	501	C2E	C5'-O5'-P1-O3A
2	A	501	C2E	O4'-C4'-C5'-O5'
2	B	501	C2E	O4'-C4'-C5'-O5'
2	A	501	C2E	C3'-C4'-C5'-O5'
2	B	501	C2E	C3'-C4'-C5'-O5'
4	A	504	FMN	C4'-C5'-O5'-P
2	A	501	C2E	O4A-C4A-C5A-O5A
2	A	501	C2E	C5'-O5'-P1-O2P


There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	C2E	1	0
2	B	501	C2E	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	392/413 (94%)	0.53	34 (8%) 10 14	20, 29, 40, 47	0
1	B	394/413 (95%)	0.69	46 (11%) 4 6	20, 29, 40, 47	0
All	All	786/826 (95%)	0.61	80 (10%) 6 9	20, 29, 40, 47	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	308	TYR	9.8
1	B	308	TYR	7.2
1	A	281	CYS	7.2
1	B	282	PHE	6.6
1	B	232	GLY	6.3
1	B	280	THR	6.2
1	A	309	SER	5.8
1	B	284	GLN	5.4
1	A	165	PRO	5.2
1	B	102	THR	5.2
1	B	276	THR	5.2
1	B	278	VAL	5.2
1	B	307	GLY	5.1
1	A	159	ALA	5.1
1	B	279	ILE	4.9
1	B	14	ASP	4.9
1	A	282	PHE	4.7
1	B	285	PHE	4.6
1	B	90	THR	4.6
1	B	231	LEU	4.2
1	B	309	SER	4.0
1	B	101	GLU	4.0
1	A	121	VAL	3.8
1	A	14	ASP	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	3.7
1	B	233	LYS	3.6
1	A	279	ILE	3.6
1	A	280	THR	3.6
1	B	277	GLU	3.5
1	A	158	THR	3.3
1	B	249	PRO	3.1
1	B	393	GLU	3.1
1	B	265	ARG	3.1
1	B	153	MET	3.0
1	A	166	GLY	2.9
1	B	237	ALA	2.9
1	B	283	ASP	2.9
1	B	238	ILE	2.8
1	A	161	PRO	2.8
1	A	26	ARG	2.8
1	B	286	ARG	2.8
1	A	286	ARG	2.8
1	A	283	ASP	2.7
1	B	25	HIS	2.7
1	B	289	LEU	2.6
1	B	313	LEU	2.6
1	A	162	GLN	2.6
1	B	281	CYS	2.6
1	B	161	PRO	2.6
1	A	163	HIS	2.6
1	A	305	GLY	2.5
1	A	155	ALA	2.5
1	B	13	PRO	2.5
1	A	149	ARG	2.4
1	B	287	LYS	2.4
1	A	306	ALA	2.4
1	B	311	LEU	2.4
1	A	316	ARG	2.4
1	A	153	MET	2.4
1	B	316	ARG	2.3
1	B	209	GLU	2.3
1	A	102	THR	2.3
1	B	230	PRO	2.3
1	B	16	PRO	2.3
1	A	160	ALA	2.3
1	A	175	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	307	GLY	2.2
1	B	399	ALA	2.2
1	B	103	ASP	2.2
1	A	247	HIS	2.2
1	B	26	ARG	2.2
1	B	100	PHE	2.1
1	A	164	LEU	2.1
1	A	195	THR	2.1
1	B	91	GLY	2.1
1	B	400	ARG	2.1
1	A	287	LYS	2.0
1	B	273	VAL	2.0
1	A	114	THR	2.0
1	B	299	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	FMN	A	504	31/31	0.86	0.17	22,23,37,37	0
4	FMN	B	504	31/31	0.88	0.14	22,23,38,38	0
3	CA	B	503	1/1	0.94	0.06	34,34,34,34	0
2	C2E	B	501	46/46	0.95	0.11	22,23,24,24	0
2	C2E	A	501	46/46	0.96	0.13	21,22,23,24	0
3	CA	A	503	1/1	0.97	0.11	30,30,30,30	0
3	CA	B	502	1/1	0.99	0.03	18,18,18,18	0
3	CA	A	502	1/1	1.00	0.04	17,17,17,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





Electron density around FMN B 504:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around C2E B 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around C2E A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.