



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 12:13 pm BST

PDB ID : 3GHB
Title : Crystal structure of anti-HIV-1 Fab 447-52D in complex with V3 peptide W2RW020
Authors : Kong, X.P.; Burke, V.J.
Deposited on : 2009-03-03
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	216	<div> <div>89%</div> <div>11%</div> </div>
1	M	216	<div>4%</div> <div>85%</div> <div>13%</div> <div>•</div>
2	H	235	<div>%</div> <div>86%</div> <div>13%</div> <div>•</div>
2	I	235	<div>2%</div> <div>79%</div> <div>20%</div> <div>•</div>
3	P	10	<div>20%</div> <div>80%</div> <div>20%</div>
3	Q	10	<div>30%</div> <div>50%</div> <div>30%</div> <div>10%</div> <div>10%</div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7431 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab 447-52D, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	216	Total	C	N	O	S	0	0	0
			1603	1007	266	325	5			
1	M	216	Total	C	N	O	S	0	0	0
			1603	1007	266	325	5			

- Molecule 2 is a protein called Fab 447-52D, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	235	Total	C	N	O	S	0	0	0
			1769	1114	296	351	8			
2	I	235	Total	C	N	O	S	0	0	0
			1769	1114	296	351	8			

- Molecule 3 is a protein called Envelope glycoprotein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	P	10	Total	C	N	O	0	0	0
			68	42	15	11			
3	Q	9	Total	C	N	O	0	0	0
			63	39	14	10			

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	H	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	194	Total	O	0	0
			194	194		
5	H	183	Total	O	0	0
			183	183		
5	P	3	Total	O	0	0
			3	3		
5	M	70	Total	O	0	0
			70	70		
5	I	100	Total	O	0	0
			100	100		
5	Q	1	Total	O	0	0
			1	1		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab 447-52D, light chain

Chain L: 





- Molecule 1: Fab 447-52D, light chain

Chain M: 





- Molecule 2: Fab 447-52D, heavy chain


Chain H: 





- Molecule 2: Fab 447-52D, heavy chain

Chain I: 






● Molecule 3: Envelope glycoprotein



● Molecule 3: Envelope glycoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.27Å 76.21Å 113.92Å 90.00° 101.25° 90.00°	Depositor
Resolution (Å)	49.26 – 2.25 49.26 – 2.24	Depositor EDS
% Data completeness (in resolution range)	96.3 (49.26-2.25) 95.5 (49.26-2.24)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.16 (at 2.24Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.196 , 0.232 0.197 , 0.199	Depositor DCC
R_{free} test set	5768 reflections (10.17%)	wwPDB-VP
Wilson B-factor (Å ²)	29.8	Xtriage
Anisotropy	0.234	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7431	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.35	0/1644	0.68	1/2247 (0.0%)
1	M	0.30	0/1644	0.59	0/2247
2	H	0.35	0/1809	0.66	0/2464
2	I	0.31	0/1809	0.62	0/2464
3	P	0.31	0/68	0.53	0/89
3	Q	0.31	0/63	0.52	0/82
All	All	0.33	0/7037	0.64	1/9593 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	96	TRP	N-CA-C	-5.37	96.51	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1603	0	1556	17	0
1	M	1603	0	1556	23	0
2	H	1769	0	1739	24	0
2	I	1769	0	1739	32	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	P	68	0	74	1	0
3	Q	63	0	69	8	0
4	H	5	0	0	0	0
5	H	183	0	0	1	0
5	I	100	0	0	1	0
5	L	194	0	0	4	0
5	M	70	0	0	2	0
5	P	3	0	0	0	0
5	Q	1	0	0	0	0
All	All	7431	0	6733	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:43:LYS:HD3	2:I:44:GLY:H	1.20	1.03
2:I:43:LYS:HD3	2:I:44:GLY:N	1.76	1.01
2:H:129:ARG:HH12	2:H:214:LYS:HD2	1.55	0.71
2:H:126:PRO:HD2	2:H:213:LEU:HD22	1.74	0.69
2:H:144:ASP:HB3	2:H:175:LEU:HD13	1.79	0.64
2:H:141:LEU:HG	2:H:143:LYS:HG3	1.80	0.62
2:I:144:ASP:HB3	2:I:175:LEU:HD13	1.82	0.62
1:M:34:LEU:HD23	1:M:49:TYR:HA	1.82	0.60
1:L:34:LEU:HD23	1:L:49:TYR:HA	1.85	0.59
2:H:150:VAL:CG1	2:H:178:LEU:HD21	2.31	0.59
2:H:129:ARG:NH1	2:H:214:LYS:HD2	2.18	0.58
1:L:189:ARG:HG3	1:L:189:ARG:HH11	1.67	0.58
1:L:18:LYS:N	1:L:18:LYS:HD2	2.18	0.57
2:I:119:PRO:HB3	2:I:145:TYR:HB3	1.87	0.56
1:M:54:ARG:HD3	1:M:62:PHE:O	2.04	0.55
2:I:33:TRP:HB2	2:I:95:ASP:HB2	1.89	0.55
1:M:4:LEU:HB2	1:M:99:GLY:HA2	1.88	0.55
3:Q:305:LYS:HD3	3:Q:305:LYS:O	2.06	0.55
2:I:183:THR:HG23	5:I:251:HOH:O	2.06	0.54
2:I:96:GLY:HA3	2:I:101:ASP:HB2	1.89	0.54
1:M:123:GLU:HG3	5:M:300:HOH:O	2.08	0.54
1:L:54:ARG:HD3	1:L:62:PHE:O	2.08	0.54
1:M:47:LEU:O	1:M:48:ILE:HG13	2.07	0.54
2:H:119:PRO:HB3	2:H:145:TYR:HB3	1.90	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:87:THR:O	2:H:88:ALA:HB2	2.08	0.53
2:I:141:LEU:HG	2:I:143:LYS:HG3	1.89	0.53
2:I:87:THR:O	2:I:88:ALA:HB2	2.09	0.53
1:L:22:SER:HB3	5:L:522:HOH:O	2.08	0.53
2:I:51:ILE:HG23	2:I:51:ILE:O	2.09	0.52
1:L:135:LEU:CD1	2:H:181:VAL:HG11	2.40	0.52
2:H:45:LEU:HD11	2:H:103:TRP:HZ3	1.73	0.52
2:I:4:LEU:HD11	2:I:94:THR:HG23	1.90	0.52
3:P:308:ARG:HB2	3:P:315:GLN:HB3	1.92	0.52
2:H:129:ARG:NH1	2:H:214:LYS:HB3	2.25	0.51
2:I:29:PHE:HB2	2:I:76:ASN:ND2	2.26	0.51
2:I:35:ASN:ND2	2:I:100(L):MET:HE2	2.26	0.51
2:H:178:LEU:HD12	2:H:178:LEU:C	2.31	0.51
2:H:171:GLN:HG2	5:H:317:HOH:O	2.12	0.50
2:I:99:MET:HA	2:I:100(G):TYR:O	2.12	0.50
1:L:123:GLU:HG3	5:L:317:HOH:O	2.12	0.50
2:I:162:GLY:O	2:I:182:VAL:HA	2.13	0.49
1:M:191:TYR:O	1:M:205:THR:HG23	2.12	0.49
1:M:91:TRP:CH2	3:Q:312:GLY:HA2	2.48	0.49
2:I:29:PHE:HB2	2:I:76:ASN:HD22	1.77	0.48
2:I:178:LEU:C	2:I:178:LEU:HD12	2.34	0.48
1:M:210:GLU:HB3	2:I:127:CYS:SG	2.53	0.48
2:I:29:PHE:CD2	2:I:76:ASN:HA	2.49	0.48
1:M:183:GLU:CD	1:M:183:GLU:H	2.16	0.48
1:L:151:ASP:OD2	1:L:188:HIS:HD2	1.97	0.48
1:L:18:LYS:HE3	5:L:284:HOH:O	2.14	0.48
1:M:95(A):SER:HB2	1:M:95(C):ASP:OD1	2.14	0.48
1:L:181:THR:OG1	1:L:184:GLN:HG3	2.14	0.47
2:H:201:LYS:NZ	2:H:204:ASN:HD21	2.11	0.47
2:I:29:PHE:CE1	2:I:34:LEU:HD11	2.51	0.46
1:M:181:THR:OG1	1:M:184:GLN:HG3	2.16	0.46
1:L:39:PHE:CE1	1:L:84:ALA:HB2	2.51	0.46
1:L:50:GLY:O	1:L:51:ASN:HB2	2.16	0.46
3:Q:315:GLN:N	3:Q:315:GLN:HE21	2.14	0.46
2:H:50:ARG:HG2	2:H:58:ASP:HB2	1.98	0.45
2:H:185:PRO:HB2	2:H:188:SER:HB3	1.99	0.45
2:H:129:ARG:HH22	2:H:214:LYS:HD2	1.82	0.45
2:H:125:ALA:HB1	2:H:213:LEU:HD22	1.99	0.45
1:M:46:LEU:HD22	2:I:100(K):TYR:CD2	2.51	0.45
1:M:150:ALA:HB2	1:M:155:VAL:HG22	1.98	0.45
1:M:210:GLU:HA	1:M:210:GLU:OE1	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:21:THR:HG22	2:I:79:TYR:CD1	2.53	0.44
2:H:1:GLU:O	2:H:26:GLY:HA3	2.18	0.44
1:M:150:ALA:O	1:M:190:SER:O	2.35	0.44
3:Q:308:ARG:CB	3:Q:315:GLN:HG2	2.48	0.44
1:L:18:LYS:HG3	5:L:284:HOH:O	2.19	0.43
2:I:97:PHE:HA	2:I:100(I):TYR:O	2.18	0.43
2:H:126:PRO:HD2	2:H:213:LEU:CD2	2.47	0.43
2:H:45:LEU:HD11	2:H:103:TRP:CZ3	2.54	0.43
1:M:156:LYS:HD3	1:M:156:LYS:N	2.34	0.43
3:Q:315:GLN:N	3:Q:315:GLN:NE2	2.66	0.43
1:M:124:GLU:HG2	1:M:129:LYS:O	2.19	0.42
2:I:6:GLU:CD	2:I:106:GLY:H	2.22	0.42
1:M:27(B):ASN:OD1	1:M:28:ILE:N	2.45	0.42
2:H:6:GLU:HA	2:H:21:THR:O	2.20	0.42
2:H:129:ARG:NH1	2:H:215:THR:O	2.53	0.42
1:L:108:GLN:HB2	1:L:109:PRO:HD2	2.02	0.42
1:M:179:SER:O	1:M:180:LEU:HD23	2.19	0.42
3:Q:315:GLN:H	3:Q:315:GLN:HE21	1.68	0.42
2:I:126:PRO:HD2	2:I:213:LEU:HG	2.02	0.41
2:I:99:MET:HE3	2:I:100(A):ARG:HE	1.86	0.41
2:I:34:LEU:HD13	2:I:78:LEU:HD13	2.01	0.41
1:L:204:LYS:HA	1:L:204:LYS:HD3	1.82	0.41
1:L:123:GLU:O	2:I:117:LYS:HE2	2.20	0.41
2:I:129:ARG:NH2	2:I:214:LYS:HD2	2.35	0.41
1:M:91:TRP:CZ3	3:Q:312:GLY:HA2	2.56	0.41
1:M:156:LYS:HD3	1:M:156:LYS:H	1.85	0.41
2:I:100(J):TYR:CE1	3:Q:308:ARG:HG3	2.56	0.41
2:H:159:LEU:HD21	2:H:182:VAL:HG21	2.02	0.40
2:I:201:LYS:HA	2:I:201:LYS:HD2	1.84	0.40
1:M:156:LYS:HG2	5:M:506:HOH:O	2.21	0.40
1:L:4:LEU:HB2	1:L:99:GLY:HA2	2.04	0.40
1:M:135:LEU:CD1	2:I:181:VAL:HG11	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	214/216 (99%)	206 (96%)	8 (4%)	0	100	100
1	M	214/216 (99%)	198 (92%)	14 (6%)	2 (1%)	17	14
2	H	233/235 (99%)	222 (95%)	11 (5%)	0	100	100
2	I	233/235 (99%)	224 (96%)	8 (3%)	1 (0%)	34	37
3	P	8/10 (80%)	8 (100%)	0	0	100	100
3	Q	7/10 (70%)	7 (100%)	0	0	100	100
All	All	909/922 (99%)	865 (95%)	41 (4%)	3 (0%)	41	46

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	M	187	SER
1	M	152	SER
2	I	76	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	180/180 (100%)	178 (99%)	2 (1%)	73	82
1	M	180/180 (100%)	177 (98%)	3 (2%)	60	71
2	H	200/200 (100%)	197 (98%)	3 (2%)	65	75
2	I	200/200 (100%)	196 (98%)	4 (2%)	55	64
3	P	6/6 (100%)	6 (100%)	0	100	100
3	Q	6/6 (100%)	5 (83%)	1 (17%)	2	0
All	All	772/772 (100%)	759 (98%)	13 (2%)	60	71

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	95(C)	ASP
1	L	189	ARG
2	H	6	GLU
2	H	45	LEU
2	H	149	PRO
1	M	95(C)	ASP
1	M	156	LYS
1	M	210	GLU
2	I	6	GLU
2	I	149	PRO
2	I	188	SER
2	I	192	GLN
3	Q	315	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	188	HIS
1	L	194	GLN
2	H	76	ASN
2	H	171	GLN
2	H	199	ASN
2	H	204	ASN
1	M	194	GLN
2	I	76	ASN
2	I	171	GLN
2	I	204	ASN
3	Q	315	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	H	218	-	4,4,4	0.28	0	6,6,6	0.06	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	L	216/216 (100%)	-0.12	1 (0%)	91 91	15, 23, 34, 57	0
1	M	216/216 (100%)	0.40	9 (4%)	36 38	25, 41, 67, 74	0
2	H	235/235 (100%)	0.03	2 (0%)	84 85	15, 25, 51, 64	0
2	I	235/235 (100%)	0.24	5 (2%)	63 66	20, 36, 64, 73	0
3	P	10/10 (100%)	1.38	2 (20%)	1 1	45, 57, 66, 69	0
3	Q	9/10 (90%)	1.31	3 (33%)	0 0	69, 74, 77, 78	0
All	All	921/922 (99%)	0.16	22 (2%)	59 62	15, 30, 60, 78	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	M	150	ALA	3.9
1	M	189	ARG	3.7
3	P	316	ALA	3.6
2	I	216	PRO	3.6
3	P	305	LYS	3.6
2	I	217	THR	3.3
1	L	1	GLN	3.0
1	M	187	SER	2.7
1	M	210	GLU	2.5
1	M	1	GLN	2.5
1	M	27	SER	2.5
1	M	207	ALA	2.5
3	Q	315	GLN	2.4
2	I	215	THR	2.4
1	M	209	THR	2.3
2	I	132	SER	2.3
3	Q	306	GLY	2.2
2	I	131	THR	2.2
1	M	18	LYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	100(C)	VAL	2.1
3	Q	312	GLY	2.1
2	H	217	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	SO4	H	218	5/5	0.98	0.16	49,50,51,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.