



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 10:50 PM BST

PDB ID : 5GH0
Title : Crystal structure of the complex of bovine lactoperoxidase with mercaptoimidazole at 2.3 Å resolution
Authors : Singh, P.K.; Sirohi, H.V.; Singh, A.K.; Bhushan, A.; Kaur, P.; Sharma, S.; Singh, T.P.
Deposited on : 2016-06-17
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.


Mol	Chain	Length	Quality of chain
1	A	595	<div> <div>5%</div> <div> <div></div> <div>87%</div> <div>10%</div> <div>.</div> </div> </div>
2	B	2	<div> <div>100%</div> </div>

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	595	Total	C	N	O	P	S	0	0	0
			4774	3037	847	863	1	26			



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
							0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O		
			14	8	1	5		
4	A	1	Total	C	N	O		
			14	8	1	5		
4	A	1	Total	C	N	O		
			14	8	1	5		
							0	0

- Molecule 5 is NITRATE ION (three-letter code: NO3) (formula: NO_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	N	O	0	0
			4	1	3		
5	A	1	Total	N	O	0	0
			4	1	3		

- Molecule 6 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	1	3		

- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Ca	0	0
			1	1		

- Molecule 8 is 1,3-dihydroimidazole-2-thione (three-letter code: MZY) (formula: $C_3H_4N_2S$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	S	0	0
			6	3	2	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	290	Total	O	0	0
			290	290		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lactoperoxidase



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.63Å 80.67Å 77.68Å 90.00° 102.60° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 24.46 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.2 (20.00-2.30) 99.3 (24.46-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.8.0151	Depositor
R, R_{free}	0.142 , 0.194 0.151 , 0.194	Depositor DCC
R_{free} test set	932 reflections (3.20%)	wwPDB-VP
Wilson B-factor (Å ²)	23.5	Xtriage
Anisotropy	0.105	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 47.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5196	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CO3, NAG, SEP, CA, MZY, HEM, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	1/4891 (0.0%)	0.92	9/6634 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	258	GLU	CD-OE1	-5.93	1.19	1.25

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	168	PRO	CB-CA-C	-8.54	90.65	112.00
1	A	11	PRO	N-CA-C	8.18	133.38	112.10
1	A	418	ARG	NE-CZ-NH1	7.54	124.07	120.30
1	A	172	TYR	N-CA-CB	-6.21	99.42	110.60
1	A	172	TYR	CA-CB-CG	5.74	124.31	113.40
1	A	297	ARG	NE-CZ-NH1	5.62	123.11	120.30
1	A	255	ARG	NE-CZ-NH1	5.34	122.97	120.30
1	A	551	ARG	NE-CZ-NH2	-5.17	117.71	120.30
1	A	173	GLN	N-CA-C	-5.05	97.36	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	10	VAL	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4774	0	4686	59	0
2	B	28	0	25	0	0
3	A	43	0	30	3	0
4	A	42	0	39	0	0
5	A	8	0	0	0	0
6	A	4	0	0	0	0
7	A	1	0	0	0	0
8	A	6	0	0	1	0
9	A	290	0	0	1	0
All	All	5196	0	4780	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:171:PRO:O	1:A:172:TYR:HB3	1.43	1.14
1:A:169:THR:HG23	1:A:170:PRO:HD3	1.17	1.10
1:A:169:THR:CG2	1:A:170:PRO:HD3	1.84	1.07
1:A:55:LEU:HD22	1:A:175:LEU:O	1.62	0.99
1:A:167:CYS:HB2	1:A:168:PRO:HD2	1.47	0.96
1:A:171:PRO:O	1:A:172:TYR:CB	2.12	0.94
1:A:170:PRO:HB2	1:A:171:PRO:HD3	1.50	0.92
1:A:170:PRO:CB	1:A:171:PRO:HD3	1.99	0.91
1:A:172:TYR:CD1	1:A:173:GLN:O	2.29	0.86
1:A:169:THR:H	1:A:170:PRO:HD2	1.38	0.85
1:A:170:PRO:HG2	1:A:171:PRO:CD	2.08	0.84
1:A:169:THR:HG23	1:A:170:PRO:CD	2.07	0.83
1:A:168:PRO:HG2	1:A:168:PRO:O	1.78	0.83
1:A:170:PRO:HG2	1:A:171:PRO:HD3	1.60	0.83

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:170:PRO:CG	1:A:171:PRO:HD3	2.11	0.81
1:A:167:CYS:HB2	1:A:168:PRO:CD	2.13	0.79
1:A:167:CYS:CB	1:A:168:PRO:CD	2.65	0.75
1:A:351:HIS:HD1	1:A:437:ASN:HD21	1.34	0.73
1:A:168:PRO:HG3	1:A:172:TYR:O	1.87	0.73
1:A:169:THR:H	1:A:170:PRO:CD	2.03	0.72
1:A:167:CYS:CB	1:A:168:PRO:HD2	2.21	0.70
1:A:169:THR:CG2	1:A:170:PRO:CD	2.66	0.70
1:A:55:LEU:CD2	1:A:175:LEU:O	2.41	0.66
1:A:170:PRO:CB	1:A:171:PRO:CD	2.77	0.62
1:A:169:THR:N	1:A:170:PRO:HD2	2.13	0.61
1:A:109:HIS:NE2	8:A:611:MZY:S2	2.67	0.61
1:A:172:TYR:CE1	1:A:173:GLN:O	2.54	0.60
1:A:170:PRO:CG	1:A:171:PRO:CD	2.77	0.59
3:A:601:HEM:HMC2	3:A:601:HEM:HBC2	1.84	0.58
1:A:123:GLU:HG3	1:A:125:SER:H	1.69	0.58
1:A:168:PRO:O	1:A:168:PRO:CG	2.47	0.58
1:A:168:PRO:HG3	1:A:172:TYR:C	2.25	0.56
1:A:170:PRO:HG2	1:A:171:PRO:HD2	1.86	0.56
1:A:55:LEU:HD22	1:A:175:LEU:C	2.25	0.56
1:A:468:GLN:HG2	1:A:474:LYS:HA	1.87	0.55
1:A:408:ASN:HB2	9:A:885:HOH:O	2.08	0.53
1:A:123:GLU:OE1	1:A:123:GLU:HA	2.09	0.52
3:A:601:HEM:HMB2	3:A:601:HEM:HBB2	1.94	0.50
1:A:106:ILE:HD11	1:A:265:ALA:CB	2.43	0.49
1:A:551:ARG:HD3	1:A:584:LYS:HA	1.95	0.49
1:A:213:MET:HG2	1:A:273:HIS:CD2	2.49	0.47
1:A:82:ILE:CD1	1:A:480:LEU:HD13	2.45	0.46
1:A:333:ASN:HD22	1:A:333:ASN:C	2.18	0.46
1:A:150:LYS:HZ2	1:A:154:GLN:HE22	1.65	0.45
1:A:467:LEU:HG	1:A:471:LEU:HD22	1.97	0.45
1:A:523:ARG:HG3	1:A:529:TRP:CE2	2.52	0.45
1:A:170:PRO:HB2	1:A:171:PRO:CD	2.35	0.45
1:A:581:THR:HG22	1:A:581:THR:O	2.17	0.45
3:A:601:HEM:CMB	3:A:601:HEM:HBB2	2.47	0.44
1:A:119:LEU:N	1:A:119:LEU:HD13	2.33	0.44
1:A:167:CYS:CB	1:A:168:PRO:HD3	2.48	0.44
1:A:169:THR:N	1:A:170:PRO:CD	2.67	0.43
1:A:221:ASP:HB2	1:A:226:TYR:CZ	2.53	0.43
1:A:169:THR:HG22	1:A:170:PRO:HD3	1.90	0.43
1:A:196:GLU:HB3	1:A:198:SEP:O2P	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:407:MET:HB3	1:A:501:MET:CE	2.49	0.43
1:A:150:LYS:NZ	1:A:154:GLN:HE22	2.17	0.43
1:A:175:LEU:HB3	1:A:176:ALA:H	1.65	0.42
1:A:123:GLU:HG3	1:A:125:SER:N	2.33	0.41
1:A:588:SER:N	1:A:589:PRO:CD	2.84	0.41
1:A:227:LEU:HD23	1:A:270:LEU:HD22	2.02	0.41
1:A:295:GLU:O	1:A:299:ILE:HD12	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	592/595 (100%)	562 (95%)	22 (4%)	8 (1%)	11 11

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	8	ALA
1	A	10	VAL
1	A	11	PRO
1	A	167	CYS
1	A	168	PRO
1	A	170	PRO
1	A	209	PRO
1	A	169	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	517/517 (100%)	488 (94%)	29 (6%)	21	29

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	CYS
1	A	57	LEU
1	A	91	VAL
1	A	98	LEU
1	A	106	ILE
1	A	119	LEU
1	A	153	THR
1	A	173	GLN
1	A	175	LEU
1	A	187	LEU
1	A	203	LEU
1	A	261	LEU
1	A	268	LEU
1	A	276	LEU
1	A	282	LYS
1	A	317	LEU
1	A	333	ASN
1	A	347	PHE
1	A	360	ARG
1	A	376	LEU
1	A	388	LYS
1	A	464	LEU
1	A	471	LEU
1	A	480	LEU
1	A	511	LEU
1	A	522	ILE
1	A	539	LYS
1	A	542	ASP
1	A	564	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	154	GLN
1	A	333	ASN
1	A	423	GLN
1	A	468	GLN
1	A	497	ASN
1	A	520	GLN
1	A	570	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	SEP	A	198	1	8,9,10	1.04	0	8,12,14	3.23	3 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	198	1	-	5/5/8/10	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	198	SEP	OG-CB-CA	7.25	115.20	108.14
1	A	198	SEP	P-OG-CB	3.63	128.28	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	198	SEP	O2P-P-OG	-3.21	98.19	106.73

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	198	SEP	CB-OG-P-O2P
1	A	198	SEP	CB-OG-P-O3P
1	A	198	SEP	CB-OG-P-O1P
1	A	198	SEP	CA-CB-OG-P
1	A	198	SEP	N-CA-CB-OG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	198	SEP	1	0

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.62	0	17,19,21	1.32	4 (23%)
2	NAG	B	2	2	14,14,15	0.76	0	17,19,21	2.68	8 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	2	NAG	C1-O5-C5	7.80	122.76	112.19
2	B	2	NAG	C1-C2-N2	-3.29	104.87	110.49
2	B	2	NAG	C4-C3-C2	3.22	115.73	111.02
2	B	2	NAG	O5-C1-C2	2.97	115.98	111.29
2	B	1	NAG	C1-C2-N2	2.90	115.44	110.49
2	B	2	NAG	O3-C3-C4	-2.83	103.81	110.35
2	B	2	NAG	C3-C4-C5	2.80	115.24	110.24
2	B	2	NAG	O5-C5-C4	2.39	116.63	110.83
2	B	1	NAG	O7-C7-N2	2.25	126.09	121.95
2	B	1	NAG	O5-C1-C2	-2.17	107.87	111.29
2	B	1	NAG	O7-C7-C8	-2.13	118.10	122.06
2	B	2	NAG	C6-C5-C4	-2.08	108.12	113.00


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	605	1	14,14,15	0.89	1 (7%)	17,19,21	2.39	9 (52%)
5	NO3	A	607	-	1,3,3	0.17	0	0,3,3	0.00	-
4	NAG	A	606	1	14,14,15	0.77	0	17,19,21	2.33	6 (35%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HEM	A	601	1,8	27,50,50	1.31	4 (14%)	17,82,82	1.48	2 (11%)
4	NAG	A	602	1	14,14,15	0.84	0	17,19,21	1.77	5 (29%)
8	MZY	A	611	3	4,6,6	2.86	1 (25%)	0,7,7	0.00	-
5	NO3	A	608	-	1,3,3	1.20	0	0,3,3	0.00	-
6	CO3	A	609	-	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	606	1	-	2/6/23/26	0/1/1/1
4	NAG	A	605	1	-	0/6/23/26	0/1/1/1
8	MZY	A	611	3	-	-	0/1/1/1
4	NAG	A	602	1	-	0/6/23/26	0/1/1/1
3	HEM	A	601	1,8	-	0/6/54/54	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	611	MZY	C2-S2	5.70	1.78	1.66
3	A	601	HEM	C3B-C2B	-3.71	1.35	1.40
3	A	601	HEM	C4B-NB	-2.46	1.31	1.36
3	A	601	HEM	C4D-C3D	2.44	1.48	1.42
4	A	605	NAG	O4-C4	2.18	1.48	1.43
3	A	601	HEM	C1A-CHA	-2.04	1.35	1.41

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	606	NAG	C1-C2-N2	-5.20	101.61	110.49
4	A	605	NAG	O5-C1-C2	-4.54	104.11	111.29
4	A	605	NAG	O3-C3-C2	-4.13	100.92	109.47
3	A	601	HEM	CBA-CAA-C2A	-3.98	105.15	112.49
4	A	606	NAG	C4-C3-C2	3.80	116.59	111.02
4	A	606	NAG	O4-C4-C5	3.64	118.33	109.30
4	A	602	NAG	O4-C4-C5	3.62	118.28	109.30
4	A	605	NAG	O4-C4-C3	3.57	118.59	110.35
4	A	606	NAG	C1-O5-C5	3.49	116.92	112.19
4	A	602	NAG	C3-C4-C5	-3.35	104.27	110.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	606	NAG	O3-C3-C4	-3.09	103.21	110.35
4	A	605	NAG	C3-C4-C5	-3.02	104.84	110.24
4	A	605	NAG	C4-C3-C2	2.90	115.26	111.02
4	A	605	NAG	O7-C7-N2	2.46	126.48	121.95
4	A	602	NAG	C1-C2-N2	-2.35	106.47	110.49
4	A	602	NAG	O5-C5-C6	2.25	110.73	107.20
4	A	602	NAG	C8-C7-N2	-2.21	112.36	116.10
3	A	601	HEM	CMC-C2C-C3C	2.17	128.73	124.68
4	A	605	NAG	C6-C5-C4	-2.09	108.11	113.00
4	A	605	NAG	O6-C6-C5	-2.09	104.14	111.29
4	A	606	NAG	O5-C5-C6	2.07	110.44	107.20
4	A	605	NAG	O4-C4-C5	2.01	114.29	109.30

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	606	NAG	C4-C5-C6-O6
4	A	606	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	601	HEM	3	0
8	A	611	MZY	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	594/595 (99%)	-0.25	32 (5%)	25 32	12, 25, 100, 211	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	174	SER	17.6
1	A	2	TRP	14.5
1	A	173	GLN	12.7
1	A	121	SER	10.7
1	A	1	SER	10.4
1	A	595	ASN	9.5
1	A	175	LEU	9.2
1	A	120	GLY	8.9
1	A	171	PRO	7.7
1	A	4	VAL	7.4
1	A	122	ASN	7.4
1	A	8	ALA	7.1
1	A	170	PRO	6.7
1	A	3	GLU	6.6
1	A	119	LEU	6.5
1	A	172	TYR	6.0
1	A	11	PRO	5.7
1	A	594	GLU	5.6
1	A	5	GLY	5.4
1	A	9	PRO	5.3
1	A	593	ARG	5.2
1	A	7	GLY	4.3
1	A	12	LEU	3.7
1	A	10	VAL	3.6
1	A	13	VAL	3.3
1	A	124	HIS	3.2
1	A	123	GLU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	64	ARG	2.6
1	A	6	CYS	2.5
1	A	118	GLU	2.4
1	A	322	GLN	2.4
1	A	18	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	A	198	10/11	0.92	0.14	24,37,98,98	0

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	B	2	14/15	0.80	0.29	57,71,83,89	0
2	NAG	B	1	14/15	0.96	0.12	29,39,48,57	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NAG	A	606	14/15	0.83	0.23	54,66,74,76	0
4	NAG	A	602	14/15	0.91	0.23	45,55,62,64	0
5	NO3	A	607	4/4	0.93	0.16	44,49,49,53	0
6	CO3	A	609	4/4	0.94	0.12	42,43,46,48	0
4	NAG	A	605	14/15	0.95	0.13	35,37,44,52	0
8	MZY	A	611	6/6	0.97	0.14	38,39,41,42	0
3	HEM	A	601	43/43	0.98	0.08	12,13,14,17	0
5	NO3	A	608	4/4	0.99	0.10	14,17,17,18	0
7	CA	A	610	1/1	1.00	0.05	14,14,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HEM A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.