



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 21, 2020 – 11:08 PM BST

PDB ID : 4GIR
Title : Crystal structure of an enolase family member from vibrio harveyi (efi-target 501692) with homology to mannonate dehydratase, with mg, ethylene glycol and sulfate bound (ordered loops, space group P41212)
Authors : Vetting, M.W.; Toro, R.; Bhosle, R.; Al Obaidi, N.F.; Morisco, L.L.; Wasserman, S.R.; Sojitra, S.; Washington, E.; Scott Glenn, A.; Chowdhury, S.; Evans, B.; Hammonds, J.; Hillerich, B.; Love, J.; Seidel, R.D.; Imker, H.J.; Gerlt, J.A.; Almo, S.C.; Enzyme Function Initiative (EFI)
Deposited on : 2012-08-08
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>5%</div> <div>6%</div> </div> </div>
1	B	422	<div> <div>%</div> <div> <div></div> <div>95%</div> <div></div> <div></div> </div> </div>
1	C	422	<div> <div>%</div> <div> <div></div> <div>88%</div> <div>6%</div> <div>6%</div> </div> </div>
1	D	422	<div> <div>5%</div> <div> <div></div> <div>89%</div> <div>8%</div> <div></div> </div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 14452 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	397	Total	C	N	O	S	0	1	0
			3199	2029	555	599	16			
1	B	411	Total	C	N	O	S	0	2	0
			3321	2104	574	625	18			
1	C	397	Total	C	N	O	S	0	1	0
			3199	2029	555	599	16			
1	D	408	Total	C	N	O	S	0	2	0
			3309	2097	573	621	18			

There are 92 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	MET	-	EXPRESSION TAG	UNP D0X4R4
A	-21	HIS	-	EXPRESSION TAG	UNP D0X4R4
A	-20	HIS	-	EXPRESSION TAG	UNP D0X4R4
A	-19	HIS	-	EXPRESSION TAG	UNP D0X4R4
A	-18	HIS	-	EXPRESSION TAG	UNP D0X4R4
A	-17	HIS	-	EXPRESSION TAG	UNP D0X4R4
A	-16	HIS	-	EXPRESSION TAG	UNP D0X4R4
A	-15	SER	-	EXPRESSION TAG	UNP D0X4R4
A	-14	SER	-	EXPRESSION TAG	UNP D0X4R4
A	-13	GLY	-	EXPRESSION TAG	UNP D0X4R4
A	-12	VAL	-	EXPRESSION TAG	UNP D0X4R4
A	-11	ASP	-	EXPRESSION TAG	UNP D0X4R4
A	-10	LEU	-	EXPRESSION TAG	UNP D0X4R4
A	-9	GLY	-	EXPRESSION TAG	UNP D0X4R4
A	-8	THR	-	EXPRESSION TAG	UNP D0X4R4
A	-7	GLU	-	EXPRESSION TAG	UNP D0X4R4
A	-6	ASN	-	EXPRESSION TAG	UNP D0X4R4
A	-5	LEU	-	EXPRESSION TAG	UNP D0X4R4
A	-4	TYR	-	EXPRESSION TAG	UNP D0X4R4
A	-3	PHE	-	EXPRESSION TAG	UNP D0X4R4
A	-2	GLN	-	EXPRESSION TAG	UNP D0X4R4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	SER	-	EXPRESSION TAG	UNP D0X4R4
A	0	MET	-	EXPRESSION TAG	UNP D0X4R4
B	-22	MET	-	EXPRESSION TAG	UNP D0X4R4
B	-21	HIS	-	EXPRESSION TAG	UNP D0X4R4
B	-20	HIS	-	EXPRESSION TAG	UNP D0X4R4
B	-19	HIS	-	EXPRESSION TAG	UNP D0X4R4
B	-18	HIS	-	EXPRESSION TAG	UNP D0X4R4
B	-17	HIS	-	EXPRESSION TAG	UNP D0X4R4
B	-16	HIS	-	EXPRESSION TAG	UNP D0X4R4
B	-15	SER	-	EXPRESSION TAG	UNP D0X4R4
B	-14	SER	-	EXPRESSION TAG	UNP D0X4R4
B	-13	GLY	-	EXPRESSION TAG	UNP D0X4R4
B	-12	VAL	-	EXPRESSION TAG	UNP D0X4R4
B	-11	ASP	-	EXPRESSION TAG	UNP D0X4R4
B	-10	LEU	-	EXPRESSION TAG	UNP D0X4R4
B	-9	GLY	-	EXPRESSION TAG	UNP D0X4R4
B	-8	THR	-	EXPRESSION TAG	UNP D0X4R4
B	-7	GLU	-	EXPRESSION TAG	UNP D0X4R4
B	-6	ASN	-	EXPRESSION TAG	UNP D0X4R4
B	-5	LEU	-	EXPRESSION TAG	UNP D0X4R4
B	-4	TYR	-	EXPRESSION TAG	UNP D0X4R4
B	-3	PHE	-	EXPRESSION TAG	UNP D0X4R4
B	-2	GLN	-	EXPRESSION TAG	UNP D0X4R4
B	-1	SER	-	EXPRESSION TAG	UNP D0X4R4
B	0	MET	-	EXPRESSION TAG	UNP D0X4R4
C	-22	MET	-	EXPRESSION TAG	UNP D0X4R4
C	-21	HIS	-	EXPRESSION TAG	UNP D0X4R4
C	-20	HIS	-	EXPRESSION TAG	UNP D0X4R4
C	-19	HIS	-	EXPRESSION TAG	UNP D0X4R4
C	-18	HIS	-	EXPRESSION TAG	UNP D0X4R4
C	-17	HIS	-	EXPRESSION TAG	UNP D0X4R4
C	-16	HIS	-	EXPRESSION TAG	UNP D0X4R4
C	-15	SER	-	EXPRESSION TAG	UNP D0X4R4
C	-14	SER	-	EXPRESSION TAG	UNP D0X4R4
C	-13	GLY	-	EXPRESSION TAG	UNP D0X4R4
C	-12	VAL	-	EXPRESSION TAG	UNP D0X4R4
C	-11	ASP	-	EXPRESSION TAG	UNP D0X4R4
C	-10	LEU	-	EXPRESSION TAG	UNP D0X4R4
C	-9	GLY	-	EXPRESSION TAG	UNP D0X4R4
C	-8	THR	-	EXPRESSION TAG	UNP D0X4R4
C	-7	GLU	-	EXPRESSION TAG	UNP D0X4R4
C	-6	ASN	-	EXPRESSION TAG	UNP D0X4R4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-5	LEU	-	EXPRESSION TAG	UNP D0X4R4
C	-4	TYR	-	EXPRESSION TAG	UNP D0X4R4
C	-3	PHE	-	EXPRESSION TAG	UNP D0X4R4
C	-2	GLN	-	EXPRESSION TAG	UNP D0X4R4
C	-1	SER	-	EXPRESSION TAG	UNP D0X4R4
C	0	MET	-	EXPRESSION TAG	UNP D0X4R4
D	-22	MET	-	EXPRESSION TAG	UNP D0X4R4
D	-21	HIS	-	EXPRESSION TAG	UNP D0X4R4
D	-20	HIS	-	EXPRESSION TAG	UNP D0X4R4
D	-19	HIS	-	EXPRESSION TAG	UNP D0X4R4
D	-18	HIS	-	EXPRESSION TAG	UNP D0X4R4
D	-17	HIS	-	EXPRESSION TAG	UNP D0X4R4
D	-16	HIS	-	EXPRESSION TAG	UNP D0X4R4
D	-15	SER	-	EXPRESSION TAG	UNP D0X4R4
D	-14	SER	-	EXPRESSION TAG	UNP D0X4R4
D	-13	GLY	-	EXPRESSION TAG	UNP D0X4R4
D	-12	VAL	-	EXPRESSION TAG	UNP D0X4R4
D	-11	ASP	-	EXPRESSION TAG	UNP D0X4R4
D	-10	LEU	-	EXPRESSION TAG	UNP D0X4R4
D	-9	GLY	-	EXPRESSION TAG	UNP D0X4R4
D	-8	THR	-	EXPRESSION TAG	UNP D0X4R4
D	-7	GLU	-	EXPRESSION TAG	UNP D0X4R4
D	-6	ASN	-	EXPRESSION TAG	UNP D0X4R4
D	-5	LEU	-	EXPRESSION TAG	UNP D0X4R4
D	-4	TYR	-	EXPRESSION TAG	UNP D0X4R4
D	-3	PHE	-	EXPRESSION TAG	UNP D0X4R4
D	-2	GLN	-	EXPRESSION TAG	UNP D0X4R4
D	-1	SER	-	EXPRESSION TAG	UNP D0X4R4
D	0	MET	-	EXPRESSION TAG	UNP D0X4R4

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	4	Total	Cl	0	0
			4	4		
4	A	2	Total	Cl	0	0
			2	2		
4	D	2	Total	Cl	0	0
			2	2		
4	C	1	Total	Cl	0	0
			1	1		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	D	1	Total	C	O	0	0
			4	2	2		
5	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Na	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	341	Total	O	0	4
			345	345		
7	B	353	Total	O	0	2
			355	355		
7	C	322	Total	O	0	3
			325	325		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	D	341	Total	O	0	4
			345	345		

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	146.84Å 146.84Å 169.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.23 – 2.00 42.39 – 2.00	Depositor EDS
% Data completeness (in resolution range)	90.9 (38.23-2.00) 90.9 (42.39-2.00)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 2.00Å)	Xtrriage
Refinement program	PHENIX 1.8_1069	Depositor
R, R_{free}	0.148 , 0.193 0.148 , 0.192	Depositor DCC
R_{free} test set	2827 reflections (2.49%)	wwPDB-VP
Wilson B-factor (Å ²)	18.9	Xtrriage
Anisotropy	0.007	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 57.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14452	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 29.72 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4779e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NA, MG, EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/3282	0.53	0/4463
1	B	0.37	0/3406	0.54	0/4631
1	C	0.35	0/3282	0.55	0/4463
1	D	0.36	0/3393	0.53	0/4609
All	All	0.36	0/13363	0.54	0/18166

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3199	0	3086	13	0
1	B	3321	0	3198	5	0
1	C	3199	0	3086	13	0
1	D	3309	0	3189	18	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	5	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
4	A	2	0	0	0	0
4	B	4	0	0	0	0
4	C	1	0	0	0	0
4	D	2	0	0	0	0
5	A	4	0	6	0	0
5	B	4	0	6	0	0
5	C	4	0	6	0	0
5	D	8	0	12	0	0
6	B	1	0	0	0	0
7	A	345	0	0	4	1
7	B	355	0	0	3	1
7	C	325	0	0	1	0
7	D	345	0	0	1	0
All	All	14452	0	12589	47	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 47 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:ARG:NH2	3:A:402:SO4:O3	2.30	0.65
1:A:240:GLU:OE1	7:A:689:HOH:O	2.14	0.65
1:D:4:ASN:OD1	1:D:30:SER:HB2	1.98	0.64
1:D:371:ARG:NH2	7:D:770:HOH:O	2.33	0.60
1:A:199:GLN:NE2	7:A:778:HOH:O	2.32	0.58

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:804:HOH:O	7:B:819:HOH:O[8_554]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	396/422 (94%)	385 (97%)	11 (3%)	0	100	100
1	B	411/422 (97%)	401 (98%)	10 (2%)	0	100	100
1	C	396/422 (94%)	388 (98%)	8 (2%)	0	100	100
1	D	406/422 (96%)	398 (98%)	8 (2%)	0	100	100
All	All	1609/1688 (95%)	1572 (98%)	37 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	345/367 (94%)	343 (99%)	2 (1%)	86	90
1	B	359/367 (98%)	357 (99%)	2 (1%)	86	90
1	C	345/367 (94%)	341 (99%)	4 (1%)	71	76
1	D	358/367 (98%)	357 (100%)	1 (0%)	92	95
All	All	1407/1468 (96%)	1398 (99%)	9 (1%)	86	90

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	40	PHE
1	D	255	LEU
1	C	255	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	40	PHE
1	C	151	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	168	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 14 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	EDO	B	406	-	3,3,3	0.53	0	2,2,2	0.31	0
3	SO4	B	403	-	4,4,4	0.16	0	6,6,6	0.13	0
5	EDO	C	404	-	3,3,3	0.50	0	2,2,2	0.31	0
3	SO4	D	402	-	4,4,4	0.14	0	6,6,6	0.15	0
5	EDO	D	404	-	3,3,3	0.44	0	2,2,2	0.42	0
5	EDO	A	405	-	3,3,3	0.52	0	2,2,2	0.31	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	C	402	-	4,4,4	0.15	0	6,6,6	0.16	0
3	SO4	A	402	-	4,4,4	0.18	0	6,6,6	0.19	0
5	EDO	D	406	-	3,3,3	0.55	0	2,2,2	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	C	404	-	-	0/1/1/1	-
5	EDO	B	406	-	-	0/1/1/1	-
5	EDO	D	404	-	-	0/1/1/1	-
5	EDO	A	405	-	-	0/1/1/1	-
5	EDO	D	406	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	397/422 (94%)	-0.35	8 (2%) 65 63	10, 16, 34, 67	0
1	B	411/422 (97%)	-0.42	5 (1%) 79 78	10, 15, 34, 64	0
1	C	397/422 (94%)	-0.36	6 (1%) 73 72	11, 17, 35, 63	0
1	D	408/422 (96%)	-0.30	19 (4%) 31 30	9, 16, 41, 88	0
All	All	1613/1688 (95%)	-0.36	38 (2%) 59 57	9, 16, 36, 88	0

The worst 5 of 38 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	-3	PHE	6.1
1	D	-4	TYR	5.9
1	D	-10	LEU	5.8
1	D	161	PRO	5.0
1	D	1	MET	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	CL	A	404	1/1	0.83	0.09	45,45,45,45	0
4	CL	B	404	1/1	0.89	0.09	39,39,39,39	0
4	CL	A	403	1/1	0.89	0.15	52,52,52,52	0
4	CL	B	405	1/1	0.89	0.22	57,57,57,57	0
4	CL	D	403	1/1	0.89	0.07	44,44,44,44	0
4	CL	C	403	1/1	0.92	0.09	45,45,45,45	0
5	EDO	D	406	4/4	0.93	0.13	27,28,29,33	0
5	EDO	B	406	4/4	0.93	0.14	22,27,28,31	0
2	MG	B	401	1/1	0.95	0.06	17,17,17,17	0
5	EDO	C	404	4/4	0.95	0.12	24,30,31,32	0
4	CL	B	407	1/1	0.95	0.30	52,52,52,52	0
5	EDO	D	404	4/4	0.95	0.11	22,25,30,31	0
4	CL	B	408	1/1	0.95	0.19	53,53,53,53	0
4	CL	D	405	1/1	0.96	0.26	51,51,51,51	0
3	SO4	C	402	5/5	0.97	0.10	35,38,44,47	0
5	EDO	A	405	4/4	0.97	0.12	18,28,30,31	0
3	SO4	B	403	5/5	0.98	0.11	36,36,42,45	0
3	SO4	D	402	5/5	0.98	0.09	38,38,45,45	0
2	MG	D	401	1/1	0.98	0.03	21,21,21,21	0
2	MG	A	401	1/1	0.98	0.04	16,16,16,16	0
3	SO4	A	402	5/5	0.98	0.08	38,38,42,45	0
6	NA	B	402	1/1	0.99	0.04	14,14,14,14	0
2	MG	C	401	1/1	0.99	0.04	19,19,19,19	0

6.5 Other polymers

There are no such residues in this entry.