



# Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2020 – 10:02 pm BST

PDB ID : 3GK7  
Title : Crystal structure of 4-hydroxybutyrate CoA-Transferase from Clostridium aminobutyricum  
Authors : Messerschmidt, A.; Macieira, S.; Velarde, M.  
Deposited on : 2009-03-10  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	448	<div> <div></div> <div>92%</div> <div>8%</div> </div>
1	B	448	<div> <div>2%</div> <div>93%</div> <div>7%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7282 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 4-Hydroxybutyrate CoA-transferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	447	Total	C	N	O	S	0	0	0
			3402	2164	567	647	24			
1	B	447	Total	C	N	O	S	0	0	0
			3410	2170	569	647	24			

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	439	SER	-	EXPRESSION TAG	UNP Q9RM86
A	440	ALA	-	EXPRESSION TAG	UNP Q9RM86
A	441	TRP	-	EXPRESSION TAG	UNP Q9RM86
A	442	SER	-	EXPRESSION TAG	UNP Q9RM86
A	443	HIS	-	EXPRESSION TAG	UNP Q9RM86
A	444	PRO	-	EXPRESSION TAG	UNP Q9RM86
A	445	GLN	-	EXPRESSION TAG	UNP Q9RM86
A	446	PHE	-	EXPRESSION TAG	UNP Q9RM86
A	447	GLU	-	EXPRESSION TAG	UNP Q9RM86
A	448	LYS	-	EXPRESSION TAG	UNP Q9RM86
B	439	SER	-	EXPRESSION TAG	UNP Q9RM86
B	440	ALA	-	EXPRESSION TAG	UNP Q9RM86
B	441	TRP	-	EXPRESSION TAG	UNP Q9RM86
B	442	SER	-	EXPRESSION TAG	UNP Q9RM86
B	443	HIS	-	EXPRESSION TAG	UNP Q9RM86
B	444	PRO	-	EXPRESSION TAG	UNP Q9RM86
B	445	GLN	-	EXPRESSION TAG	UNP Q9RM86
B	446	PHE	-	EXPRESSION TAG	UNP Q9RM86
B	447	GLU	-	EXPRESSION TAG	UNP Q9RM86
B	448	LYS	-	EXPRESSION TAG	UNP Q9RM86

- Molecule 2 is SPERMIDINE (three-letter code: SPD) (formula: C<sub>7</sub>H<sub>19</sub>N<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			10	7	3		
2	B	1	Total	C	N	0	0
			10	7	3		

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula:  $C_4H_{12}NO_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			8	4	1	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	205	Total	O	0	0
			205	205		
4	B	229	Total	O	0	0
			229	229		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4-Hydroxybutyrate CoA-transferase



- Molecule 1: 4-Hydroxybutyrate CoA-transferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	116.59Å 116.59Å 174.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.85 48.46 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.6 (15.00-1.85) 98.6 (48.46-1.85)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.97 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.186 , 0.211 0.189 , 0.213	Depositor DCC
$R_{free}$ test set	5039 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.4	Xtriage
Anisotropy	0.327	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 55.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7282	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.67 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.4341e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, SPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/3476	0.58	0/4706
1	B	0.49	0/3484	0.58	0/4714
All	All	0.49	0/6960	0.58	0/9420

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3402	0	3354	16	0
1	B	3410	0	3376	20	0
2	A	10	0	19	0	0
2	B	10	0	19	0	0
3	A	8	0	12	0	0
3	B	8	0	12	0	0
4	A	205	0	0	0	0
4	B	229	0	0	3	0
All	All	7282	0	6792	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.



All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:308:CYS:HB2	4:B:676:HOH:O	1.19	1.29
1:B:154:GLU:OE2	1:B:181:HIS:HE1	1.59	0.85
1:B:186:ILE:HG21	1:B:271:MET:HE3	1.64	0.80
1:A:61:LEU:HD21	1:A:271:MET:CE	2.11	0.80
1:B:308:CYS:CB	4:B:676:HOH:O	1.94	0.72
1:A:205:LEU:O	1:A:353:LYS:HE3	1.92	0.70
1:A:154:GLU:OE2	1:A:181:HIS:HE1	1.77	0.68
1:B:296:HIS:HD2	1:B:298:SER:H	1.42	0.66
1:B:115:HIS:HE1	1:B:149:LYS:HE3	1.63	0.64
1:A:296:HIS:HD2	1:A:298:SER:H	1.44	0.63
1:B:154:GLU:OE2	1:B:181:HIS:CE1	2.50	0.60
1:A:83:SER:HB2	1:A:84:PRO:CD	2.33	0.58
1:B:308:CYS:SG	4:B:676:HOH:O	2.56	0.57
1:B:186:ILE:HG21	1:B:271:MET:CE	2.33	0.57
1:B:115:HIS:CE1	1:B:149:LYS:HE3	2.41	0.55
1:A:61:LEU:HD21	1:A:271:MET:HE2	1.88	0.53
1:A:237:SER:O	1:A:267:ILE:HA	2.09	0.52
1:B:186:ILE:HD13	1:B:271:MET:HE2	1.96	0.48
1:A:83:SER:HB2	1:A:84:PRO:HD2	1.95	0.48
1:B:237:SER:O	1:B:267:ILE:HA	2.14	0.47
1:B:83:SER:HB2	1:B:84:PRO:HD2	1.97	0.47
1:A:316:ASP:HA	1:A:374:VAL:O	2.15	0.46
1:B:83:SER:HB2	1:B:84:PRO:CD	2.45	0.46
1:A:247:LEU:HG	1:A:252:VAL:HB	1.98	0.46
1:A:181:HIS:HD2	1:A:182:PRO:O	1.99	0.46
1:B:316:ASP:HA	1:B:374:VAL:O	2.16	0.45
1:B:247:LEU:HG	1:B:252:VAL:HB	1.99	0.45
1:B:205:LEU:O	1:B:353:LYS:HE3	2.18	0.44
1:A:267:ILE:HG13	1:A:288:LEU:HD23	1.98	0.44
1:B:258:LYS:HE2	1:B:258:LYS:HB3	1.88	0.42
1:B:321:ILE:HB	1:B:385:THR:HB	2.02	0.42
1:A:238:GLU:OE1	1:A:336:GLY:HA3	2.20	0.41
1:A:363:LYS:HE2	1:A:367:GLY:O	2.20	0.41
1:A:35:GLU:O	1:A:37:PRO:HD3	2.21	0.41
1:A:246:ASP:OD1	1:A:275:ARG:NH2	2.50	0.41
1:B:300:VAL:HB	1:B:343:GLY:HA3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	445/448 (99%)	433 (97%)	11 (2%)	1 (0%)	47	33
1	B	445/448 (99%)	434 (98%)	10 (2%)	1 (0%)	47	33
All	All	890/896 (99%)	867 (97%)	21 (2%)	2 (0%)	47	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	139	ASP
1	B	139	ASP

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	370/376 (98%)	362 (98%)	8 (2%)	52	36
1	B	372/376 (99%)	365 (98%)	7 (2%)	57	43
All	All	742/752 (99%)	727 (98%)	15 (2%)	55	40

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	LEU
1	A	65	GLU
1	A	102	PHE
1	A	214	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	227	LEU
1	A	230	LYS
1	A	233	LEU
1	A	285	LYS
1	B	39	LEU
1	B	65	GLU
1	B	102	PHE
1	B	214	LEU
1	B	227	LEU
1	B	233	LEU
1	B	444	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	46	ASN
1	A	74	ASN
1	A	181	HIS
1	A	296	HIS
1	B	46	ASN
1	B	74	ASN
1	B	181	HIS
1	B	296	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SPD	B	501	-	9,9,9	0.32	0	8,8,8	0.67	0
2	SPD	A	501	-	9,9,9	0.33	0	8,8,8	0.67	0
3	TRS	A	502	-	7,7,7	0.28	0	9,9,9	0.64	0
3	TRS	B	502	-	7,7,7	0.19	0	9,9,9	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SPD	B	501	-	-	4/7/7/7	-
2	SPD	A	501	-	-	6/7/7/7	-
3	TRS	A	502	-	-	0/9/9/9	-
3	TRS	B	502	-	-	0/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	501	SPD	C3-C4-C5-N6
2	A	501	SPD	C3-C4-C5-N6
2	A	501	SPD	C8-C7-N6-C5
2	B	501	SPD	C7-C8-C9-N10
2	A	501	SPD	C7-C8-C9-N10
2	A	501	SPD	C4-C5-N6-C7
2	B	501	SPD	C4-C5-N6-C7
2	A	501	SPD	C2-C3-C4-C5
2	B	501	SPD	N1-C2-C3-C4
2	A	501	SPD	N1-C2-C3-C4

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	447/448 (99%)	-0.14	6 (1%) 77 78	10, 17, 29, 41	0
1	B	447/448 (99%)	-0.10	11 (2%) 57 56	10, 17, 29, 46	0
All	All	894/896 (99%)	-0.12	17 (1%) 66 66	10, 17, 29, 46	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	441	TRP	8.7
1	B	440	ALA	7.4
1	A	441	TRP	6.7
1	B	1	MET	6.2
1	A	1	MET	5.0
1	B	442	SER	3.5
1	B	444	PRO	3.3
1	B	2	ASP	3.3
1	B	446	PHE	2.9
1	A	2	ASP	2.6
1	B	447	GLU	2.5
1	B	9	ASP	2.5
1	A	228	LYS	2.4
1	B	228	LYS	2.1
1	A	229	ASP	2.1
1	B	229	ASP	2.1
1	A	9	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SPD	B	501	10/10	0.59	0.30	47,50,51,51	0
2	SPD	A	501	10/10	0.76	0.23	34,43,47,47	0
3	TRS	B	502	8/8	0.95	0.11	13,14,15,15	0
3	TRS	A	502	8/8	0.96	0.10	13,14,14,14	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.